Vol. 94, Sect. 15, 1957

PISCES

COMPILED BY

G. PALMER (Recent) and E. I. WHITE, D.Sc., F.R.S. (Fossil)

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE EIGHT SHILLINGS AND SIXPENCE

March, 1959

SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The following are the Sections and their prices, viz .:-

						s.	d.
1.	Comprehens	sive Zo	ology		 	 3	0
2.	Protozoa				 	 10	0
3.	Porifera				 	 2	0
4.	Coelenterata	a			 	 4	0
5.	Echinoderm	a			 	 4	0
6.	Vermes				 	 10	0
7.	Brachiopoda	a			 	 4	0
8.	Bryozoa				 	 2	6
9.	Mollusca				 	 10	0
10.	Crustacea				 	 6	6
11.	Trilobita				 	 5	0
12.	Arachnida				 	 10	0
*13.	Insecta				 	 50	0
14.	Protochorda	ıta			 	 2	0
15.	Pisces				 	 8	6
16A	. Amphibia				 	 6	0
	. Reptilia				 	 79	6
	Aves				 	 10	0
18.	Mammalia				 	 10	0
19.	List of Nev					4	0
				0			

^{*} Obtainable only from the Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7.

15. PISCES

COMPILED BY

G. PALMER (Recent) and

E. I. WHITE, D.Sc., F.R.S. (Fossil)

CONTENTS

						PAGE	
I.	TITLES	• • •	 			•••	1
II.	SUBJECT INDEX	•••	 	•••	•••		48
III.	Systematic Index		 				59

I.—TITLES.

The year of publication of a title is omitted from the reference where it is the same as the volume year of the 'Record' namely 1957.

Direction 30. Determination of the gender to be attributed to the names of fifty-seven genera of fishes, cyclostomes, reptiles and amphibia placed on the Official List of Generic Names in Zoology in the period up to the end of 1936. Opin. int. Comm. zool. Nom. 1 C 1955: 287–298.

Direction 33. Addition to the Official List of Generic Names in Zoology of the generic name Pleuronectes Linnaeus, 1758 (Class Actinopterygii) (Direction supplementary to Opinion 68). Opin. int. Comm. zool. Nom. 1 C 1956: 329-340.

Direction 34. Addition to the Official List of Generic Names in Zoology of the generic name Sparus Linnaeus, 1758 (Class Actinopterygii) (Direction supplementary to Opinion 69). Opin. int. Comm. zool. Nom. 1 C 1956: 341-

© Zoological Society of London, 1959.

Direction 56. Completion and in certain cases correction of entries relating to the names of genera belonging to the Classes Pisces, Amphibia and Reptilia made in the Official List of Generic Names in Zoology in the period up to the end of 1936. Opin. int. Comm. zool. Nom. 1 D 1956: 337-364.

Direction 57. Addition to the Official List of Specific Names in Zoology (a) of the specific names of forty-seven species belonging to the Classes Cyclostomata, Pisces, Amphibia and Reptilia, each of which is the type species of a genus, the name of which was placed on the Official List of Generic Names in Zoology in the period up to the end of 1936 and (b) of the specific name of one species of the Class Amphibia which is currently treated as a senior subjective synonym of the name of such a species. Opin. int. Comm. zool. Nom. 1 D 1956: 365-388.

Direction 69. Validation under the Plenary Powers of the generic name Osmerus as a subgeneric status as from Linnaeus, 1758 (Class Pisces) (validation of an error in the Ruling given in Opinion 77). Opin. int. Comm. zool. Nom. 1 E: 137-150.

Opinion 375. Designation under the plenary powers of a type species in harmony with current usage for the genus *Heterandria* Agassiz (J.L.R.), 1853 (Class Osteichthyes). Opin. int. Comm. zool. Nom. 11 1955: 379–390.

Opinion 438. Validation under the Plenary Powers of the generic name Rhinopteraspis Jaekel, 1919 (Class Ostracodermi). Opin. int. Comm. zool. Nom. 15: 41-50.

Opinion 461. Addition to the Official List of Generic Names in Zoology of the generic name Cubiceps Lowe (R.T.), 1843, a name published conditionally for a species described simultaneously in the genus Seriola Cuvier (G.L.C.F.D.), [1816] (Class Pisces) (Opinion supplementary to Declaration 30). Opin. int. Comm. zool. Nom. 15: 475–484.

Opinion 485. Determination of the relative priority to be assigned to the names Cheirodon axelrodi Schultz (L.P.) and Hyphessobrycon cardinalis Myers (G.S.) and Weitzman (S.H.) (Class Pisces) both being names published in February, 1956. Opin. int. Comm. zool. Nom. 17:87-104.

Opinion 487. Validation under the Plenary Powers of the generic names Gempylus Cuvier, 1829 (Class Pisces) and Acinaces Gerstaecker, 1858 (Class Insecta, Order Coleoptera). Opin. int. Comm. zool. Nom. 17: 119-142.

Anon (1). Mistaken journey. Angling Times No. 207:1.

Anon (2). Mystery of the Red Tide. Sea Frontiers 3: 21-31 6 figs.

(Abakumov, V. A.). [The mode of life of the Baltic migratory lamprey]. Ichth. Probl., Moscow 6 1956: 122–128. [In Russian.]

(Abdullaev, M. A.). [Fishes of Kuyu-Mazar water reservoir]. Ichth. Probl., Moscow 8:107-115 4 graphs. [In Russian].

Abe, T. (1). New, rare or uncommon fishes from Japanese waters. VI. Notes on the rare fishes of the family Histiopteridae. Jap. J. Ichthyol. 6:35–40 and 71–74 1 fig.

Abe, T. (2). Studies on the fishes whose young are believed to be carried in large numbers by the "Kuro-shiwo." Rec. oceanogr. Wks. Japan (Spl. No.): 93–96.

Abe, T. (3). Notes on fishes from the stomachs of whales taken in the Antarctic. I. Xenocyttus nemotoi, a new genus and new species of Zeomorph fish of the subfamily Oreosominae Goode and Bean, 1895. Sci. Rep. Whale Res. Inst. Tokyo No. 12: 225–233 2 pls.

Abe, T. see Tomiyama, I.

Abolins, L. & Abolins-Krogis, A. Studies on skin carotenoids of *Crenilabrus pavo* C.V. Pubbl. Staz. zool. Napoli **29**: 389–406 4 figs.

Abolins-Krogis, A. see Abolins, L.

(Abrosov, V. N. & Agapov, I. D.). [On the autoregulation of the number of smelt in Zhizhitzk lake]. Ichth. Probl., Moscow 8:160-178 fig. [In Russian.]

Acolat, L. Recherches relatives à la reaction du coeur isolé du Brochet (Esox lucius L.) au passage du courant galvanique. Ann. Sci. Univ. Besançon (2) Zool. & Phys. No. 4 1955: 9–16 6 figs.

Acolat, L. & Muller, T. Essais en vue de provoquer artificiellement des formations pathologiques au niveau des téguments chez quelques Poissons d'eau douce. Ann. Sci. Univ. Besançon (2) Zool. & Phys. No. 5 1956: 3–12.

Adam, H. Der III Ventrikel und die mikroskopische Struktur seiner Wände bei Lampetra (Petromyzon) fluviatilis L. und Myxine glutinosa L., nebst einigen Bemerkungen über des Infundibularorgan von Branchiostoma (Amphioxus) lanceolatum Pal. Progress in Neurobiology, London, N.Y. 1956: 146–158 6 figs.

Afzelius, B. A. & Olsson, R. The fine structure of the subcommissural cells and of Reissner's fibre in *Myxine*. Z. Zellforsch. 46: 672–685.

Agapov, I. D. see Abrosov, V. N.

Ahmad, N. Mortality of fish during transport and cultural operation: Causes and prevention. Pakistan J. Sci. 6 1954:102-106.

Alabaster, J. G., Herbert, D. W. M. & Hemens, J. The survival of the rainbow trout (Salmo gairdnerii Richardson) and perch (Perca fluviatilis L.) at various concentrations of dissolved oxygen and carbon dioxide. Ann. appl. Biol. 45: 177-188 6 figs.

Albuquerque, R. M. Peixes de Portugal e ilhas adjacentes chaves para a sua determinação. Portug. acta biol. 5B 1956: i-xvi, 1-1167 445 figs.

Alderdice, D. F. & Brett, J. R. Some effects of Kraft mill effluent on young Pacific salmon. J. Fish. Res. Bd. Can., 14:783-795.

(Aleev, Y. G.) (1). [The role played by low temperatures in the stimulation of trophoplasmatic growth of oocytes in fish]. C.R. Acad. Sci. URSS 110 1956: 491-493 Pl. [In Russian.]

(Aleev, Y. G.) (2). [On the functional importance of lateral (horizontal) body orientation in Pleuronectiformes]. C.R. Acad. Sci. URSS 110 1956: 707–709 3 figs. [In Russian.]

(Aleev, Y. G.) (3). [Some principles of fish growth.] Ichth. Probl., Moscow 6 1956: 75-95. [In Russian.]

(Aleev, Y. G.) (4). [The systematic position of *Trachurus* in the Black Sea.] Ichth. Probl., Moscow 7 1956: 174–184 2 figs. [In Russian.]

(Aleev, Y. G.) (5). [Function of fins and their characteristics.] Ichth. Probl., Moscow 8:55-76 9 figs. [In Russian.]

(Aleev, Y. G.) (6). [On ten species of Trachurus in the U.S.S.R. seas]. Trav. Sta. biol. Sebastopol 9:167-242 figs. pls. [In Russian.]

(Aleev, Y. G.) (7). [On the evolution of the pelagic Carangidae (Carangidae, Perciformes).] Trav. Sta. biol. Sebastopol 9:274–281 4 figs. [In Russian.]

Alexander, K. M. A comparative study of the total protein content in the skeletal muscles of some representative vertebrates. J. zool. Soc. India 8 1956: 149-156.

Alfert, M. Chemical differentiation of nuclear proteins during spermatogenesis in the salmon. J. biophys. & biochem. Cytol. 2 1956: 107-1143 pls.

Allison, L. N. Variation in strength of pyridylmercuric acetate technical and its effect on rainbow trout. Prog. Fish. Cult. 19:108-111 2 figs.

Alm, G. [The yield of the pike and perch fisheries along the Baltic coasts of Sweden during the years 1914–1955.] Rep. Inst. Freshwater Res. Drottningholm, No. 38, : 5–69. 24 figs. [English Summary.]

Altamirans, M. Electrical properties of the innervated membrane of the electroplax of electric eel. J. cell. comp. Physiol. 46 1955: 249-277 9 figs.

Alvariño, A. (1). Estudio del zooplancton recogido en la campaña "Vendaval", en Terranova, Marzo, abril y mayo de 1953. Bol. Inst. esp. Oceanogr. No. 76 1956: 1-28.

Alvariño, A. (2). Zooplancton de Terranova (Febrero, marzo y junio de 1955). Bol. Inst. esp. Oceanogr. No. 77 1956: 1–18.

Alvariño, A. (3). Estudio del zooplancton del Mediterraneo Occidental. Bol. Inst. esp. Oceanogr. No. 81: 1-26 9 figs.

Alvariño, A. (4). Zooplancton del Atlantico Iberico. Bol. Inst. esp. Oceanogr. No. 82: 1–51 7 figs.

Alverson, D. L. & Chatwin, B. M. Results from tagging experiments on a spawning stock of petrale sole, *Eopsetta jordani* (Lockington). J. Fish. Res. Bd. Can. 14:953-974 3 figs.

Alverson, D. L. see Welander, A. D.

Alverson, F. G. & Shimada, B. M. A study of the Eastern Pacific Fishery for tuna baitfishes, with particular reference to the anchoveta (*Cetengraulis mysticetus*). Bull. Inter-Amer. Tuna Comm. 2:25-79.

Aminul Huq, A. K. M. see Bhatti, H. K.

(Amosov, V. A.) [Fundamental indicators for external features of fish,] Iehth. Probl., Moscow 6 1956; 55-74 10 figs. [In Russian.]

Anadón, E. Anatomia e histologia de las placas luminosas caudales de Lampadena nitida (Taaning). Bol. Soc. esp. Hist. nat. Secc. Biol. 55: 129-144 6 pls.

Ancona, U. D'. (1). Inversions spontanées et expérimentelles dans les gonades des téléostéens. Année Biol. (3) 32 1956: 89-99.

Ancona, U. d' (2). The gonad differentiation in lower teleostomes. Int. Congr. Zool. 14 1953 (1956): 237-239.

Ancona, U. d' (3). Morphogénèse et differenciation sexuelle chez les poissons téléostéens. Bull. Soc. zool. Fr. 81: 219–229 2 pls.

Ancona, U. d' (4). La base génétique et l'influence du milieu sur la différenciation sexuelle de l'Anguille. C.R. Acad. Sci. Paris 245: 23–264.

Ancona, U. d' (5). Nuove ricerche sperimentali sull'azione di ormoni steroidi sulla gonade dell' anguilla. Pubbl. Staz. zool. Napoli 29:307-322 5 pls. [English summary.]

Andréka, F. see Gyurko, S.

(Andriyashev, A. P.). [The role of sense organs in searching for food by fish.] Trud. Sovesh. Ikhtiol. Komm. 6 1954 (1955):135-142. [In Russian.]

(Anpilova, V. I.). [On the taxonomic position of Coregonus sardinella baunti Muchomedijarov, 1948.] C.R. Acad. Sci. URSS 111 1956: 898-900 fig. [In Russian.]

Aoyagi, H. One new species of the Pomacentridae, Pisces, from the Palau Islands. Zool. Mag., Tokyo 53, 1941: 180-181 1 fig.

Aoyama, T. [On the regeneration of the teleost scale, with special reference to the yellow sea bream, *Taius tumifrons*.] Bull. Jap. Soc. Sci. Fish. 22: 679–684 4 figs. [English summary.]

Arachelian, I. see Busnita, T.

Aramburu, A. S. A. de (1). Nota sobre *Cyrtocharax squamosus* (Eigenmann y Kennedy). (Pisces: Characoidei). Notas Mus. Univ. nac. La Plata, Zool. 19:87-92.

Aramburu, A. S. A. de. (2). Una nueva cita de *Mola mola* (Linné) para el atlantico sur (Pisces: Plectognathi). Notas Mus. Univ. nac. La Plata, Zool. 19 93–99 5 figs.

Aramburu, A. S. A. de. (3). Porotergus ellisi una nueva especie de gimnotido de la Argentina (Pisces: Gymnotoidei). Notas Mus. Univ. nac. La Plata, Zool. 19:153-159 4 figs.

Aramburu, A. S. A. de. (4). Nota sobre Auchenipterus nuchalis (Spix) (Pisces: Siluroidei). Notas Mus. Univ. nac. La Plata Zool. 19: 171–176 3 figs.

Arbocco, G. Contributo alla conoscenza dei pesci d'acqua dolce della Liguria. 1. Le trotte del Rio Baracca Ann. Mus. Civico Stor. nat. Genova 68 1955: 143-148 pls. iii and iv.

Arnold, D. C. Notes on the fishes of the family Carapidae (Fierasferidae) in the Museum of the Institut Océanographique, Monaco. Bull. Inst. océanogr. Monaco No. 1084 1956: 1-5.

Arnoult, J. Sur quelques poissons rare et peu connus des eaux douces de France. Bull. Mus. Hist. nat. Paris (2) 29:464-466.

Aronson, L. R. see Brown, M. E.

Arora, H. L. & Sperry, R. W. Myotypic respecification of regenerated nerve fibres in cichlid fishes. J. Embryol. Exp. Morph. 5:256-263 2 figs.

Arvy, L., Fontaine, M. & Gabe, M. Fonction thyroidienne et complexe hypothalamo-hypophysaire chez la truite. C.R. Soc. Biol. Paris 150 1956: 625-627.

Ashby, K. R. (1). The early development of the reproductive system of the brown trout *Salmo trutta* L. Int. Congr. Zool. 14 1953 (1956): 239–242.

Ashby, K. R. (2). The effect of steroid hormones on the brown trout (Salmo trutta L.) during the period of gonadal differentiation. J. Embryol. Exp. Morph. 5: 225-249 2 pl. 4 figs.

(Aslanova, N. E.). [Behaviour of pelagic fish in Black Sea during the winter.] Trud. Sovesh. Ikhtiol. Komm. 5 1953 (1955): 144-153 8 figs. [In Russian.]

Atz, E. H. and Pickford, G. E. Failure to elicit the Galli-Mainini reaction in Rana pipiens with spawning reflex fractions and other teleostean pituitary preparations, and observations on the response to mammalian gonadotrophins. Zoologica, N.Y. 39 1954:117-122.

Atz, J. W. & Coates, C. W. A remarkable instance of parallelism among Teleost fishes. Copeia 1957: 302-303.

Austin, P. Notes on the growth of

Austin, P. Notes on the growth of Irish pike. Fishing Gazette 139:420.

Avnimelech, M. Découverte de Stratodus (Teleostei: Dercetidae) dans le Sénonien supérieur d'Israël. C.R. Soc. géol. Fr. 1957: 23–24.

Avondhu'. Notes on some large Irish salmon and brown trout. Salm-Trout Mag. No. 150: 156-161.

Axelrod, H. R. & Vorderwinkler, W. Encyclopedia of tropical fishes, with special emphasis on techniques of breeding. New York (Sterling Publ. Co.) pp. 733, text illust. (some col.).

Azazaki, M. [Biological studies on a dragonet, Synchiropus altivelis (Temminck et Schlegel).] Jap. J. Ichthyol. 5:146-152 2 figs. [English summary.]

(Azbelev, V. V. & Lagunov, I. I.). [Some data regarding the migrations of semga salmon.] Ichth. Probl., Moscow 6 1956: 113-120. [In Russian.]

(Baburina, E. A.). [Peculiarities in structure and function of the fish eye.] Trud. Sovesh. Ikhtiol. Komm. 5 1953 [1955]: 90–103 12 figs. [In Russian.]

(Baburina, E. A. & Buznikov, G. A.). [The hatching glands of Kuban embryos of *Vimba vimba carinata* as a source of hyaluronidase and of the hatching enzyme.] C.R. Acad. Sci. URSS 113: 1387–1390 2 figs. [In Russian.]

Backus, R. H. (1). The fishes of Labrador. Bull. Amer. Mus. nat. Hist. 113: 273-332 2 pls. 2 figs.

Backus, R. H. (2). Notes on Western North Atlantic sharks. Copeia 1957: 246-248.

Backus, R. H. see Gordon, M. S. Backus, R. H. see Johnson, H. R.

Bader, H. Erfahrungen mit der Netzschmerle Botia lohachata Chaudhuri. Aquar. Terrar. Z. 10: 203–205.

Baerends, G. P. see Brown, M. E.

Bagenal, T. B. (1). The breeding and fecundity of the long rough dab *Hippoglossoides platessoides* (Fabr.) and the associated cycle in condition. J. Mar. biol. Ass. U.K. 36:339-375 10 figs.

Bagenal, T. B. (2). Annual variations in fish fecundity. J. Mar. biol. Ass. U.K. 36: 377–382.

Bailey, R. M. Cichlaurus versus Cichlasoma as the name for a genus of perciform fishes. Copeia 1957: 303–304.

Bailey, W. B. see Leim, A. H.

Baird, D. A *Physonemus* spine from the Pennsylvanian of West Virginia. J. Paleont. **31**:1010-1018, figs.

Baker, H. see Nigrelli, R. F.

Bal, D. V. see Marathe, V. B.

(Balabai, P. P.) (1). [On the phylogeny of the Agnatha.] Zool. Zh., Moscow 35 1956: 874-890 12 figs. [Russian with English summary.]

[Balabai, P. P.] (2). [K voprosu o vertika'nom rasprostranenii pteraspid podol'skogho Old-Reda.]. Geol. Sbornik 4:305–309 2 t-figs.

Balabai, P. P. & Opalatenko, L. K. On the fauna of the Old Red and the Podalian bed. Rep. Acad. Sci. Ukr. 1957: 406–409 3 figs. [English summary.]

Balon, E. (1). The development of scales in Rutilus rutilus (L.) and Leucaspius delineatus (Heck.). Polsk. Arch. Hydrobiol. 3 1956:175–187 17 figs. [English summary.]

Balon, E. (2). Zur ichthyofauna des Orava-Flusses. Zool. Listy 5 1956: 325– 337 1 fig. [German summary.]

Bănărescu, P. (1). Variation du torus longitudinal du cerveau chez les poissons téléostéens. Comun. Acad. Republ. Pop. rom. 6 1956: 893-899 6 figs. [French summary.]

Bănărescu, P. (2). Distribution et rôle biologique des Gobies dans le complexe lagunaire Razelm. Bul. Inst. Cercet. Pisc. Roman. 16 No. 2: 36-45. [French summary.]

Bănărescu, P. & Nalbant, T. Eidonomische und taxionomische Untersuchungen an *Cobitis elongata* (Pisces, Cobitidae). Senckenbergiana biol. 38: 283–294 7 figs.

Baraduc, M. M. Influence de la thyroxine sur les échanges de l'ion chlore de la Triute arc-en-ciel (Salmo gairdnerii Rich.). C.R. Soc. Biol. Paris 151: 303-305.

(Baranenkova, A. S.). [Data on the distribution of sea perches of the genus Sebastes.] C.R. Acad. Sci. URSS 113: 468-471 3 figs. [In Russian.]

(Barannikova, I. A.). [Completion of spawning condition of autumn migrating female and male winter sturgeon after exclusion from freshwater during the spawning migration.] C.R. Acad. Sci. URSS N.S. 99 1954: 641–644 2 figs. [In Russian.]

Bardach, J. E. The sensitivity of the goldfish (*Carassius auratus* L.) to point heat stimulation. Amer. Nat. 90 1956: 309-317 1 fig.

Bardach, J. E. & Bjorklund, R. G. The temperature sensitivity of some American freshwater fishes. Amer. Nat. 91:233-251 2 figs.

Barets, A. Les récepteurs intramusculaires des nageoires chez les sélaciens. Arch. Anat. micr. Morph. exp. 45 1956: 254–260 5 figs.

Barets, A. & Le Touzé, S. Identification dans le muscle latéral d'un Téléostéen (*Tinca tinca* L.) de deux systèmes moteurs, "lent" et "rapide". C.R. Acad. Sci. Paris 242 1956 : 1230–1233 2 figs.

Barnoud, R. see Cordier, D.

Barr, L. M. see Prosser, C. L.

Barraclough, W. E. An abnormal long-nosed skate (*Raja rhina*) from British Columbia. Rep. prov. Mus. Vict., B.C. 1955 1956: D57-58 1 fig.

Barrett, J. H. see Bassindale, R.

Barrington, E. J. W. see Brown, M. E.

Barsukov, V. V. (1). Biological information on the increase of White Sea wolf fish (Anarhichas lupus). Zool. Zh., Moscow 32 1953:1211-1216. [In Russian.]

Barsukov, V. V. (2). White Sea wolf-fish (Anarhichas lupus maris-albi Barsukov). Ichth. Probl., Moscow No. 6 1956: 129–135 1 pl. [In Russian.]

(Barsukov, V. V.) (3). [Distribution of Anarhichus lupus, A. minor and A. latifrons in Barents Sea.] Ichth. Probl. Moscow 8: 28-50 13 figs. [In Russian.]

Baschieri-Salvadori, F. Spedizione subacqua Italiani nel Mar Rosso. Ricerche zoologiche IX. Pomacentridi. Riv. Biol. colon. 15:57–68 6 pls.

Bassindale, R. & Barrett, J. H. The Dale Fort marine fauna. Proc. Bristol Nats' Soc. 29: 227-328 3 figs.

Bauch, G. Der Elbachs (Salmo salar L.), seine Biologie und wirtschaftliche Bedeutung. Z. Fisch. 6:241-250, 3 figs.

Bauchot, M.-L. & R. & Lubet, P. Étude de la fauna ichthyologique du Bassin d'Arcachon (Gironde). Bull. Mus. Hist. nat. Paris (2) 29:385–406 2 figs.

Bauchot, R. see Thomopoulos, A.

[1957]

Bauer, K. & Schubert, P. Proterorhinus marmoratus Pallas (Gobiidae) ein für die österreichische Fauna neuer Fisch. Burgenländ. Heimat Bl. 19:6-9 3 figs.

Baxter, E. W. The development of the intestinal blood system of lampreys (Petromyzonidae), with an account of the origin of vascular couples. Proc. zool. Soc. Lond. 129: 371-396 3 pls., 13 figs.

(Beklemishev, K. V.). [The effect of sound waves in the sea on the vertical distribution of fishes and plankton]. Adv. mod. Biol. Moscow 41 1956: 90-96 3 figs. [In Russian.]

Belton, L. Étude sommaire d'un moulage natural de la cavité cranienne d'un Borcosomus de l'Eotrias de Madagascar. C.R. Acad. Sci. Paris 245: 549-551 1 fig.

Bemmel, A. C. V. van. Vissoorten der uit de Nederlandse fauna dreigen te verdwijnen. Levende Nat. 60: 109–114.

Benl, G. (1). Tetraodon leiurus brevirostris, subsp. nova. Ein Nachtrag zu den Süsswasser-Kugelfischen in Datz 9 (6-8), 1956. Aquar.-Terrar. Z. 10:63-65, 4 text-figs.

Benl, G. (2). Carinotetraodon chlupatyi nov. gen., nov. spec. ein Kugelfisch mit Kamm und Kiel. Opusc. Zool. No. 5:1-4, 1 fig.

Ben-Tuvia, A. see Steinitz, H.

Bergman, P. see Welander, A. D.

Bergquist, H. On early cell migration processes in the embryonic brain. J. Embryol. exp. Morph. 4 1956: 152–160, 1 pl., 8 figs.

Bernard, F. Existence d'une région à épithélium gastrique dans la tube digestif de l'alevin de Gardon (*Leuciscus rutilus* L.). C.R. Acad. Sci. Paris 243 1956: 1161-1162.

Bernard, F. Recherches sur l'organogenèse de la bouche, des fentes branchiales et du coelome antérieur chez l'embryon de truite (Salmo trutta L.). Trav. Lab. hydrobiol. Grenoble 48: 109-140 6 pls.

Berrill, N. J. The origin of Vertebrates. Oxford (O.U.P.) 1955 pp. viii + 257 31 text-figs.

(Bervald, E. A.). [Influence of industry on productivity of bream shoals in Aral sea.] Ichth. Probl., Moscow 7 1956: 21-32 3 figs. [In Russian.]

Best, E. A. (1). Recent occurrences of the red brotula, *Brosmophycis marginata* (Ayres), in Californian waters. Calif. Fish Game 43 No. 1:97–98.

Best, E. A. (2). Tagged dover sole (*Microstomus pacificus*) at liberty six years. Calif. Fish Game 43:147.

Bhatti, H. K. Development of tailfin in Teleostei. Proc. 4th Pakistan sei. Conf. pt. 3 1952: 68-69. [Abstract.]

Bhatti, H. K. & Aminul Hug, A. K. M. On the structure of scales in Ophicephalidae. Proc. 4th Pakistan sci. Conf. pt. 3 1952: 67-68. [Abstract.]

Bhatti, I. H. (1). On the history of barbels of a common siluroid fish, *Rita rita* Hamilton. Proc. 4th Pakistan sci. Conf. pt. 3 1952:65-66.

Bhatti, I. H. (2). On the cutaneous sense organs of a common siluroid fish, *Rita rita* Hamilton. Proc. 4th Pakistan sci. Conf. pt. 3 1952: 66-67. [Abstract.]

Bhuiya, M. A. H. & Siddiqui, A. A. On the afferent branchial arteries in *Ophicephalus punctatus* Bloch. Proc. 4th Pakistan sci. Conf. pt. 3 1952: 69. [Abstract.]

Bigelow, H. B. & Schroeder, W. C. A study of the sharks of the suborder Squaloidea. Bull. Mus. comp. Zool. Harv. 117: 1-150 4 pls.

(Birman, I. B.) (1). [On the causes of one of the peculiarities in Oncorhynchus gorbuscha (Walb.) in the Sea of Japan.] Zool. Zh., Moscow 35 1956: 1681–1684. [Russian with English summary.]

(Birman, I. B.) (2). [Local shoals of the autumn Keta Salmon in the Amur basin.] Ichth. Probl., Moscow 7 1956: 158-173. [In Russian.]

(Birman, I. B.) (3). [The Kuro-Sio and the stock of Oncorhynchus keta infra sp. autumnalis.] Ichth. Probl. Moscow 8: 3-7 2 figs. [In Russian.]

(Biryukov, N. P.) [Utilisation of growth-rate studies for distinguishing migrations and populations of Atlantic-Scandinavian herring.] Ichth. Probl. Moscow 6 1956: 47–54 4 figs. [In Russian.]

Bjerkan, P. Notes on the Greenland shark Acanthorhinus carcharias (Gunn).

1. The reproduction problem of the Greenland shark, Fiskeridir, Skr. Havundersøk. 11 No. 10: 1-7.

Bjorklund, R. G. see Bardach, J. E.

Black, E. C. (1). Alterations in the blood level of lactic acid in certain salmonoid fishes following muscular activity. I. kamloops trout, Salmo gairdneri. J. Fish. Res. Bd. Can. 14: 117-134 4 figs.

Black, E. C. (2). Alterations in the blood level of lactic acid in certain salmonoid fishes following muscular activity. II. Lake trout, Salvelinus namaycush. J. Fish. Res. Bd. Can. 14:645-649.

Black, V. S. see Brown, M. E.

Blair, A. A. (1). Salmon tagging at Francis Harbour Bight, Labrador. J. Fish. Res. Bd. Can. 14: 135-140 1 fig.

Blair, A. A. (2). Salmon tagging at Cape Charles, Labrador. J. Fish. Res. Bd. Can. 14: 141-144 1 fig.

Blanc, M. Sur les poissons des grands lacs africains. C.R. Soc. Biogéogr. 31 1954: 35–38.

Blanco, G. J. & Montalban, H. R. A bibliography of Philippine fishes and fisheries. Philipp. J. Fish. 1 1956: 107–130.

Blanco, G. J. & Villadalid, D. V. The young of some fishes of Luzon. Philipp. J. Fish. 1 1951:67-93 35 figs.

Bloch, J. see Gros, C. M.

Blok, J. W. de. On the architecture of the asymmetrical head in *Pleuronectes platessa* L. IV. Proc. Akad. Wet. Amst. 60C: 77-89 5 figs.

Boden, B. P. see Kampa, E. M.

(Bodrova, N. V., Krayukhin, B. V., Malyarevskaya, A. Y.). [Importance of artificial food for young carp.] Ichth. Probl. Moscow 6 1956: 149-155. [In Russian.]

Boeseman, M. (1). On a collection of East Asian fishes. Zool. Meded. 35: 69-78 1 pl.

Boeseman, M. (2). On a collection of fishes from Stanley Pool (Belgian Congo). Zool. Meded. 35:139-151 1 pl.

Böhlke, J. (1). A new shallow-water brotulid fish from the Great Bahama Bank. Notul. nat. Acad. Philad. No. 295: 1-8 1 fig.

8

Böhlke, J. (2). The Bahaman species of Emblemariid blennies. Proc. Acad. nat. Sci. Philad. 109: 25-57 3 pls. 3 figs.

Böhlke, J. (3). On the occurrence of garden eels in the Western Atlantic, with a synopsis of the Heterocongrinae. Proc. Acad. nat. Sci. Philad. 109: 59-79 1 pl. 7 figs.

Böhlke, J. (4). A review of the blenny genus *Chaenopsis*, and the description of a related new genus from the Bahamas. Proc. Acad. nat. Sci. Philad. 109:81-103 2 pls. 3 figs.

Böhlke, J. & Chaplin, C. C. G. Oral incubation in Bahaman jawfishes *Opisthognathus whitehursti* and *O. maxillosus*. Science 125: 353 1 fig.

Böhlke, J. & Cohen, D. M. The Eastern Pacific species of *Callionymus*. Copeia **1957**: 159–160.

Böhlke, J. & Mead, G. W. A new blenny from the coast of western Mexico. Notul. nat. Acad. Philad. No. 301: 1-8 1 pl. 1 fig.

Bolin, R. L. Age and growth of the myctophid fish *Lampanyctus leucopsarus* (Eigenmann & Eigenmann). Int. Congr. Zool. 14 1953 (1956): 541-542.

Bondy, P. K., Upton, G. V. and Pickford, G. E. Demonstration of cortisol in fish blood. Nature, London 179: 1354-1355.

Bont, A. F. de (1). Premiers essais de rizipisciculture à la station de recherches piscicoles (Katanga). Bull. Agric. Congo belge 46 1955 : 1550–1553 2 figs.

Bont, A. F. de (2). Lutte contre les mollusques dans les eaux africaines. Bull. Agric. Congo belge 47 1956: 377-380.

Bont, A. F. de (3). Controle biologique des mollusques d'eau douce et des maladies qu'ils transmettent. Ann. Soc. belge Méd. trop. 36 1956: 667-672.

Bont, A. F. de & Maes, H. Relation entre le frai du *Labeo altivelis* Peters et la conductivité des eaux. Hydrobiologia 8 1956 : 288-292.

Bont-Hers, A. F. & M. J. de. *Haplo-chromis mellandi* Blgr. Poisson malacophage. (Fam. Cichlidae). Rev. zool. bot. Afr. 53 1956: 370–376 1 fig.

Borisov, P. G. & Ovsyannikov, N. S. [Key to fishes of economic importance in the USSR]. Moscow, 1954 pp. 260 306 figs. [In Russian.]

(Borutzki, E. V.). (Method of studying the feeding of herbivorous fishes.] Trud. Sovesh. Ikhtiol. Komm. 6 1954 (1955): 54-61. [In Russian.]

Boschung, H. T. Some interesting records of marine fishes from Alabama. J. Ala. Acad. Sci. 29: 33-35.

Bouillon, J. see Brien, P.

Boursot, J. B. The poraqué. Aquarist & Pondkpr. 22: 99-100; 130-131 2 figs.

Boute, A. Sur une cause possible des hécatombes de Poissons. C.R. Soc. géol. Fr. 1957: 104–105.

Bowers, A. B. (1). Age and size of Manx herring in 1954. Rep. Mar. biol. Sta. Pt. Erin. 67 1954 [1955]: 31-33.

Bowers, A. B. (2). Age and size of Manx herring in 1955. Rep. Mar. biol. Sta. Pt. Erin No. 68, 1955 [1956]: 17–19.

Bowers, A. B. (3). Age and size of Manx herrings in 1956. Rep. Mar. Biol. Sta. Pt. Erin. 69 1956: 18-20.

(Braginskaya, R. Y.). [The development of Alosa kessleri volgensis and A. kessleri kessleri.] Trud. Inst. Morf. Zhiv. 16:171-180 fig. [In Russian.]

Braginskaya, R. Y. see Vasnetzov, V. V.

Brandon, A. M. see Cordier, D.

Brânzei, P. & Jitariu, P. Notes sur l'epilepsie experimentale. La dynamique des chromatophores pendant l'epilepsie experimentale chez certaines espèces de poissons marins. Stud. Cercet, Acad. române. Fil. Iasi 2 1–2 1951: 210–225. [French summary.]

Brayshaw, J. D. see Hartley, W. G. Breder, C. M. A note on preliminary stages in the fossilization of fishes. Copeia 1957: 132-135 1 pl.

Brett, J. R. Salmon research and hydro-electric power development. Bull. Fish. Res. Bd. Can. No. 114: 1–26 2 figs.

Pisces 9

Brett, J. R. see Alderdice, D. F.

Brett, J. R. see Brown, M. E.

Breuer, J. P. Ecological survey of Baffin and Alazan Bays, Texas. Publ. Inst. Mar. Sci. Univ. Tex. 4 No. 2: 134-155.

Brien, P., Poll, M. & Bouillon, J. Éthologie de la reproduction d'un Dipneuste africain *Protopterus dolloi* (Boulenger). C.R. Acad. Sci. Paris **245**: 1988–1990.

Briggs, J. C. A new genus and two new species of Eastern Atlantic clingfishes. Copeia 1957: 204-208 3 figs. 1 pl.

Briggs, J. C. & Caldwell, D. K. Acanthurus randalli, a new surgeon fish from the Gulf of Mexico. Bull. Fla St. Mus. 2:43-515 figs.

Briggs, J. C. see Caldwell, D. K.

Broadhead, G. C. Changes in the size structure of the yellow fin tuna population of the tropical Eastern Pacific Ocean from 1947–1955. Bull. Inter-Amer. Tuna Comm. 2: 1–20.

Brown, D. G. & Rosen, W. G. A progress report on the study of perch movement in Lake Okoboji, Dickinson County, Iowa. Proc. Iowa Acad. Sci. 58 1951: 423-434 2 figs.

Brown, J. L. (1). Distinguishing characters of the cyprinodont fishes, Fundulus cingulatus and Fundulus chrysotus. Copeia 1956 No. 4 [1957]: 251-255.

Brown, J. L. (2). A key to the species and subspecies of the cyprinodont genus Fundulus in the United States and Canada east of the Continental Divide. J. Wash. Acad. Sci. 47: 69-77.

Brown, M. E. & others. The physiology of fishes. Vol. 1. Metabolism pp. xiii + 447; Vol. 2. Behaviour pp. xi + 526 figs. New York (Academic Press).

Brown, W. H. see Hubbs, C.

Brull, L. L'azote non-protéique du plasma de Scyliorhinus canicula. Vie et Milieu 7 1956 : 521-522.

Bsharah, L. Plankton of the Florida Current. V. Environmental conditions, standing crop, seasonal and diurnal changes at a station forty miles east of Miami. Bull. mar. Sci. Gulf & Caribbean 7: 201-251 45 pls.

Buchholz, M. Age and growth of river carpsucker in Des Moines River, Iowa. Proc. Iowa Acad. Sci. 64: 589-600 1 fig.,

Buckridge, T. N. see Donaldson, L. R.

Budker, P. see Le Danois, E.

Buen, F. de (1). La oceanografia frente a las costas del Uruguay. Ann. Mus. Hist. nat. (2) 6 1953: 1-37 12 figs.

Buen, F. de (2). Los peces de la familia Molidae. Invest. zool. Chil. 4:66-76 6 figs.

Buen, F. de (3). Suborden Luvaroidei, nuevo para la fauna Chilena. Invest. zool. Chil. 4:83-88 2 figs.

[Buistrov, A. P.]. [Evolyntziya zubov pozvonchnuikh.] Annu. Soc. paléont. russe 14 1953 : 39–60 figs.

Bull, H. O. see Brown, M. E.

Bullis, H. R. see Springer, S.

Burd, A. C. see Cushing, D. H.

(Burdak, V. D.). [Peculiarities in the ontogenetic and phylogenetic relationship in Black Sea mullet (Mugil saliens, M. auratus, M. cephalus).] Trav. Stabiol. Sebastapol 9:243-273 16 figs. [In Russian.]

(Burmakin, E. V.). [On changes in the morphology of carp acclimatized in the Lake Balkhash basin.] Zool. Zh., Moscow 35 1956:1887-1891. [Russian with English Summary.]

Buschendorf, C. Eine neue Zuchtform des Scalare. Aquar. Terrar. Z. 10:200, 1 text-fig.

Buşniţă, Th. & Arachelian, I. Quelques données concernant la pêche maritime roumaine et bulgare dans l'intervalle 1940-1954. Bul. Inst.-Cercet. Pisc. Roman. 16 (1):13-35. [French summary].

Buşniţă, Th. & others (1). Premiers essais d'acclimatation des Corégones dans les eaux de la Republique Populaire Roumaine. Bul. Inst. Cercet. Pisc. Roman 16 No. 2:5-19. [French summary.]

Buşniţă, T., & others (2). La gynogénèse chez Carassius auratus gibelio Bloch. Comun. Acad. Rep. Pop. Romîne 7:81-87 3 pls. [French summary.]

Butler, R. L. The development of a vinyl plastic subcutaneous tag for trout. Calif. Fish Game 43: 201-212, 4 figs.

(Butzkaya, N. A.). [On the peculiarities of the function of the testis in fish in various types of spawning.] C.R. Acad. Sci. URSS 100 1955: 809-812 fig. [In Russian.]

Buznikov, G. A. see Baburina, E. A. Bystrow, A. P. The microstructure of dermal bones in arthrodires. Acta Zool. Stockholm, 38: 239-275, 25 text-figs.

Cailleux, A. see Chavan, A.

Cairns, J. & Scheier, A. The effects of temperature and hardness of water upon the toxicity of zinc to the common bluegill (*Lepomis macrochirus* Raf.). Notul. Nat. Philad. No. 299:1-12 tabs.

Caldito, E. B. see Sulet, J. I.

Caldwell, D. K. The biology and systematics of the pin fish, Lagodon rhomboides (Linnaeus). Bull. Fla. St. Mus. Biol. Sci. 2:77–173 22 figs.

Caldwell, D. K. & Briggs, J. C. Range extensions of Western North Atlantic fishes with notes on some soles of the genus *Gymnachirus*. Bull. Fla. Acad. Sci. 2 1:1-11.

Caldwell, D. K. see Briggs, J. C. Caldwell, D. K. see Kilby, J. D.

Callan, H. G. The lampbrush chromosomes of Sepia officinalis L., Anilocra physodes L. and Scyllium catalus Cuv. and their structural relationship to the lampbrush chromosomes of Amphibia. Pubbl. Staz. zool. Napoli 29: 329-346 2 pls. 4 text-figs.

Cameron, I. A preliminary report on examination of salmon and scale readings. Salm. Trout Mag. No. 151: 256-274 15 figs.

Canagaratnam, P. & Medcof, J. C. Ceylon's beach seine fishery. Bull. Fish. Res. Sta. Ceylon No. 4 1956: 1–32 5 figs.

Cantacuzene, A. M. see Devillers, C.

Capco, S. R. & Manacop, P. R. The Abuyan, an improved type of goby fishing gear, used in Laguna de Bay with notes on the composition of the commercial catches. Philipp. J. Sci. 3 1955: 65-84 7 pls., 8 figs.

Caraiman, I. G. Observations sur le rythme de croissance des petits de la carpe de culture de l'étang Tataraseni. Bul. Inst. Cercet. Pisc. Roman 16 No. 2,: 53-60.

[1957]

Carlander, K. D. An unusually large population of fish in a gravel pit lake. Proc. Iowa Acad. Sci. 58 1951: 435-440.

Carlander, K. D. & Moorman, R. B. Standing crops of fish in Iowa ponds. Proc. Iowa Acad. Sci. 63 1956: 659–668.

Carlo, J. M. de & Lopez, R. B. La valvula intestinal del pejerrey. Notas mus. Univ. nac. La Plata, Zool. 19: 161-169 6 figs.

Carmin, J. Treasures of our seashore. Tel-Aviv (Yavnek Publ. House) pp. 368, text illust.

Carter, G. S. see Brown, M. E.

Casasco, E. Primi dati sulla innervazione recettrice e vegetativa del labirinto membranoso dei pesci. Monit. Zool. Ital. 65 Suppl.: 164-166 2 figs.

Cassie, R. M. Condition factor of snapper, *Chrysophrys auratus* Forster, in Hauraki Gulf. N.Z. J. Sci. Tech. 38B: 375-388 3 figs.

Cautis, I. see Nicolae, I.

Cautis, I. see Pora, E. A.

Cavaliere, A. Su rari stadi larvali e giovanili di *Pomatomichthys constanciae* Gigl. e relativa differenziazione da stadi giovanili della specie affine *Pomatomus telescopium* Risso. Boll. Pesca Piscic. Idrobiol. 32 fasc. 1 1956 [1957]: 121–127 1 pl.

Chambers, R. & E. L. see Kao, C. Y.

Chaplin, C. C. G. see Böhlke, J.

Chaturin, B. M. see Alverson, D. L.

Chavan, A. & Cailleux, A. Détermination practique des fossiles Paris, : 1-385 figs.

Chi, Y. S. Upper Devonian Bothriolepis beds of Yunnan. Sci. Rec. Acad. Sinica 1 1944:171-174.

Chlupaty, P. Dascyllus aruanus, der Preussenfisch. Aquar. Terrar. Z. 10: 209 figs. 1 and 2.

Chopard, L. Un poisson qui peut vivre sans eau. Le protoptère. Nature, Paris No. 3261 : 30-34 6 figs.

(Chubareva, L. A.) [Cytological and cytochemical investigations on fertilization in Lampetra fluviatilis,] C.R. Acad. Sci. URSS 112:945-948 3 figs. [In Russian.]

Čihăr, J. Systematical and biological notes on the pike (Esox lucius L.). Univ. Carolina, Pragae 1 Biol. 1 1955: 1-18 3 figs. [English summary.]

Citterio, P. & others. On the presence of actin and myosin in the electric organ of the *Torpedo*. Pubbl. Staz. zool. Napoli 29: 434–438 2 figs.

Citterio, P. see Maldacea, L.

Clancey, J. F. A contribution to the life history of the fish, Bregmaceros atlanticus Goode and Bean, from the Florida current. Bull. mar. Sci. Gulf and Caribbean 6 1956: 233-260 8 figs.

Clark, E. E. see Humm, D. G.

Clark, M. R. see Musacchia, X. J.

Clavert, J. & Zahnd, J. P. (1). Modifications hépatiqués durant la gestation chez deux poissons ovovivipares (Xiphophorus helleri et Lebistes reticulatus). C.R. Soc. Biol., Paris 150: 2257–2259.

Clavert, J. & Zahnd, J. P. (2). Modifications hépatiques survenant pendant la vitellogenèse chez deux espèces de poissons ovovivipares (Xiphophorus helleri et Lebistes reticulatus). C.R. Soc. Biol. Paris 150 1956: 1261–1263.

Cleary, R. E. An annotated checklist of fishes of the Wapsipinicon River drainage system in Iowa. Proc. Iowa Acad. Sci. 59 1952: 435–441 1 fig.

• Clemens, H. B. Fishes collected in the tropical eastern Pacific, 1954. Calif. Fish Game 43: 299-307 1 fig.

Clements, M. The proteolytic activity of the thyroid and other tissues of the dogfish, Scyliorhinus (Scyllium) canicula. Bull. Inst. Oceanogr. Monaco No. 1091: 1-10 3 figs.

Coates, C. W. see Atz, J. W.

Coffin, G. W. see Scattergood, L. W.

Cohen, D. M. Notes on the identity of several salariine blennies of the genus *Scartichthys* from western South America. Copeia 1956 No. 4 [1957]: 246-248.

Cohen, D. M. see Böhlke, J.

Colas, J. see Devillers, C.

Concha, B. F. & Vargas, F. F. Actividad eléctrica del corazón del Teleósteo Sicyases sanguineus. Invest. Zool. Chil. 3:96–105 6 figs.

Concha, B. F. see Vargas, F. F.

Cooper, L. B. & Hemphill, A. F. Record soft he lamprey Lampetra aepyptera from Alabama. Copeia 1957: 54

Coraboeuf, E. see Gargouil, Y. M.

Cordier, D., Barnaud, R. & Brandon, A. M. Influence du passage de l'eau de mer à l'eau douce et de l'eau de mer sursalée sur les échanges respiratoires du Gobie (Gobius auratus et Gobius lota). C.R. Soc. Biol. Paris 151: 739-742 2 figs.

Cordier, D. & Maurice, A. (1). Étude sur l'absorption intestinale des Sucres chez l'anguille (*Anguilla vulgaris* L.) vivant dans l'eau de mer ou dans l'eau douce. C.R. Soc. Biol. Paris 150: 1957-1959.

Cordier, D. & Maurice, A. (2). Recherches sur l'absorption intestinale des solutions de pentoses et d'hexoses chez le Brochet (*Esox lucius*, L.). C.R. Soc. Biol. Paris 151:116-118.

Cordier, D. & Worbe, J. F. Influence de pH du milieu intestinal sur l'absorption du glucose chez la Rascasse (Scorpaena porcus, L.). C.R. Soc. Biol. Paris 151: 121-124.

Costea, E. see Vasiliu, G. D.

Couteaux, R. & Laurent, P. Étude au microscope électronique du coeur de l'Anguille, observations sur la structure du tissu musculaire de l'oreillette et son innervation. C.R. Acad. Sci. Paris 245: 2097–2100 10 figs.

Creteanu, A. see Gyurko, S.

Cristea, A. & E. see Teodorescu-Leonte, R.

Cristian, A. see Busnita, T.

Crossman, E. J. Growth, mortality and movements of a sanctuary population of maskinonge (*Esox masquinongy* Mitchill). J. Fish. Res. Bd. Can. 13 1956: 599-612.

Cruz, J. A. M. (1). A sardinha do norte de Portugal (Sardina pilchardus Walb.). Características etárias e de reprodução. An. Fac. Ciênc. Porto 37 1953: 217–250 3 pls. [English summary.] Cruz, J. A. M. (2). Sur la succéssion de générations dans la pêche à la sardine, au nord du Portugal, et ses possibles conséquences d'ordre économique. An. Fac. Ciênc. Porto 38 1954: 72–75 fig.

Curitan, A. see Gyurko, S.

Curry-Lindahl, K. Fiskarna i Färg-Stockholm (Almquist & Wiksell) 1953, pp. viii + 189, 64 pls. col. [incl. in pagination].

Cushing, D. H. (1). The interpretation of echo traces. Fish. Invest. Lond. Ser. 2, 21 No. 3:1-16 14 figs.

Cushing, D. H. (2). The number of pilchards in the Channel. Fish. Invest. Lond. Ser. 2, 21 No. 5: 1-27, 15 figs.

Cushing, D. H. & Burd, A. C. On the herring of the Southern North Sea. Fish. Invest. Lond. Ser. 2, 20 No. 11: pp. iv + 31, 16 figs.

Cushing, J. E. & Durall, G. L. Isoagglutination in fish. Amer. Nat. 91:121-126.

Daget, J. Données récentes sur la biologie des poissons dans la delta central du Niger. Hydrobiologia 9: 321-347 5 figs.

Dalenius, P. Studium av en elritspopulation i en lappländsk fjällsjö. Fauna och Flora 52:125–136 2 figs.

Danil'chenko, P. G. Novuii rod semeistva Bregmacerotidae. Mater. K. Osnovam Palaeont., Akad. Nauk. SSSR 1:57-59 1 text-fig.

David, L. R. Fishes (other than Agnatha) in Treatise on marine ecology and paleoecology. Vol. 2. Mem. Geol. Soc. America 67: 999–1010.

Davies, D. H. (1). The South African pilchard (Sardinops occillata). Size composition of the commercial catches, 1950–55. Invest. Rep. Div. Fish. S. Afr. No. 26:1–59 14 pls. 6 figs. tabs.

Davies, D. H. (2). The South African pilchard (Sardinops ocellata). Preliminary report on feeding off the west coast, 1953–56. Invest. Rep. Div. Fish. S. Afr. No. 30:1–40, 17 pls.

Davies, D. H. (3). The biology of the South African pilchard (Sardinops ocellata). Invest. Rep. Div. Fish S. Afr. No. 32: 1-11 3 figs.

Dawson, E. Y. see Mohr, J. L.

Day, L. R. see Leim, A. H.

(Demetrashvih, M. G.). [The fishes of Khramisk water reservoir and of the rivers Khrami and Agri-Chai]. Trud. Inst. Zool. Tiflis 13 1954:133-178. [In Russian.]

Denton, E. J. Recherches sur l'absorption de la lumière par le cristallin des poissons. Bull. Inst. oceanogr. Monaco No. 1071 1956: 1–10 2 figs.

Denton, E. J. & Warren, F. J. The photosensitive pigments in the retinae of deep-sea fish. J. Mar. Biol. Ass. U.K. 36:651-662 2 figs.

Denzer, W. Elektro-fischerei. Kosmos, Stuttgart 49 1953: 49-52 5 figs.

de Sylva, D. P. Studies on the age and growth of the Atlantic sail fish, *Istiophorus americanus* (Cuvier), using length-frequency curves. Bull. mar. Sci. Gulf & Caribbean 7:1-20 7 figs.

Devillers, C., Colas, J. & Cantacuzène, A. M. Gastrulation de l'oeuf de Truite (Salmo irideus) en l'absence d'oxygène on en présence d'inhibiteurs du métabolisme. C.R. Acad. Sci. Paris 245: 1461-1462.

Devillers, C. & others. Différentiation in vitro de blastodermes de truite (Salmo irideus) dépourvus de couche enveloppante. J. Embryol. Exp. Morph. 5:264-273 3 figs.

Dieuzeide, R. Les myxines (Myxine glutinosa L.) en Méditerranée. Bull. Sta. Aquic. Pêche Castiglione No. 8 1956 [1957]: 75–80.

Dieuzeide, R., & Roland, J. (1). Opérations de draguages et de chalutages effectuées au large des Côtes Algériennes au cours des années 1954 et 1955. Bull. Sta. Aquic. Pêche Castiglione No. 8 1956 [1957]: 11-27, 2 figs.

Dieuzeide, R. & Roland, J. (2). Complément au catalogue des poissons des Côtes Algériennes. Bull. Sta. Aquic. Pêche Castiglione No. 8 87-106 4 pl.

Dieuzeide, R. & Roland, J. (3). Étude biométrique de Sardina pilchardus Walb. et de Sardinella aurita C.V. capturées dans la Baie de Castiglione. Bull. Sta. Aquic. Pêche Castiglione No. 8 1956 [1957]: 111-225 28 figs.

Dieuzeide, R. & Roland, J. (4). Observations relevées sur les sardines (Sardina pilchardus Walb.) de la Baie de Castiglione parasitées par Peroderma cylindricum Heller. Bull. Sta. Aquic. Pêche Castiglione No. 8 1956 [1957]: 239-249 1 fig.

Dimitriu, M. see Busnita, J.

Dimoftache, M. see Gyurko, S.

Dineen, C. F. & Stokely, P. S. The osteology of the sacramento perch, *Archoplites interruptus* (Girard). Copeia 1956 No. 4 [1957]: 217-230 f. 1-13.

(Disler, N. N.) [The development of autumn keta (Oncorhynchus keta (Walb.)) in the river Amur.] Trud. Inst. Morf. Zhiv. 20: 3-70 16 figs. [In Russian.]

(Dmitrieva, E. N.) (1). [Stages of development of Abramis brama, Rutilus rutilus heckeli and Lucioperca lucioperca in Taganrog Bay.] Trud. Inst. Morf. Zhiv. 16:77-101 4 figs. [In Russian.]

(Dmitrieva, E. N.) (2). [Morphological and ecological analysis of two species of crucian carp.] Trud. Inst. Morf. Zhiv. 16:102-170 9 figs. [In Russian.]

Dmitrieva, E. N. see Vasnetzov, V. V.

Döderlein, L. Bestimmungsbuch für deutsche Land-und Süsswassertiere. Wirbeltiere...2, erweiterte Auflage. München (R. Oldenbourg) 1955, pp. 304, 266 figs.

Domurat, J. Embryonic development of trout (Salmo trutta L.) pike (Esox lucius L.) and roach (Rutilus rutilus L.) in the environment deprived of water. Polsk. Arch. Hydrobiol. 3 1956: 167-173. (English summary.)

Donaldson, L. R., Hansler, D. D. and Buckridge, T. N. Interracial hybridization of cutthroat trout, Salmo clarkii, and its use in fisheries management. Trans. Amer. Fish. Soc. 86 (1956) 1957: 350–360, 2 text-figs.

Donaldson, L. R. & Olson, P. R. Development of rainbow trout brood stock by selective breeding. Trans. Amer. Fish. Soc. 85 (1955): 93–101 4 figs.

Dorn, E. Das Zwischenhirn von Protopterus. Verh. dtsch. zool. Ges. 1956: 226–230, 7 figs. Dorr, J. A. A pleuracanth shark spine from the early Pennsylvanian, Saginaw formation, of Michigan. Pap. Mich. Acad. Sci. 42:99-104 1 fig.

Dôtu, Y. (1). A new species of a goby with a synopsis of the species of the genus *Luciogobius* Gill and its allied genera. J. Fac. Agric. Kyushu Univ. 11:69-76.

Dôtu, Y. (2). The life history of the small, transparent goby, *Gobius lidwilli* McCulloch. Sci. Bull. Fac. Agric. Kyushu Univ. 16:85–92 3 figs.

Dôtu, Y. (3). The life-history of the goby, Luciogobius guttatus Gill. Sci. Bull. Fac. Agric. Kyushu Univ. 16: 93-100 6 figs. [English Summary.]

Dôtu, Y. (4). On the bionomics and life history of the eel-like goby, Odontamblyopus rubicundus (Hamilton). Sci. Bull. Fac. Agric. Kyushu Univ. 16: 101-110 9 figs. [English summary.]

Dôtu, Y. (5). The bionomics and life history of the goby, *Triaenopogon barbatus* (Günther) in the innermost part of Ariake Sound. Sci. Bull. Fac. Agric. Kyushu Univ. 16: 261-274 5 figs.

Doudoroff, P. see Brown, M. E.

Doumenge, F. La pêche à Bermeo (Espagne). Vie et Milieu 7 1956 : 307-325 5 figs.

Downing, K. M. & Merkens, J. C. The influence of temperature on the survival of several species of fish in low tension of dissolved oxygen. Ann. appl. Biol. 45: 261-267 3 figs.

Downing, K. M. see Merkens, J. C.

Dragotoiu, C. see Busnita, T.

(Drapkin, E. I.). [On the nuptial attire of female sticklebacks]. Ichth. Probl., Moscow 7 1956: 119-122 1 fig. [In Russian.]

Drilhon, A. & **Fine,** J.-M. Les albumines chez quelques espèces de poissons. C.R. Acad. Sci. Paris **245**: 1676–1679.

Drilhon, A. Fine, J., Uriel, J. & Le Bourdelles, F. Étude electrophorétique des constituents du serum de l'Anguille. C.R. Acad. Sci. Paris 243 1956: 1802–1805.

Dubinina, V. N. see Shulman, G. E.

Dudek, J. Beitrag zur Bestimmung des Lichteinflusses auf die Intensität der Nahrungsaufnahme beim Güster (*Blicca bjoerkna* L.). Biólogia, Bratislava 12: 374–376). [German summary.]

Duijn, C. W. A case of spontaneous sarcoma in fish. Microscope 11; 133-138 9 figs.

Durall, G. L. see Cushing, J. E.

Dürr, W. Untersuchungen über die verschiedene Gestalt der Schuppen beim Karpfen, Cyprinus carpio L. (nach Beobachtungen an Schuppenkarpfen). Z. Fisch. 5: 325–421 69 figs.

Dziekonska, J. Studies on embryonic development of fish. I. Observations on the spawning and the embryonic development of bream in the Vistula Lagoon. Polsk. Arch. Hydrobiol. 3 1956: 291–305 18 figs. [English summary.]

Ebeling, A. W. The dentition of Eastern Pacific mullets, with special reference to adaptation and taxonomy. Copeia 1957: 173–185 3 pls. 7 figs.

Ebeling, A. see Kanwisher, J.

Edinger, T. Paired pineal organs. "Progress in Neurobiology" (Proc. 1st Internat. Meeting Neurobiol., Amsterdam, 1955, 1956: 121-129 4 text-figs.

Egami, N. Inhibitory effect of thiourea on the development of male characteristics in females of the fish *Oryzias latipes*, kept in water containing testosterone propionate. Annot. zool. jap. 30:26-30.

Egami, N. see Ishii, S.

(Egorov, A. G.). (Stocking the river Angare with grayling]. Ichth. Probl. Moscow 6 1956:121. [In Russian.]

Eibl-Eibesfeldt, I. Bericht von einer Reise zu den Galápagos-Inseln unter besonderer Berücksichtigung verhaltenskundlicher, herpetologischer und ichthyologischer Beobachtungen (2.) Aquar. Terrar. Z. 10: 38–43 figs. 3–8.

(Elanidge, R. F.) (1). [A study of the fish fauna of the river Alazan.] Trud. Inst. Zool. Tiflis 10 1951:161-186. [In Gruzian with Russian summary.]

(Elanidge, R. F.) (2). [The fish fauna of the river Sups.] Trud. Inst. Zool. Tiflis 13 1954:197-214. [In Gruzian with Russian summary.]

Ellis, R. J. & Gowing, H. Relationship between food supply and condition of wild brown trout, Salmo trutta Linnaeus, in a Michigan stream. Limnol. and Oceanogr. 2:299-308 3 figs.

Enns, T. see Scholander, P. F.

Erdman, D. S. Recent fish records from Puerto Rico. Bull. mar. Sci. Gulf & Caribbean 6 1956: 315-340.

(Eremeeva, E. F.). [The horn teeth of young Cypriniformes and their adaptive significance in various stages of fish development.] Trud. Inst. Morf. Zhiv. 16: 299-31311 fig. [In Russian.]

Eremeeva, E. F. see Vasnetzov, V. V.

(Ereshchenko, V. I.) [Fish fauna in the river Sari-Su basin.] Sborn. Rabot. Ikhtiol. Ghidrobiol. 1 1956:94-123. [In Russian.]

Ermin, R. & Gordon, M. On the regeneration of piscine melanomas. Int. Congr. Zool. 14 1953 (1956): 236.

Fahlbusch, K. Pteraspis dunensis Roemer eine heubebeitung der Pteraspiden funde (Agnathen) von Overath (Bez. Köln) Palaeontographica 108A 1-4:1-56 7 pl. figs.

(Faktorovich, K. A.) (1). [Liver regeneration in Salmo irideus]. C.R. Acad. Sci. URSS 110 1956: 301-303. [In Russian.]

(Faktorovich, K. A.) (2). [Diseases of two year old trout kept on artificial food.] Ichthyol. Prob. Moscow 6 1956: 156-164 3 figs. [In Russian.]

Fast, T. N. The occurrence of the deep-sea anglerfish, *Cryptopsaras couesii* in Monterey Bay, California. Copeia **1957**: 237–240 1 fig.

(Fedorov, A. V.) [On the adaptations of predatory fish to feeding on large and armed animals.] Zool. Zh., Moscow 35 1956: 939-940. [In Russian.]

Fernandez Yepez, A. Contribución al estudio de la familia Belonidae. Mem. Soc. Cienc. nat. La Salle 8 1948: 142-144 6 figs.

Fields, P. E. Guiding migrant salmon. Scientific Monthly, Pa. 85 July: 10–22 text-figs. 1–5.

Fine, J. M. see Drilhon, A.

Fisk, A. Experiments on the development of the acoustico-facialis complex and associated structures in the lamprey Lampetra planeri (Bloch). Proc. zool. Soc. Lond. 128: 267-278 6 figs.

Fitzgerald, J. W. Range extension for the western mudminnow, *Novumbra hubbsi* Schultz. Copeia 1957: 248.

Foersch, W. Aphyosemion cognatum Meinken, der "rote Prachtkärpfling." Aquar. Terrar. Z. 10: 197–200, colourplate and text-figs. 1–3.

Fontaine, M. & Motais, R. Influence du glucoside de disoxycorticosterone sur l'halotropisme de la jeune Anguille (civelle) et sa résistance aux eaux de mer sursalées. C.R. Acad. Sci. Paris 242 1956: 1359–1361.

Fontaine, M., Querrière, F. de la & Raffy, A. Action de l'hypophysectomie sur le métabolisme respiratoire de l'Anguille (Anguilla anguilla L.). C.R. Soc. Biol. Paris 151: 232-233.

Fontaine, M. see Arvy, L.

Ford, E. River fishing thrills in Tanganyika. Fishing Gazette (Beckenham, England) 139:1115.

Forselius, S. Studies of Anabantid fishes. I–III. Zool. Bidr. Uppsala 32: 93-597 18 pls. 78 figs.

(Fortunatova, K. R.) (1). [Method of studying feeding of predatory fishes.] Trud. Sovesh. Ikhtiol. Komm. 6 1954 [1955]: 62-84 6 figs. [In Russian.]

Fortunatova, K. R. (2). [Some data on the effect of the predators upon the size of fish population.] Zool. Zh., Moscow 36:575-586. [English summary.]

Fosberg, W. see Starr, T. J.

Fowler, H. W. (1). A synopsis of the fishes of China. Part VII. The perch like fishes (completed). Quart. J. Taiwan Mus. 9 1956: 161–354 61 figs.

Fowler, H. W. (2). Notes on Australian Fishes. Ichthyol. Notes Qd. 1:65-72, 2 figs.

Fox, D. L. see Brown, M. E.

Franca, P. da (1). Contribuição para o conhecimento do Género Mer-luccius, no atlantico oriental, ao suldo Equador. An. Ita Inv. Ultramar, Lisbon 9 No. 2 1956: 47-98 11 figs.

Franca, P. da (2). Contribuição para o conhecimento dos Carangédeos (Carangidae, Pomatomidae e Coryphaenidae) de Angola. An. Ita. Inv. Ultramar, Lisbon 9 No. 2 1956: 99–150 6 pls.

Français, Y. Le déterminisme de la "concentration" dans la nageoire dorsale de *Salmo*. C.R. Acad. Sci. Paris 243 1956: 196–199.

Francesco, M. Contributo allo studio della biologia dello scombro (Scomber scombrus L.) del mare Tirreno. Boll. Pesca Piscic. Idrobiol. 31 1956:120–148 14 figs.

François, Y. Quelques problèmes relatifs a la nageoire dorsale des Téléostéens. Année biol. (3) 33:413-4267 figs.

Frank, S. The food of common bullhead (Ameiurus nebulosus Le Seuer 1819) in the Zabe (Elbe) region. Univ. Carolina, Pragae 1 Biol. 1 1955: 19-24. [English summary.]

Frank, S. see Oliva, O.

Friedrich, G. Die Lebervenen des Karpfens (*Cyprinus carpio* L.). Z. Fisch. **5**: 303–313 3 pls.

Frost, W. E. (1). The growth of brown trout (Salmo trutta L.) in Haweswater before and after the raising of the level of the lake. Salm. Trout Mag. 1956: 266-274.

Frost, W. E. (2). A note on eels (Anguilla spp.). Piscator No. 38: [1-3] 1 fig.

Fry, D. H. see Hallock, R. J.

Fry, F. E. J. see Brown, M. E.

Fryer, G. A new species of *Gephyrochromis* from Lake Nyasa, with notes on its ecology and affinities. Rev. Zool. Bot. afr. 55: 347-352, 1 fig.

Fujioka, J. see Yoshihara, T.

Gabe, M. Contribution à l'histologie du rein de l'hippocampe (*Hippocampus guttulatus* Cuvier). Arch. Biol., Paris 68 fasc. 1:29-43 1 pl.

Gabe, M. see Arvy, L.

Gadea, E. see Vallmitjana, L.

(Gaevskaya, N. S.). [The basic tasks in the study of fish feeding in connection with the more important biological principles of fish breeding.] Trud. Sovesh. Ikhtiol. Komm. 6 1954 [1955]: 8-21. [In Russian.]

- Gail, R. & Rageau, J. Premières observations sur un poisson marin venimeux de la Nouvelle-Calédonie: la synancée (Synanceia verrucosa Bloch). Bull. Soc. Path. exot. 49 1956: 846-854 1 pl. 4 figs.
- Gargouïl, Y.-M. & Coraboeuf, É. Enrigistrement intracellulaire de l'activité électrique du ventricule de Roussette (Scyliorhinus canicula Linné). C.R. Acad. Sci. Paris 245: 1949–1952.
- Garnaud, J. (1). Note sur le squelette céphalique de *Balistes capriscus* L. Bull. Inst. océanogr. Monaco No. 108 1956: 1–12 11 figs.
- Garnaud, J. (2). Ethologie de Dascyllus trimaculatus (Rüppell). Bull. Inst. Océanogr. Monaco No. 1096: 1–10 3 figs. [English summary.]
- Garrick, J. A. F. Studies on New Zealand elasmobranchii, Part VI. Two new species of *Etmopterus* from New Zealand. Bull. Mus. Comp. Zool. Harv. 116: 171–190 4 figs.
- Garrod, D. J. A review of Lake Victoria Fishery service records 1951– 1956. E.A.F.R.O. Pamph. No. 1:1-10.

Garside, E. T. see Pitt, T. K.

- Gasowska, M. Research on a reciprocal cross-breeding of Coregonus lavaretus maraenoides Poliakow and Coregonus albula (L.). Food of the hybrids in breeding ponds. Polsk. Arch. Hydrobiol. 3 1956: 253–268. [English summary.]
- Gehringer, J. W. Observations on the development of the Atlantic sailfish Istiophorus americanus (Cuvier) with notes on an unidentified species of istiophorid. Fish. Bull. U.S., 57 1956: 139-171 40 figs.
- Genovese, S. Ulteriori osservazioni sui caratteri biometrici del tonno (Thunnus thynnus L.) Tirrenico—Il tonno intergenetico dello Stretto di Messina. Boll. Pesca Piscic. Idrobiol. 32 fasc. 1 1956 [1957]: 83–120.
- George, M. G. The anatomy and histology of the swim-bladder in *Ophicephalus* (Bloch) or *Channa* Scopoli. J. zool. Soc. India 8 1956:181-194 6 figs.
 - Georgescu, I. see Popescu-Gorj, A.

[Ghlikman, L. S.] (1). [O sistematicheskom zhachenii dobavochnenkh zubtzov na Zubakh akul semeīstv Lamnidae i Scapanorhynchidae.] Trudui Geol. Muz. Karpinskogho 1:103-109 figs

[1957]

- [Ghlikman, L. S.] (2). [O gheneticheskoi svyazi semeistv Lamnidae i Odontaspidae i novuikh rodakh verkhnemelovuikh Lamnid.] Trudui Geol. Muz. Karpinskogho 1:110-117 figs.
- [Ghorbach, L. P.] [Ob Olighotzenovuikh Macrurida Vostochnuikh Karpat.] Annu. Soc. paléont. russe 16: 286–289, fig.
- Gibbs, R. H. (1). Cyprinid fishes of the subgenus Cyprinella of Notropis. I. Systematic status of the subgenus Cyprinella, with a key to the species exclusive of the lutrensis-ornatus complex. Copeia 1957: 185-195.
- Gibbs, R. H. (2). A taxonomic analysis of Myctophum affine and M. nitidulum, two lantern-fishes previously synonymized, in the western North Atlantic. Deep-Sea Res. 4:230-237 5 figs.
- Gibbs, R. H. (3). Cyprinid fishes of the subgenus Cyprinella of Notropis. III. Variation and subspecies of Notropis venustus (Girard). Tulane Stud. Zool. 5: 175-203 6 figs.
 - Gibbs, R. H. see Mather, F. J.
- Gibson, F. A. (1). Further specimens of shads from Irish waters. Irish Nats'. J. 12 1956: 109.
- Gibson, F. A. (2). Further specimen of Greater weaver, *Trachinus draco* L. Irish Nats'. J. 12 1956: 110.
- Gibson, F. A. (3). Further specimens of Red snake-fish, Cepola rubescens L. Irish Nats'. J. 12: 138-139.
- Gibson, F. A. (4). Specimen of *Urophycis blennoides* Brunn. taken off Dunmore East, Co. Waterford. Irish Nats'. J. 12: 202-203.
- (Ginzburg, A. S.). [Lipids in the occytes and in the eggs of *Acipenser stellatus*.] C.R. Acad. Sci. URSS **111** 1956: 236–239 Pl. [In Russian.]
- Godfrey, H. Mortalities among developing trout and salmon ova following shock by direct-current electrical fishing gear. J. Fish. Res. Bd. Can., 14: 153–164.

Gogol', V. A. Systematic characters of Gambusia acclimatized in Uzbekistan. Zool. Zh., Moscow 36: 459-462. (English summary.)

Good, R. Australasian floras. Nature, London, 179: 926.

Goodwin, D. Picture the fish. Nat. Hist., N.Y. 65 1956: 440-442 figs.

Gordon, M. Hereditary differences in seven natural populations of the platyfish, *Xiphophorus maculatus*. Int. Congr. Zool. 14 1953 (1956): 172–176 3 figs.

Gordon, M. see Brown, M. E.

Gordon, M. see Ermin, R.

Gordon, M. S. Observations on osmoregulation in the arctic char (Salvelinus alpinus L.). Biol. Bull. Woods Hole 112:28-33

Gordon, M. S. & Backus, R. H. New records of Labrador fishes with special reference to those of Hebron Fjord. Copeia 1957:17-20.

Gossington, R. An aid to fish handling—tricaine. Aquarium J. 28:318-321 2 figs.

(Gosteeva, M. N.) [Peculiarity of the development of vobla, the Aral sea roach.] Ichth. Probl. Moscow 6 1956: 105-112 4 figs. [In Russian.]

Gottenbos, J. J. On the correlation between the muscles in the skull structure in the regio occipitalis and the regio otica in *Raja batis* L. I–III. Proc. Akad. Wet. Amst. **59C** 1956: 541–553; 723–737 11 figs.

Gottlieb, E. The age and growth of the red mullet *Mullus barbatus* L. in Israel waters 1953–1955. Bull. Sea Fish. Res. Sta. Israel No. 12 1956: 1–20 11 figs.

Gowing, H. see Ellis, R. J.

Graham, D. H. A treasury of New Zealand fishes 2nd edition, Wellington, N.Z. (A. H. & A. W. Reed) 1956, pp. 424 frontis. (port.) text illust.

Grangaud, R., Vignais, P., Massonet, R. & Moatu, J.-P. Biosynthèse de la vitamine A chez *Gambusia holbrooki* Grd. C.R. Acad. Sci. Paris **243** 1956: 1170–1172.

Grant, E. M. see Stephenson, W.

Grassl, E. F. (1). Pelleted dry rations for trout propagation in Michigan hatcheries. Trans. Amer. Fish. Soc. 86 (1956): 307–322 2 figs.

Grassl, E. F. (2). Possible value of continuous feeding of medicated dry diets to prevent and control pathogens in hatchery reared trout. Progr. Fish Cult. 19:85–88.

Grave, E. V. Fisch-Schuppen unter dem Mikroskop. Mikrokosmos 46:184–188 5 figs.

Greenbank, J. Aggregational behavior in a freshwater sculpin. Copeia 1957: 157.

Greenwood, P. H. (1). A revision of the Lake Victoria Haplochromis species (Pisces, Cichlidae) Pt. II. H. sauvagei (Pfeffer), H. prodromus Trewavas, H. granti Blgr and H. xenognathus sp. n. Bull. Brit. Mus. (nat. Hist.) Zool. 5: 75-97 1 pl. 8 figs.

Greenwood, P. H. (2). Fish remains from the Mumba Cave, Lake Eyasi. in Lehmann U. Eine jungpleistozäne Wirbeltier fauna aus Ostafrika. Mitt. geol. (St.) Inst. Hamb. 26: 125–130.

Greenwood, P. H. (3). The fishes of Uganda—III & IV. Uganda J. 21: 47-80 22 figs. & 191-219 11 figs.

Greven, U. Wirbelsäulenverkrummung einer Plötze (*Leuciscus rutilus*). Gewässer u. Abwässer No. 1 1953: 70–76 9 figs.

Grey, M. New records of deep sea fishes including a new species, *Oneirodes bradburyae*, from the Gulf of Mexico. Copeia 1956 No. 4 [1957]: 242-246 2 figs.

Groenewald, A. A. v. J. The tiger-fish (*Hydrocyon vittatus*). Fauna & Flora, Pretoria, No. 6 1955:19-25 3 figs.

Gros. C. M. & Bloch, J. (1). Influence de la dose sur la survie d'un poisson, Carassius carassius (L.) après irradiation totale à 18°. C.R. Soc. Biol., Paris 151: 169-172 2 figs.

Gros, C. M. & Bloch, J. (2). Influence de la température sur la survie d'un poisson *Carassius carassius* (L.) après irradiation totale, C.R. Soc. Biol. Paris 151: 602-604.

Gross, W. Mundzähne und Hautzähne der Acanthodier und Arthrodiren. Palaeontographica A 109 1-2:1-406 pls. figs.

Guffroy, S. Notes Paléoichthyologiques. Bull. Soc. géol. Fr. (6) 6:847–854 2 pls. figs.

18

Guibé, J. & Spillmann, J. Au sujet de quelques types de Cyprinidés (Poissons) de Cuvier et Valenciennes. Bull. Mus. Hist. nat. Paris (2) 29: 462-463.

Gunter, G. Predominance of the young among marine fishes found in fresh water. Copeia 1957: 13-16.

Gyurko, S. & others. La reproduction artificielle du chevaine et du barbeau a Manastur-Cluj. Bul. Inst. Cercet. Pisc. Roman. 16 No. 2:46-52. [French summary.]

Hachey, H. B. see Leim, A. H.

Hagen, D. W. see Warburton, B.

Hajny, R. A. see Welander, A. D.

Hale, H. M. Der Fetzenfisch, ein seltsam esüdaustralische Seepferdehen. Natur u. Volk 87: 382–385 2 figs.

Hall, D. N. F. Recent developments in the Barbadian flying-fish fishery and contributions to the biology of the flying-fish, Fish. Publ. Lond. No. 7, 1956: 1-41.

Hallock, R. J. & others. The use of wire fyke traps to estimate the runs of adult salmon and steelhead in the Sacramento River. Calif. Fish Game 43:271-298, 10 figs.

Halstead, B. W. Weever stings and their medical management. U.S. Armed Forces Med. J. 8:1441-1451 8 figs.

Halstead, B. W. see Kawabata, T.

Hamada, K. A new osmerid fish, Hypomesus sakhalinus new species, obtained from Lake Taraika, Sakhalin, Jap. J. Ichthyol. 5:136-142 3 figs.

Hamon, M. Sur le trajet des fibres nerveuses optiques chez le poisson Cyprinodonte *Gambusia holbrooki* Gir. C.R. Acad. Sci. Paris **245**: 2396-2399 1 fig.

Haneveld, G. T. Injuries caused by garfish-like fishes in Netherlands New Guinea. Docum, Med. geogr. trop. 8 1956: 96-98.

Hansen, H. Let's go on an African safari (fish collecting, that is). Aquarium, Philad. 26: 129-133.

Hansler, D. D. see Donaldson, L. R.

Hardisty, M. W. Osmotic conditions during the embryonic and early larval life of the brook lamprey. (*Lampetra planeri*). J. exp. Biol. 34:237-252 3 figs.

Harrington, R. W. Sexual photoperiodicity of the Cyprinid fish, Notropis bifrenatus (Cope), in relation to the phases of its annual reproductive cycle. J. exp. Zool. 135: 529-556 l pl. 4 figs.

Harrison, H. M. Growth of the channel catfish, *Ictalurus punctatus* (Rafinesque), in some Iowa waters. Proc. Iowa Acad. Sci. **64**:657-666 3 figs.

Hartley, W. G. & Brayshaw, J. D. Movements of salmon off the North-East Coast of England, Fish. Invest. Lond. (1) 5 No. 8 1955: 1-17 2 figs.

Harvey, E. N. see Brown, M. E.

Hasan, G. Contributii la studiul musculaturii laterale la Stavrid (*Trachurus trachurus trachurus* L.). Bul. sti. Acad. Repub. rom. Sect. Biol., 3 1951: 703–714 3 figs. [French résumé.]

Hashitani, S. see Imamura, J.

Hasler, A. D. see Brown, M. E.

Haven, D. S. Distribution, growth, and availability of juvenile croaker. *Micropogon undulatus* in Virginia. Ecology 38: 88-97 8 figs.

Hayama, K. see Kubota, Z.

Hayama, K. see Syazuki, K.

Hayashi, S. & Suzuki, H. Growth of the Japanese anchovy. III. Vertebral counts of the post larvae. Bull. Tokai Fish. Res. Lab. 15: 21-28, 3 figs.

Hayes, F. R. On the variation in bottom fauna and fish yield in relation to trophic level and lake dimensions. J. Fish. Res. Bd. Can., 14: 1-32 6 figs.

Haysom, N.M. Notes on some Queensland fishes. Ichthyol. Notes Qd. 1: 139-144, 2 figs.

Hazelwood, A. Pisces. Naturalist No. 860: 10.

Healey, E. G. see Brown, M. E.

Healy, A. (1). A specimen of tench, *Tinca tinca*, L., from Lough Lein, Killarney. Irish Nats'. J. 12 1956: 109.

Healy, A. (2). Fishes of Lough Rea, Co. Galway, Ireland. II. Pike and rudd with general conclusions. Salm. Trout Mag. No. 151 1956: 246-252.

Healy, A. (3). Fishes of Lough Rea, Co. Galway, Ireland. I. Trout and perch. Salm. Trout. Mag. No. 150: 107-116.

Healy, A. (4). Brown trout of the Fergus system, County Clare, Ireland. Salm. Trout. Mag. No. 151:193-1982 figs.

Heck, L. Über den Gabelbart (Osteoglossum bicirrhosum Vandelli). Aquar. Terrar. Z. 10: 121-122, fig.

Hedgpeth, J. W. Treatise on marine ecology and paleoecology Vol. 1, Ecology. Mem. geol. Soc. Amer. 67: 1-1296, figs.

Heide, S. van der. In W. J. Jongmans & S. van der Heide. Flore et faune du Carbonifère inférieur de l'Égypte. Med. Geol. Sticht., N.S. 8 1955: 73-75 pl. xvii.

Hemens, J. see Alabaster, J. S.

Hemming, F. Proposed validation under the plenary powers of Osmerus (Class Actinopterygii) as of subgeneric status as from Linnaeus, 1758 (proposed validation of an erroneous entry on the Official List of Generic names in Zoology made by the ruling given in Opinion 77. Bull. zool. Nom. 11 1955: 281–282.

Hempel, G. Okologische Studien zum tagesrhythmischen Verhalten von Meeresfischen. Verh. dtsch. zool. Ges. 1956: 415-421 3 figs.

Hemphill, A. F. The southern redbelly dace, *Chrosomus erythrogaster* erythrogaster from the lower Mississippi river drainage. Copeia 1957: 53.

Hemphill, A. F. see Cooper, L. B.

Henderson, G. T. D. The distribution of young Gadus poutassou (Risso). Bull. mar. Ecol. 4:179-202 5 figs.

Hennemuth, R. C. An analysis of methods of sampling to determine the size composition of commercial landings of Yellowfin Tuna (Neothunnus macropterus) and Skipjack (Katsuwonus pelamis). Bull. Inter-Amer. Tuna Comm. 2:174-243.

Henriques da Silva, G. La Faune Miocène de l'Ile du Prince. Mem. Mus. min. geol. Univ. Coimbra 42 1956: 29-53, 7 pls.

Hepburn, R. L. see Pitt, T. K.

Herald, E. S. The reported emergence of fishes from California's subterranean waters. Calif. Fish Game 43: 235-237, 1 fig.

Herbert, D. W. M. see Alabaster, J. S.

Herbst, H. V. Bauchflossenlose Hechte. Gewässer u. Abwässer No. 8: 77 1 fig.

Hersey, J. B. see Johnson, H. R.

Heuts, M. J. see Meuwis, A. L.

Hibiya, T. see Oguri, M.

Hill, R. N. Victory over the shark. Nat. Hist. N.Y. 65 1956: 352-356.

Hoar, W. S. see Brown, M. E.

Hodgson, W. C. The herring and its fishery. London, (Routledge & Kegan Paul) pp. xiii + 197 9 pls. 24 figs.

Hoedeman, J. J. (1). Die bisher beschriebenen Formen und Arten der Gattung Rivulus Poey. Aquar. Terrar. Z. 7 1956: 199–202 8 figs.

Hoedeman, J. J. (2). Hyphessobrycon griemi, eine farbenprächtige Neuheit der bifasciatus—Gruppe aus Gojas. Aquar. Terrar. Z. 10:87-89 2 figs.

Hoedeman, J. J. (3). Le statut du genre *Barbus* Cuvier et Cloquet 1816. L'Aquar. et les poissons 6 No. 3 1956: 13-16.

Hoedeman, J. J. (4). Notes on the ichthyology of Surinam (Dutch Guiana) 4. Additional records of Siluriform fishes. Beaufortia 6 No. 71:147-160 31 figs.

Hoedeman, J. J. (5). The status of the Callichthyid genus Cascadura Ellis, 1913 (Pisces-Siluriformes-Callichthyidae). Bull. aquat. Biol. 1:15-19 2 figs.

Hoehne, K. Fischschuppen im Vitrit unterrotliegender Steinkohlenflöze von Stockheim im Oberfranken und Manebach in Thüringen. Geologie **6**:528– 540 8 pls.

Hoffmann, J. Notizen und Bemerkungen zu dem Begriff der Haxl (Squalius leuciscus L.). Bull. Soc. Nat. luxemb. 59 1955: 171-184 2 figs.

Holstvoogd, C. The post embryonic development of the pronephros in Clupea harengus L. Arch. néerl. Zool. 12 (3): 455-466, 1 pl. 7 figs.

Honma, Y. (1). [On the pituitary gland of a northern Japanese blenny, Stichaeus grigorjewi Herzenstein.] Jap. J. Ichthyol. 5:93-98 2 figs. [English summary.]

Honma, Y. (2). [On the pituitary gland of some Japanese elasmobranchiate fishes.] Jap. J. Ichthyol. 5: 107–113 12 figs. [English summary.]

Honma, Y. (3). [On the pituitary gland of some Japanese teleosts.] Jap. J. Ichthyol. 5:175-181 11 figs. [English summary.]

Honma, Y. (4). [A case of the Janus monstrosity found in the chum salmon embryo.] Jap. J. Ichthyol. 5:182 2 figs. [In Japanese.]

Horstmann, E. Familien-und Kernspezifische Synapsenformen bei Teleosteern. Progress in Neurobiology, London etc. 1956: 26–33 5 figs.

Houdemer, F. E. (1). Liste commentée des poissons de mer observés à Ajaccio. Bull. Soc. hist. Corse 1949:
—?. [Not seen.]

Houdemer, F. E. (2). Notes complémentaires sur quelques poissons observés en Corse. Cah. Nat. Bull. N.P. (N.S.) 8 1953: 83.

Huang, W. Early Pleistocene fish remains from Sanmen Area, Honan. Vertebrata Palasiatica 1:313-319 figs. [In Chinese with English summary p. 319.]

Hubbs, C. (1). Distributional patterns of Texas freshwater fishes. Southwestern Nat. 2:89-104 2 figs.

Hubbs, C. (2). A check-list of Texas fresh-water fishes. Texas Game Fish Comm. Ser. 1F No. 3: 1-11.

Hubbs, C. (3). Duration of sperm viability as a factor in frequency of fish hybridization. Texas J. Sci. 9: 472-474.

Hubbs, C. (4). Gambusia heterochir, a new Poeciliid fish from Texas, with an account of its hybridization with G. affinis. Tulane Stud. Zool. 5 No. 1: 1-16 7 figs.

Hubbs, C. & Brown, W. H. Dionda diaboli (Cyprinidae), a new minnow from Texas. Southwestern Nat. 1 1956 [1957]: 69-77 4 figs.

Hubbs, C. & Kemp, R. J. Three shiners, Notropis ortenburgeri, N. maculatus and N. blennius, added to the known East Texas fauna. Texas J. Sci. 8 1956: 110-112.

Hubbs, C. & Reynolds, R. A. Copulatory function of the modified pectoral fin of Gambusiin fishes. Amer. Nat. 91:333-335.

Hubbs, C. & Springer, V. G. A revision of the *Gambusia nobilis* species group, with descriptions of three new species, and notes on their variation, ecology and evolution. Texas J. Sci. 9: 279-327 15 figs.

Hubbs, C. & Strawn, K. (1). The effects of light and temperature on the fecundity of the greenthroat darter, Etheostoma lepidum. Ecology 38:596-602 2 figs.

Hubbs, C. & Strawn, K. (2). Relative variability of hybrids between the darters, *Etheostoma spectabile* and *Percina caprodes*. Evolution 11:1-10 2 figs.

Hubbs, C. & Strawn, K. (3). Survival of F1 hybrids between fishes of the subfamily Etheostominae. J. exp. Zool. 134:33-62.

Hubbs, C. see Warburton, B.

Humm, D. G., Clark, E. E. & Humm, J. H. Transplantation of melanomas from platyfish-swordtail hybrids into embryos of swordtails, platyfish and their hybrids. J. exp. Biol. 34:518–528 2 pls. 1 fig.

Huzum, I. V. Consideratiuni asupra unei diformatiuni la aripioarcle pectorale dela Pastruga si Nisetru. (Acipenser stellatus Pall.) si (Acipenser guldenstaedti Brandt). Bul. Soc. Nat. Român. No. 15 1941: 30-33 3 figs.

(Ignatyeva, G. M.) (1). [Morphophysiological studies of the hatching gland in *Huso huso* (L.). C.R. Acad. Sci. URSS 114:908-911 3 figs. [In Russian.]

(Ignatyeva, G. M.) (2). [Morphophysiological studies of the hatching gland in *Acipenser stellatus* at different incubation temperatures.] C.R. Acad. Sci. URSS 114: 1132-1135 2 figs. [In Russian.]

Ikehara, I. I. see King, J. E.

(Ilyin, B. S.) (1). [Remarks on and alterations to the Gobioidei in L. S. Berg's Freshwater fishes of the USSR and adjacent countries. Ed. 4 1948–1949 pp. 1055–1125.] Ichth. Prob. Moscow 7 1956: 185–192. [In Russian.]

(Ilyin, B. S.) (2). [Gobius (Macrogobius) paganellus in Black Sea.] Ichth. Probl. Moscow, 8:25-27 fig. [In Russian.]

Imamura, T. & Hashitani, S. On the food habits of four fishes in Marsh Hinuma. Bull. Fac. Lib. Arts. Ibaraki Univ. Nat. Sci. No. 7:45-56. [English summary.]

Inger, R. F. Report on a collection of marine fishes from North Borneo. Fieldiana, Zool. 36:341-405 2 figs.

Inger, R. F. see Woods, L. P.

(Irkimovich, A. I. & Zelenin, A. M.). [Histological changes occurring in the pituitary body in the process of sexual maturation of *Cyprinus carpio*.] C.R. Acad. Sci. URSS 114:655-657 pl. [In Russian.]

Ishida, J. Salvelinus in southern Saghalien. Zool. Mag., Tokyo 54 1942: 431-438 2 pls.

Ishida, J. & Taguchi, S. Effect of 2, 4-Dinitrophenol and sodium azide on lactic acid production during early development of the teleost *Oryzias latipes*. Annot. zool. jap. 30:1-7.

Ishii, S. Seasonal changes in the ovary and testis of the viviparous teleost, *Ditrema temmincki*. Annot. zool. Jap. 30: 204–210 8 figs

Ishii, S., & Egami, N. Effect of testosterone on dorsal fin of the file fish, *Monacanthus cirrhifer*. Annot. zool. Jap. 30: 77–82 13 figs.

Ito, T. see Yoshida, H.

(Ivanova, M. N.) [Some data on the food of perch in the Volga delta.] Ichth. Probl. Moscow 7 1956: 96-106 5 figs. [In Russian.]

(Ivanova, S. A.) [Histological study of gonads of gorbuscha and summer keta salmon.] Ichth. Probl. Moscow 6 1956: 96-99 3 pls. [In Russian.]

Ivasik, V. M. & Roi, P. E. Experiences pour enrager l'hydropisie de la carpe dans les fermes piscicoles des regions à l'ouest de la R.S.S. Ukraine. Bul. Inst. Cercet. Pisc. Roman. 16 No. 2:61-68. [French summary.]

(Izhevski, G. K.) [Methods and ways of developing research on exploration and behaviour of fishes.] Trud. Sovesh. Ikhtiol. Komm. 5 1953 [1955]: 8-11 [In Russian.]

Jakowska, S. see Nigrelli, R. F.

James, W. W. A further study of dentine. Trans. Zool. Soc. Lond. 29: 1-45 10 pls. 4 figs.

Jarvik, E. Crossopterygierna och de forsta fyrfota djuren. Svensk Faunistik Revy 1955 No. 3:80-96 7 figs.

Jászfalusi, L. Descrierea limnobiologică piscicolă a Muresculin intre hotarele Comunei Godea și a pâraielor din imprejunni. Notul. biol. București 5 1947: 287–321 3 pls, 2 figs, map.

Jayaram, K. C. Nomenclatural status of the names *Bagre* Cuvier (Oken), *Bagrus* Valenciennes and *Porcus* Geoffroy St. Hilaire. Copeia **1956** No. 4 [1957]: 248–249.

Jean, Y. Présence de larves de Sebastes marinus dans la Baie-des-Chaleurs et leurs caractères distinctifs. Nat. canad. 82 1955: 33-43 2 figs.

Jensen, P. T. An extension of the range of the long-finned smelt. Calif. Fish Game 43:99 1 fig.

Jhingran, V. G. Age determination of the Indian major carp *Cirrhina mrigala* (Ham.) by means of scales. Nature London 179: 468–469 text-figs. 1 and 2.

Jitariu, P. see Branzei, P.

Joensen, J. S. Merking av toski vestan fyri Suduroy i 1952. Fróskaparrit **5** 1956: 85–97 8 maps. [English summary.]

John, K. R. Observations on the behavior of blind and blinded fishes. Copeia 1957: 123-132, 4 figs.

Johnson, H. R., and others. Suspended echo sounder and camera studies of midwater sound scatterers. Deep Sea Res. 3 1956: 266–272.

Johnson, R. C. see Welander, A. D.

Johnson, W. E. On the distribution of young sockeye salmon (*Oncorhynchus nerka*) in Babine and Nilkitkwa Lakes, B.C. J. Fish. Res. Bd. Can. 13 1956: 695–708.

Jones, F. R. H. Rotation experiments with blind goldfish. J. exp. Biol. 34: 259-275 8 figs.

Jones, F. R. H. see Brown, M. E.

Jones, R. W. Char. Fishing Gazette (Beckenham, England) 139:1122.

Jónsson, J. Er Ysan stadbundin i Faxaflóa? Aegir **49** 1956 : [1–2.] [English summary.]

22

Jubb, R. A. The freshwater eel Anguilla obscura Gunther, 1871, in Southern Africa. Nature, Lond. 180: 1216.

Judefind, T. F. see Kawabata, T.

Julin, E. Vanliga havsnålens förekomst i Bottenhavet. Fauna och Flora No. 3 1956: 140–144 2 figs.

June, F. C. see Yuen, H. S. H.

Junker, M. Vitamin A in Fischen, Krebsen und Muscheln. Arch. Fisch. Wiss. 7: 248-260.

Kafuku, T. [Ecological and phyletic significance of anal fin between cyprinids and catostomid fish.] Jap. J. Ichthyol. 5:163-173. 6 figs. [English summary.]

Kähling, J. see Thines, G.

Kähsbauer, P. Viktor Pietschmann. Ann. naturh. Mus. Wien 61: 1-3 ports.

Kakimoto, D. [Studies on B-vitamins of pyloric coeca of Skipjack, Katsuwonus vagans IV. On vitamin B12.] Bull. Jap. Soc. Sci. Fish. 22:634-636. [English summary].

Kakimoto, D., & others. [Amino acid composition of the pyloric coeca of skipjack, *Katswonus vagans*.] Bull. Jap. Soc. Sci. Fish. **22**:631-633. [English summary.]

Kamer, J. C. van de. The pineal organ in fish and amphibia. Progress in Neurobiology, London, N.Y. 1956: 113-120 5 figs.

Kamohara, T. (1). On the fishes of the family Chlorophthalmidae. Res. Rep. Kôchi Univ. 5 No. 15 1956: 1-8 3 pls. [English summary.]

Kamohara, T. (2). Notes on twenty additions to the marine fish fauna of Prov. Tosa, Japan, including one new genus (family Peristediidae). Res. Rep. Kôchi. Univ. 6 No. 5: 1-61 fig.

Kamohara, T. (3). List of fishes from Amami-Oshima and adjacent regions, Kagashima Prefecture, Japan. Rep. Usa mar. Biol. Sta. 4: 1-65 38 figs.

Kampa, E. M. & Boden, B. P. Light generation in a sonic-scattering layer. Deep Sea Res. 4:73-92 8 figs. (Kanaeva, I. P.) [Daily changes in feeding of *Percarina demidoffi maeotica*.] Ichth. Probl., Moscow **7** 1956: 52–59 8 fig. [In Russian.]

Kanazawa, A. see Kakimoto, D.

Kanazawa, R. H. A new species of eel, Coloconger meadi, and new records for the ateleopid fish, Ijimaia antillarum Howell Rivero, both from the Gulf of Mexico. Copeia 1957: 234-235 1 fig.

Kanoh, Y. & Yamamoto, T. S. Removal of the membrane of the dog salmon egg by means of proteolytic enzymes. Bull. Jap. Soc. sci. Fish. 23: 166-172, 3 figs.

Kanwisher, J. & Ebeling, A. Composition of the swim-bladder gas in bathypelagic fishes. Deep Sea Res. 4: 211-217 2 figs.

Kao, C. Y., Chambers, R. & E. L. Internal hydrostatic pressure of the Fundulus egg. II. Permeability of the chorion. J. cell. comp. Physiol. 44 1954: 447-461 2 figs.

Kapoor, A. S. Sensory canals of the head in *Wallagonia attu* (Day). Experientia 13:459-460 1 fig.

Kapoor, B. G. (1). Secretory cells in the gills of an Indian fresh-water spiny eel, *Mastacembelus armatus* (Lacép.). Jap. J. Ichthyol. 5:123-126 1 fig.

Kapoor, B. G. (2). Oral valves of teleosts. Jap. J. Ichthyol. 5:127-131 7 figs.

Kapoor, B. G. (3). Pharyngeal pockets of *Gadusia chapra* (Ham). Jap. J. Ichthyol. 5:132–135 2 figs.

Kapoor, B. G. (4). The callous pad of the pharyngeal masticatory apparatus in Indian carps. Jap. J. Ichthyol. 6: 26-30 2 pls.

Kapoor, B. G. see Sarkar, H. L.

Kappers, J. A. On the problem of the presence of a paraphysis cerebri in selachians. Pubbl. Staz. zool. Napoli 29:41-70 4 pls.

Karapetkova, M. S. see Martino, C. V.

(Karpevich, A. F.) [Method of prognosis of food level and conditions of fish feeding in changing water reservoir conditions.] Trud. Sovesh. Ikhtiol. Komm. 5 1954 (1955): 85–92. [In Russian.]

(Karzinkin, G. S. & Krivobok, M. N.) [Method of studying the physiology of nutrition and food requirements of fish.] Trud. Sovesh. Ikhtiol. Komm. 6 1954 (1955): 40-53 6 figs. [In Russian.]

Kasahara, S. Notes on the young of the Black Porgy, Mylio latus (Houttuin). Jap. J. Ichthyol. 6:20-25 3 figs. [English summary.]

Kashiwada, K. see Kakimoto, D.

Kawabata, T., Halstead, B. W. and Judefind, T. F. A report of a series of recent outbreaks of unusual cephalopod and fish intoxications in Japan. Amer. J. trop. Med. Hygiene 6: 935-939.

Kemp, R. J. Occurrence of the ocean sunfish *Mola mola* (Linnaeus) in Texas. Copeia **1957**: 250–251.

Kemp, R. J. see Hubbs, C.

Kennedy, H. D. see Nielson, R. S.

Kessler, R. Heredozelluläre Strahlenschäden im Hoden des Zahnkarpfens Lebistes reticulatus. Naturwissenschaften 44: 19–20 2 figs.

Ketchen, K. S. Factors influencing the survival of the lemon sole (*Parophrys vetulus*) in Hecate Strait, British Columbia. J. Fish. Res. Bd. Can. **13** 1956: 647-694.

Keynes, R. D. see Brown, M. E.

Khabakov, A. V. (1). Pisces section in Atlas of the leading forms of the fossil fauna of the USSR, Vol. 4, Lower Carboniferous, 1941. [Omitted from previous record.]

[Khabakov, A. V.] (2). [O dvukh novuikh formakh problematicheskikh iskopaemuikh *Crookallia* iz karbona donetzkogho bassina i *Palaeoxyris* iz yurui Kruima.] Ann. Soc. paléont russe 13 1949: 83–87 figs.

Kilby, J. D. & Caldwell, D. K. A list of fishes from the southern tip of the Florida Peninsula. Quart. J. Fla. Acad. Sci. 18 1955: 195-206 1 fig.

Kimsey, J. B. (1). The status of the redeye bass in California. Calif. Fish Game 43:99-100.

Kimsey, J. B. (2). Large mouth bass tagging at Clear Lake, Lake County, California. Calif. Fish Game 43:111-118. King, J. E. & Ikehara, I. I. Comparative study of food of big-eye and yellow-fin tuna in the central Pacific. Fish. Bull. U.S. 57 1956: 59-85.

(Kirpichnikov, V. S., Svetovidov, A. N. & Troshin, A. S.) [The labelling of *Cyprinus carpio* with phosphorous and calcium radioactive isotopes.] C.R. Acad. Sci. URSS 111 1956: 221-224 4 graphs. [In Russian.]

(Kirshenblat, Y. D.) [The action of purified hormones of the pituitary body upon fish ovary.] C.R. Acad. Sci. URSS 111 1956: 504-506.] [In Russian.]

Klausewitz, W. (1). Botia modesta, Botia lecontei und Botia horae. Aquar. Terrar. Z. 10: 33-35, 3 text-figs.

Klausewitz, W. (2). Roloff's Zwergbarbe (*Puntius roloffi*), eine neue Art. Aquar. Terrar. Z. 10:89-90 1 fig.

Klausewitz, W. (3). Eine neue Kugelfisch-Art aus Siam: Tetraodon somphongsi. Aquar. Terrar. Z. 10: 115-116 figs. 1-2.

Klausewitz, W. (4). Nachtrag zu Botia modesta und lecontei. Aquar. Terrar. Z. 10:184.

Klausewitz, W. (5). Die Lohachata-Schmerle (*Botia lohachata*) eine interessante Neueinführung aus Pakistan. Aquar. Terrar. Z. 10:201–203, text-figs. 1–5.

Klausewitz, W. (6). Feuerfische der Gattungen Dendrochirus und Pterois. Aquar. Terrar. Z. 10:319-323 2 figs.

Klausewitz, W. (7). Anisochaetodon (Linophora) pictus Forskål eine gute Art der Borstenzähner (Pisces, Chaetodontidae.) Bull. aquat. Biol. 1:1-12 2 figs.

Klausewitz, W. (8). Über einige Fische aus Thailand. Senckenbergiana, Biol. 38: 33-39 4 figs.

Klausewitz, W. (9). Neue Süsswasserfische aus Thailand. Senckenbergiana, Biol. 38:193–204 3 pls. 5 figs.

Klausewitz, W. (10). Tetraodon somphongsi n.sp. ein weiterer neuer Kugelfisch aus Thailand (Pisces, Tetraodontidae). Senckenbergiana, Biol. 38: 205–208 2 figs. Klausewitz, W. (11). Barbus schneemanni und Aplocheilichthys maculatus, zwei neue Fische aus Ost-Afrika (Pisces, Cyprinidae und Cyprinodontidae.) Senckenbergiana Biol. 38: 279–282 2 figs.

Kleerekoper, H. & Sibakin, K. An investigation of the electrical "spike" potentials produced by the sea lamprey (*Petromyzon marinus*) in the water surrounding the head region. II. J. Fish. Res. Bd. Can., 14: 145-151 3 figs.

Klyuchareva, O. A.) [On some problems regarding intraspecific relationships in fishes.] Zool. Zh., Moscow 35 1956: 275–289. [In Russian.]

Koefoed, E. Notes on the Greenland shark Acanthorhinus carcharias (Gunn.) 2. A uterine foetus and the uterus from a Greenland shark. Fiskeridir, Skr. Havundersøk. 11 No. 10:8-12 5 figs.

Koenig, O. Erfahrungen mit Korallenfischen. Aquar. Terrar. Z. 10: 154-156.

(Kokhanova, N. A.). (Development of *Cobitis taenia*.] Ichthyol. Problems, Moscow 8: 89–101 4 figs. [In Russian.]

(Kokhiya, A. B.) (1). [Food of some fishes in Khramsk water reservoir and adjoining rivers.] Trud. Inst. Zool. Tiflis 13 1954:179-195. [Gruzian with Russian summary.]

(Kokhiya, A. B.) (2). [The food of fish in the basin of the river Rioni.] Trud. Inst. Zool. Tiflis 15 1956: 261–287. [In Gruzian with Russian summary.]

Koli, L. Kirjoeväsimpeon (Cottus poecilopus Heck. esüntymisestä Suomessa, Luonnon Tutkuja 60:152–154 2 figs.

(Konstantinov, V. G.) [Comparative morphology and biology of *Perca fluviatilis*, *Perca schrenki* and *Lucioperca lucioperca* in different stages of development.] Trud. Inst. Morf. Zhiv. **16**: 181–236 10 figs. [In Russian.]

Konta, J. Crystal orientation of two phosphates in the scales of Permian Fish of the group Palaeoniscidae. Act. Geol., Budapest, 4:1956:113-121 6 text-figs.

(Koryakov, E. A.) [Certain ecological adaptations in the reproduction of Comephoridae.] C.R. Acad. Sci. URSS 111 1956: 1111-1114 2 graphs. [In Russian.]

Koster, W. J. Guide to the fishes of New Mexico. Albuquerque (University of New Mexico Press), pp. vii + 116: text-figs.

Kosto, B. see Pickford, G. E.

Kotthaus, A. Die deutschen Schollenuntersuchungen von 1948–1954. Ber. dtsch. Komm. Meeresforsch. 14 1956: 83–108 11 figs.

Kotthaus, A. & Krefft, G. Fischfaunenliste der Fahrten mit F.F.S. "Anton Dohrn" nach Island-Grönland. Ber. Dtsch. Wiss. Komm. Meeresforsch. 14:169–191.

(Kozhin, N. I.) [Fish-fauna of the Iranian shores of the Caspian Sea.] Ichthyol. Problems, Moscow 8:8-18. [In Russian.]

(Kozlov, V. V.) [The asphyxiation of fishes in Oka river.] Zool. Zh., Moscow 35 1956: 936-937. [In Russian.]

[Kozlovski, D. A.) [On the migratory instinct in fishes.] Zool. Zh., Moscow 35 1956: 266-274. [In Russian.]

Krayukhin, B. V. see Bodrova, N. V.

Krebs, W. see Sauerteig, E.

Krefft, G. Ein Beitrag zur Kenntnis des Gronlandrochens, *Raja spinicauda* Jensen 1914. Arch. Fisch. Wiss. **7** 1956: 139–145 2 figs.

Krefft, G. see Kotthaus, A.

Kristensen, I. Koornaarvissen (Atherina presbyter) langs en Nederlandse kust. Levende Nat. 60: 159-162 2 figs.

Krivobok, M. N. see Karzinkin, G. S.

(Krizhanovski, S. G.) (1). [Development of herring in water of high salinity]. Ichthyol. Problems Moscow 6 1956: 100–104 2 figs. [In Russian.]

(Krizhanovski, S. G.) (2). [Parthenogenetic development in Sakhalin herring, found in the spawning material of 1955.] Ichthyol. Problems, Moscow 8:81-88. [In Russian.]

(Krogius, F. V. & Krokhin, E. M.) [Results of biological investigations on red-nerka salmon, and the fluctuations of stock in Kamtchatka waters.] Ichthyol. Problems, Moscow 7 1956; 3-20 5 figs. [In Russian.]

Krokhin, E. M. see Krogius, F. V.

Kruckow, T. Die stratigraphische und paläogeographische Bedeutung der miozänen Elasmobranchier-Fauna von Baja California, Mexico. Neues Jb. Geol. Paläont. B. 1957: 444–449.

(Kruglova, V. M.) [Feeding grounds of bottom feeding fishes in Veselovsk reservoir and the reason for slow growth of carp.] Ichthyol. Problems, Moscow 8:116-128. [In Russian.]

Krumholz, L. A. (1). A record of the night shark, *Hypoprion signatus*, from Bimini, Bahamas. Copeia 1957: 300–301.

Krumholz, L. A. (2), Measurements of a large sharp-nose mackerel shark, *Isurus oxyrinchus*, from Bimini, Bahamas, Copeia 1957: 302.

Kubota, S. S. see Okada, Y.

Kubota, Z., & others. [Influence of pH and agricultural chemicals on the loach, Misgurnus anguillicaudatus (Cantor)].
J. Shimonoseki Coll. Fish. 6: 327-336. [English summary.]

[Kudrin, L. N.] [O nakhodkakh zubov akul v otlozheniyakh miotzena yugho-zapadnoi okrainui russkoi platformui.] Geol. Sbornik 4:349.

Kuenzer, P. Brutpflege beider Elterntiere bei Nannacara anomala. Aquar. Terrar. Z. 10:175–177 textfigs. 1-4,

Kulczycki, J. Upper Devonian fishes from the Holy Cross Mountains (Poland). Acta palaeont. polon. 2:285–382 13 pls. figs.

Kurazhskovski, Y. N. see Pidoplichko, I. G.

(Kurilov, S. V.). [The sturgeons in Pechora river.] Priroda 3:116-117. [In Russian.]

Kuriyama, Z. see Kubota, Z.

Kuroda, N. (1). [On the life colors of some fishes. V.] Jap. J. Ichthyol. 5:118-122 2 figs. [English summary.]

Kuroda, N. (2). Additions and corrections to the fishes of Suruga Bay (XIII). Zool. Mag., Tokyo 66:1-2. [English summary.]

Kuroda, N. (3). A list of the fishes of Nabuto, Boshu, Japan. Zool. Mag., Tokyo 66: 3-5 2 figs. [English summary.]

Kuronuma, K. & Yamaguchi, M. Descriptions of a new iniomid fish from Choshi, Japan. Zool. Mag., Tokyo 53 1941: 272-274.

Kusa, M. Osmotic behaviour of the isolated cortical alveoli of stickleback eggs. Annot. Zool. Jap. 30:67-70 l fig.

Kuthalingam, M. D. K. Observations on the food and feeding habits of the Indian Mackerel Rastrelliger kanagurta (Russel). J. zool. Soc. India 8 1956: 99-106.

Kutkuhn, J. H. Food and feeding habits of some fishes in a dredged Iowa lake. Proc. Iowa Acad. Sci. **62** 1955: 576–588.

Kuwabara, S. Acetylcholine (ACl) cycle in the electric organ of fish. J. Shimonoseki Coll. Fish. 6: 149-240 12 figs.

(Kuznetzov, V. V.). [On the so called phenomenon of Lee.] Ichthyol. Problems, Moscow 8:143-154. [In Russian.]

Ladiges, W. Über einige Farbabweichungen bei Fischen. Aquar. Terrar. Z. 10:153-156 text-figs. 1-3.

La Faunce, D. A. see Hallock, R. J.

Lagler, K. F. & Steinmetz, C. Characteristics and fertility of experimentally produced sunfish hybrids, *Lepomis gibbosus* × *L. macrochirus*. Copeia 1957: 290–292.

Lagunov, I. I. see Azbelev, V. V.

Lambert, J. G. Sur deux Electridae d'eau douce de Guinée Française, avec une révision du genre *Kribia*, Herre. Rev. Zool. Bot. afr., **56**:101-110 2 figs.

Lange, N. O. see Vasnetzov, V. V.

Lanzing, W. J. R. Biologie van de Petromyzonidae. Vakbl. Biol. 37: 133-142 1 fig.

Lapparent, A. F. de. Étude paléontologique des vertébrés du Jurassique d'El Mers (Moyen Atlas). Notes Serv. Min. Maroc 124 1955: 1–36, 5 pls. figs. [Pisces p. 10.]

Larimore, R. W. Ecological life history of the warmouth (Centrarchidae). Bull. Ill. nat. Hist. Surv. 27: 1–83 l pl. 27 figs.

Larkin, P. A., Terpenning, J. G. and Parker, R. R. Size as a determinant of growth rate in rainbow trout, Salmo gairdneri. Trans. Amer. Fish. Soc. 86 (1956) 1957: 84-96, 19 text-figs., 3 tables.

Lauer, C. Y. see Prosser, C. L. Laurent, P. see Couteaux, R.

Lauzier, L. see Leim, A. H.

Lawrence, J. M. A trapping experiment to estimate the bluegill population in a farm pond. Proc. Iowa Acad. Sci. 59 1952: 475-479.

Leary, T. R. (1). An occurrence of the parrotfish, *Sparisoma radians*, in a Texas bay. Copeia 1956 No. 4 [1957]: 249-250.

Leary, T. R. (2). The bonefish, Allula vulpes in Texas. Copeia 1957: 248-249.

Lecal, J. Recherches expérimentales sur les blennidés de Castiglione. Bull. biol. 91:115-157.

Le Danois, E. & others. Fishes of the world, London (G. G. Harrap) pp. 190 15 pls. col., text illust.

Lee, A. British fishery research in the Barents Sea. Polar Rec. 8 1956: 109–117 3 figs.

Legaspi, V. A. A contribution to the life history of the nomeid fish *Psenes cyanophrys* Cuvier and Valenciennes. Bull. mar. Sci. Gulf & Caribbean 6 1956: 179–199 5 figs.

Legeza, M. I. see Lindberg, G. U.

Leghissa, S. Contribution ultérieure à une meilleure connaissance de l'appareil de Mauthner chez les poissons et observations sur la morphologie de la fibre. Progress in Neurobiology, London etc. 1956: 45–62 6 figs.

Lehman, J.-P. (1). L'Evolution des Dipneustes et l'origine des Urodèles. Colloques Internat., Centre nat. Recherches Sci., Paris, 60 1956: 69-76.

Lehman, J.-P. (2). Complements à l'étude des genres *Ecrinesomus* et *Bobasatrania* de l'Eotrias de Madagascar. Ann. Paléont., Paris 42 1957 (1956): 65–94 6 text-figs pls. 5–13.

Lehman, J.-P. (3). Un problème non résolu: l'origine des Vertebrés. La Nature No. 3265: 174-7 4 figs. Leim, A. H. & others. Report of the Atlantic herring Investigation Committee. Bull. Fish. Res. Bd. Can. No. 111: 1-317.

Leiner, M. Stichlingsbastarde. Natur u. Volk 87: 294-300 5 figs.

Lemasson, M. J. Second symposium on African Hydrobiology and inland fisheries. C.S.A. Brazzaville No. 25 1956: 1-268 tabs.

Leonte, V. Contributions à l'étude de la biologie des esturgeons marins de la république populaire Roumaine. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 167-185.

Leonte, V. see Teodorescu-Leonte, R. Le Touzé, S. see Barents, A.

Lévèque, R. Notes sur la faune ichthyologique de Camargue. Terre et la Vie 104: 231-240 map.

Lieder, U. Die Ergebnisse der im Jahre 1956 durchgeführten Karpfen-Karauschenkreuzungen. 1. Beitrag zur Anwendung der 'entfernten' Hybridisation in der Fischzüchtung. Z. Fisch. 6: 283–299, 7 figs.

Lillelund, K. & Tiews, K. Die Dokumentation in der Fischereiliteratur und die Möglichkeit ihrer praktischen Verwirklichung. Arch. Fisch. Wiss. 7:129-138.

(Lindberg, G. U. & Legeza, M. I.) [On the two forms of the spiny dogfish Squalus acanthias L.] Zool. Zh., Moscow 35 1956: 1685–1688 2 figs. [Russian with English summary.]

Lindroth, A. (1) A study of the whitefish (Coregonus) of the Sundsvall Bay district. Rep. Inst. Freshwater Res. Drottningholm, No. 38:70-108 8 figs.

Lindroth, A. (2). Baltic salmon fluctuations: a reply. Rep. Inst. Freshwater Res. Drottningholm, No. 38:109-130 3 figs.

Lindsey, C. C. Possible effects of water diversions on fish distribution in British Columbia. J. Fish. Res. Bd. Can. 14:651-668 2 figs.

Liu, H. T. (1). Fossil fishes from Locality 14 of Choukontien. Palaeont. sinica N.S.C. 14, 1954: 1-21 8 pls. figs.

[Liu, H.-T.] (2). [A new fossil Cyprinid fish from Maoming, Kwangtung.] Vertebrata Palasiatica 1:151-153 1 pl. [English summary p. 153.]

Liu, H. & Yeh, S. Two new species of *Ceratodus* from Szechuan, China. Vertebrata Palasiatica 1:305-311 1 pl. [In Chinese with English summary p. 310.]

Liu, T.S. A new Cretaceous Palaeoniscoid fish from the Yumen of the Chiuchuan Basin, Western Kansu. Vertebrate Palasiatica 1:103-122 pls. i-vi 2 figs.

(Lobintzev, K. S.) [A contribution to the problem of tonic and untonic muscle fibres.] C.R. Acad. Sci. U.R.S.S. 112:1116-1118 2 figs. [In Russian.]

Locke, A. Beyond the National Park. Angling in Malaya 4 1956: 61-67.

Lockley, A. S. Adrenal cortical hormones and osmotic stress in three species of fishes. Copeia 1957: 241-242.

Lofts, B. & Marshall, A. J. Cyclical changes in the distribution of the testis lipids of a teleost fish, Esox lucius. Quart. J. micr. Sci. 98:79-88 2 pls. 1 fig.

Lopez, R. B. Pez aguja Scomberesox saurus (Walbaum) pescado en Necochea. Notas Mus. Univ. nac. La Plata 19: 145–151 8 figs.

Lopez, R. B. see Carlo, J. M.

Love, R. M. see Brown, M. E.

Lowe, R. H. Observations on the diagnosis and biology of *Tilapia leucosticta* Trewavas in East Africa. Rev. Zool. Bot. afr. 55: 353-373.

Lowenstein, O. see Brown, M. E.

Lubet, P. see Bauchot, M. L.

Lühmann, M. & Mann, H. Beobachtungen über die Blumenkohlkrankheit der Aale. Arch. Fisch. Wiss. 7:229–239 9 figs.

Lüling, K. H. Caecobarbus geertsi Boulenger—seine Entdeckung, Verbreitung und Lebensweise—und nochmals Anoptichthys jordani Hubbs und Innes. Aquar. Terrar. Z. 10:145–150 text-figs. 1–5.

Luscan, S. see Busnita, T.

Lyall, A. H. (1). The growth of the trout retina. Quart. J. micr. Sci. 98: 101-110 2 pls. 7 figs.

Lyall, A. H. (2). Cone arrangements in teleost retinae. Quart. J. micr. Sci. 98: 189–201 8 figs.

(Lyubitskaya, A. I.) [Influence of different parts of the visible spectrum region on the developmental stages of embryos and larvae of fish.] Zool. Zh., Moscow 35 1956: 1873–1886 6 figs. [Russian with English summary.]

McMynn, R. G. see Vernon, E. H.

Maes, H. see Bont, A. F. de.

Magnússon, J. (1). Mikroskopischanatomische Untersuchungen zur Fortpflanzungsbiologie des Rotbarsches (Sebastes marinus Linné). Z. Zellforsch. 43 1955: 121-167 24 figs.

Magnússon, J. (2). Fiskimidaleit med b. v. Fylki. Aegir 49 1956: 235– 240 3 figs. [English summary.]

Magnússon, J. (3). Fiskimidaleit med b. v. Fylki. Aegir 49 1956: 281– 283 1 fig. [English summary.]

(Makkoveeva, I. I.) [The food of young pike in Rybinsk reservoir.) Ichthyol. Problems Moscow 7 1956: 60-95 12 figs. [In Russian.]

Maldacea, L., Citterio, P. & Ranzi, S. Sulla presenza di actina e miosina nell' organo elettrico della torpedine. R.C. Ist. lombardo Cl. Sci. mat. e nat. 90 1956: 512

Maldacea, L. see Citterio, P.

Malyarevskaya, A. Y. see Bodrova, N. V.

Manacop, P. R. see Capco, S. R.

Manea, G. I. & Mihai, M. Premières expériences de reproduction artificielle des Esturgeons de la Roumaine (Résultats préliminaires obtenus en printemps 1955 et 1956). Bul. Inst. Cercet. Pisc. Roman. 16 (1): 57–72.

Manea, G. see Busnita, T.

Mann, H. see Lühmann, M.

Marathe, V. B. & Bal, D. V. The chondrocranium of *Eleutheronema tetradactylum* (Shaw). J. zool. Soc. India 8 1956: 107-118 7 figs.

Marine Biological Association of the U.K. Plymouth Marine Fauna, Edit. 3, Plymouth (Mar. Biol. Assoc.) 1957.

Marshall, A. J. see Lofts, B.

Marshall, J. R. The lateral musculature and the swimming of fish. Proc. zool. Soc. Lond. 126 1956:127-143 9 figs.

Marshall, N. B. (1). Evolutionary aspects of fish classification. Baer, J. G. First Symposium on host specificity among parasites of vertebrates, Neuchatel Univ.,: 173-184 3 figs.

Marshall, N. B. (2). Tiefseebiologie, Jena (Gustav Fischer) pp. viii + 334, 5 pls. col., 103 figs.

Marshall, T. C. Ichthyological Notes. Ichthyol. Notes Qd. 1:117-137 1 pl.

Martino, C. V. & Karapetkova, M. S. Étude quantitative des relations nutritives chez les poissons. Zool. Zh., Moscow 36:425-431. [French summary.]

Maruyama, K. Rare 'deep-water fishes from off Tohoku and adjacent regions. II. Record of a rare species of the genus *Melamphaes* (family Melamphaidae). Jap. J. Ichthyol. 6: 31-34 l fig. [English summary.]

Massmann, W. H. New and recent records for fishes in Chesapeake Bay. Copeia 1957: 156-157.

Massonet, R. see Grangaud, R.

Massuti, M. O. Biologia de la sardina de Baleares. Bol. Inst. esp. Oceanogr. No. 80 1956: 1–21 14 figs.

Matei, D. see Busnita, T.

Mather, F. J. & Gibbs, R. H. Distributional records of fishes from waters off New England and the middle Atlantic states. Copeia 1957: 242-244.

Matta, F. L'accrescimento del merluzzo del Mediterraneo. Boll. Pesca Piscic. Idrobiol. 32 fasc. 1 1956 [1957]: 31–35.

Maul, G. E. Further additions to the previously revised family Searsidae. Bol. Mus. Funchal No. 10:5-21 5 figs.

Maurice, A. see Cordier, D.

Maxunov, V. A. Certain data on morphology and biology of *Capoeto*brama kuschakewitschi (Kessler) of the river Syr-Darva. Zool. Zh., Moscow 36:238-245. [English summary.]

Mayhew, J. (1). The bluegill, Lepomis macrochirus (Rafinesque) in West Okoboji Lake, Iowa. Proc. Iowa Acad. Sci. 63 1956: 705-713 2 figs.

Mayhew, J. (2). The occurrence of blindness in the black bullhead, *Ictalurus melas* (Rafinesque), of East Okoboji Lake, Iowa. Proc. Iowa Acad. Sci. **64**: 654-656.

Mazumdar, S. R. On the spawning habits and early development of Wallagonia attu (Bloch and Schneider). Proc. zool. Soc. India 10:99-104 1 pl.

Mead, G. W. (1). An Atlantic record of the zeoid fish *Parazen pacificus*. Copeia 1957: 235–237 1 pl.

Mead, G. W. (2). On the bramid fishes of the Gulf of Mexico. Zoologica, N.Y. 42: 51-61 3 pl.

Mead, G. W. see Bohlke, J.

Medcof, J. C. Nuptial or pre-nuptial behaviour of the shad, *Alosa sapidissima* (Wilson). Copeia 1957: 252–253.

Medcof, J. C. see Canagaratnam, P.

Medcof, J. C. see Sivalingam, S.

(Mednikov, B. M. & Prokhorov, V. G.) [A new species of Cyclopteropsis (Pisces Cyclopterinae) from the Bering Sea.] C.R. Acad. Sci. URSS 111 1956: 717-719 fig. [In Russian.]

Meinken, H. (1). Über zwei der Liebhaberei bislang unbekannte Corydoras-Neuheiten (Callichthyidae—Ostariophysi). Aquar. Terrar. Z. 10:4-7 1 fig.

Meinken, H. (2). Rasbora nigromarginata n. spec., die "Rote Rasbora". Aquar. Terrar. Z. 10: 65-68 2 figs.

Meinken, H. (3). Moenkhausia sanctae filomenae (Steindachner) 1907, die Rotaugen-Moenkhausia. Aquar. Terrar. Z. 10: 141–143, fig.

Meinken, H. (4). Wieder einmal die Barbus—Puntius—Frage. Aquar. Terrar. Z. 10: 180–184, text-figs. 1–3.

(Mekhanik, F. Y.). [The growth and metabolism of the fry of Ladoga salmon and trout at various temperatures.] Zool. Zh., Moscow 35 1956: 290-299 2 graphs. [In Russian.]

(Melnikov, G. B.) [Trial of introducing bream in Khristoforov water reservoir (Kpivorozhye).] Trud. Ghidrobiol. Obshch. 7 1956: 209–215. [In Russian.]

Menzel, H.-U. & Steffens, W. Über das Wachstum von Spiegel und Nachtkarpfen. Z. Fisch. 6:385–392 4 figs.

Merkel, T. J. Food habits of the king salmon, Oncorhynchus tshawytscha (Walbaum), in the vicinity of San Francisco, California. Calif. Fish Game 43: 249-270 4 figs.

Merkens, J. C. & Downing, K. M. The effect of tension of dissolved oxygen on the toxicity of un-ionized ammonia to several species of fish. Ann. appl. Biol. 45: 521-527 4 figs.

Merkens, J. C. see Downing, K. M.

Meuwis, A. L. & Heuts, M. J. Temperature dependence of breathing rate in carp. Biol. Bull. Woods Hole 112: 97-107.

Meyer-Waarden, P. F. & Rübenberg, L. Untersuchungen über den verwertbaren Fleischanteil und Fettgehalt bei Ostseeflundern. Arch. Fisch. Wiss. 7:240-247.

Miegel, H. (1). Fischerei und Fische in den Trinkwassertalsperren im Einzugsgebiet der Wupper. Gewässer u. Abwässer No. 7 1955: 39-71 2 maps.

Miegel, H. (2). Bescreibung der Talsperrenform eines Cypriniden aus der Abramisgruppe. Gewässer u. Abwässer No. 13 1956: 7–22 13 figs.

Migdalski, E. C. Contribution to the life history of the South American fish Arapaima gigas. Copeia 1957: 54-56.

Mihai, M. see Manea, G. I.

Miller, H. W. Petalodus jewetti, a new species of fossil Bradyodont fish from Kansas. Trans. Kansas Acad. Sci. 60:82-85 2 figs.

Miller, H. W. and Swineford, A. Paleoecology of nodulose zone at top of Haskell Limestone (Upper Pennsylvanian) in Kansas. Bull. Amer. Ass. Petrol. Geol. 41: 2012–2036 3 pls. figs.

Miller, R. B. (1). Permanence and size of home territory in stream-dwelling cutthroat trout. J. Fish. Res. Bd. Can., 14:687-691.

Miller, R. B. (2). Have the genetic patterns of fishes been altered by introductions or by selective fishing? J. Fish. Res. Bd. Canada 14:797-806.

Miller, R. R. (1). The scientific name of the Australian lungfish. Turtox News 32 1954:69.

Miller, R. R. (2), Fishes from Laguna Ocotal. Bull. Mus. Comp. Zool. Harv. 116: 238-241.

Miller, R. R. (3). Utilization of X-rays as a tool in systematic zoology. Syst. Zool. 6:29-40.

Miller, R. R. (4). Origin and dispersal of the alewife, Alosa pseudoharengus, and the gizzard shad, Dorosoma cepedianum, in the Great Lakes. Trans. Amer. Fish. Soc. 86 (1956): 97-111.

Miller, R. R. see Robins, C. R.

Millot, J. see Le Danois, E.

Milne, D. J. Recent British Columbia spring and coho salmon tagging experiments, and a comparison with those conducted from 1925 to 1930. Bull. Fish. Res. Bd. Can. No. 113: 1-56 13 figs.

(Milokhin, A. A.) (1). [Relationship between the structure of the digestive tract nervous system and the kind of food in some Cyprinidae.] C.R. Acad. Sci. URSS 105 1955:182-184 4 figs. [In Russian.]

(Milokhin, A. A.) (2). [On a new plexus discovered in the digestive tube of *Perca fluviatilis*.] C.R. Acad. Sci. URSS 110 1956: 1100-1102 2 figs. [In Russian.]

Mircea, M. see Gyurko, S.

Mironova, N. V. (1). [Nourishment and size of young cod-fish in the littoral zone of the eastern Murman.] Moscow, 1956, pp. 100.

(Mironova, N. V.) (2). [Remarks on the biology and fishery of the coal-fish (*Pollachius virens* L.) in the Barents Sea.] Ichth. Probl. Moscow 8: 155-159 4 figs. [In Russian.]

Mišik, V. Biometrik des Frauenfisches aus der Donau Rutilus pigus (Lacépède 1804) virgo (Heckel 1852). Biológia, Bratislava 12:177–190 1 fig. [German summary.]

Mitani, F. [Correlation between catches of fishing grounds and secular variation in the catch of the yellowtail, Seriola quinqueradiata, along the Pacific coast of Japan.] Bull. Jap. Soc. Sci. Fish. 22:605-608. [English summary.]

Moatu, J. P. see Grangaud, R.

Maen, T. Food of the freshwater drum, Aplodinotus grunniens Rafinesque, in four Dickinson County, Iowa, lakes. Proc. Iowa Acad. Sci. 62 1955: 589-598.

Mohr, E. Vliegende Vissen, Amsterdam (Uitgeverij Breughel) [1955] pp. 63 50 figs.

Mohr, J. L., Willimovsky, N. J. & Dawson, E. Y. An arctic Alaskan kelp bed. Arctic 10:45-52 2 text-figs.

Mohsen, T. Etude morphologique des éléments du systéme suspensorial dans une souche particulière de *Lebistes* reticulatus R. Proc. Egypt. Acad. Sci. 11 1955 (1956): 67-73 3 figs.

(Moiseev, P. A.) (1). [Peculiarities of population dynamics of the commercial fauna in the north-western part of the Pacific, and the causes of these peculiarities.] Zool. Zh., Moscow 35 1956: 1601-1607. [Russian with English summary.]

(Moiseev, P. A.) (2). [Oncorhynchus masu morpha formosanus (Jordan et Oshima) as a non-migratory formin the basin of the river Sedanka.] C.R. Acad. Sci. URSS 112: 163-164. [In Russian.]

Moldenhauer, D. Ein Beitrag zur Systematik der Gattung Acanthophthalmus. Aquar. Terrar. Z. 10:119-1216 figs.

Monod, T. see Le Danois, E.

Montalban, H. R. see Blanco, G. J.

Montalban, H. R. see Rabanal, H. R.

Monteiro, R. Contribuição para o estudo da biologia do Clupeidae de Angola. 1—Sardinella aurita C. & V. An. Jta Inv. Ultramar, Lisbon 9 No. 2 1956: 151-177 6 figs.

Moody, H. L. A fisheries study of Lake Panasoffku, Florida. Quart. J. Fla. Acad. Sci. 20: 21-88.

Moorman, R. B. see Carlander, K. D.

Morawa, F. W. F. Die Verteilung des Fettes bei einigen Plattfischen (Flunder, Kliesche und Heilbutt). Z. Fisch. 5: 269-273 1 fig.

Mori, H. Regeneration des Schwanzes und der Schwanzflosse bei der Knochenfische (Oryzias latipes). Zool. Mag., Tokyo 54 1942:119-124 1 pl. 8 figs.

Mori, K. see Okada, Y.

Mori, S. see Yanagashima, S.

Mori, T. A new species and a new subspecies of Cyprinidae from North China. Zool. Mag., Tokyo 53 1941: 182-184 2 figs.

Mori, T. see Saiki, M.

Morovic, D. Jadranski Mugilidi (Cipli, Skocci) sa bibliografijom Mugilida. Les muges de l'Adriatique avec la bibliographie des muges. Zagreb (Hrvatska Seijacka Tiskara): 1–22 26 figs.

(Morozova, T. A.) [Biology and systematics of the Pacific lamprey.] Ichth. Probl. Moscow 7 1956: 149–157 5 figs. [In Russian.]

(Morozova, T. E.) [The role of external and internal factors in the sex cycle of *Perca fluviatilis* L. C.R. Acad. Sci. U.R.S.S. 112:1129-1132. [In Russian.]

Morris, R. Some aspects of the structure and cytology of the gills of *Lampetra fluviatilis*. Quart. J. micr. Sci. 98:473-485 6 figs.

Morris, R. W. Some aspects of the problem of rearing marine fishes. Bull. Inst. océanogr. Monaco No. 1082 1956: 1-61 9 figs.

Morrow, J. E. (1). Fish records from Long Island Sound. Copeia, 1957: 240-241.

Morrow, J. E. (2). A redefinition of the subspecies of *Fodiator acutus*. Postilla No. 29: 1–10 1 fig.

(Moskalenko, B. K.) [On the influence of several years' fluctuations of the water level in the Ob river on the fecundity and reproduction of certain feedbases.] Zool. Zh., Moscow 35 1956: 746-752. [Russian with English summary.]

Motais, R. (1). Modifications de l'halotropisme et de l'euryhalinité de la jeune anguille sous l'action d'une hormone corticosurrénale. Bull. Inst. océanogr. Monaco No. 1078 1956: 1–14 9 figs.

Motais, R. (2). Sur l'absorption de la lumière par le cristallin de quelques poissons de grande profondeur. Bull. Inst. océanogr. Monaco No. 1074: 1956 1-4.

Motais, R. see Fontaine, M.

Mott, J. C. see Brown, M. E.

Moulton, J. M. A collection of drawings of fishes ascribed to J. P. Kirtland (1793–1877) in the library of Bowdoin College. Breviora No. 80: 1–43 figs.

Moustafa, Y. S. Siluroid fish remains from near Wadi Hoff. Egypt. Bull. Inst. Desert Egypt 3 1953:141-142 (Movchan, V. A. & Yarmoshenko, N. P.) [The eminent Russian pisciculturist O. A. Grimm and his role in the development of Russian fish breeding and fish farming.] Zool. Zh., Moscow 35 1956: 1370-1378 port. [Russian with English summary.]

Muller, T. see Acolat, L.

Munro, I. S. R. (1). Handbook of Australian fishes. Nos. 1–7; 14. Fish. Newsl. 10 No. 7:13–20 figs.; 15 Nos. 8–12 1956:15–18 figs.; 16 No. 1:15–18 figs.; No. 8:15–18 figs.

Munro, I. S. R. (2). Occurrence of the big-eyed tuna, *Parathunnus mebachi* (Kishinouye) in Queensland. Ichthyol. Notes Qd. 1:145-148 1 fig.

Munteanu, I. see Teodorescu-Leonte, R.

Munthe, H. On the development of the Baltic herring in the light of the Late-Quaternary history of the Baltic. Ark. Zool. Stockholm, (4) 9 4 1956: 333-341.

(Murina, V. V.) [The food of eel in Kournian and Vistula bays of the Baltie.] Trud. Ghidrobiol. Obshch. 7 1956: 148-162 6 fig. [In Russian.]

Murray, R. W. Evidence for a mechanoreceptive function of the ampullae of Lorenzini. Nature, London 179: 106-7 figs. 1 & 2.

Musacchia, X. J. & Clark, M. R. Effects of elevated temperatures on tissue chemistry of the arctic sculpin, Myoxocephalus quadricornis. Physiol. Zool. 30:12-17.

Musacchia, X. J. & others. A comparison of liver and muscle lipids in Arctic fishes. Copeia 1957: 10–12.

Mužinic, R. Sur la croissance de la jeune sardine (Sardina pilchardus Walb.) dans les eaux de Split. Notes Inst. oceanogr. Split. No. 12:1-4.

Myers, G. S. Annual fishes. Aquarium J. 23 1952: 125-141 figs.

Myers, G. S. & Storey, M. H. Curatorial practices in zoological research collections. 2. System followed in filing specimens of recent fishes in the Natural History Museum of the Stanford University. Circ. nat. Hist. Mus. Stanf. Univ. No. 6 1956: 1-44.

Nalbandoğlu, U. Über einen Bastard zwischen Rutilus rutilus and Chalcalburnus chalcoides. Hidrobiologi, Istanbul 4: 70-75 3 figs.

Nalbant, T. (1). Contributions à l'étude de la faune ichtyologique de la Dimbovita inférieure. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 345-359 figs. pls. [French summary.]

Nalbant, T. (2). Cobitis aurata vallachica, eine neue Unterart des Balkan-steinpeitzgers (Pisces, Cobitidae). Senckenbergiana biol. 38: 209–212 3 figs.

Nalbant, T. see Banarescu, P.

Navarro, O. B. see Sulet, J. I.

Nedelea, M. see Busnita, T.

(Neïfakh, A. A.) (1). [The change in X-ray sensitivity during fertilization in Misgurnus fossilis.] C.R. Acad. Sci. URSS 109 1956: 943-946 3 graphs. [In Russian.]

(Neïfakh, A. A.) (2). [The influence of ionizing radiation upon the egg cells of *Misgurnus fossilis*.] C.R. Acad. Sci. URSS 111 1956:585-588 2 figs. [In Russian.]

Nelson, E. M. An early review article on the swim bladder of fishes. Copeia 1957: 301–302.

Newell, B. S. see Williams, F.

Newth, D. R. On the neural crest of the lamprey embryo. J. Embryol. exp. Morph. 4 1956: 358-375 2 pls. 12 text-figs.

Nicol, J. A. C. Observations on photophores and luminescence in the teleost *Porichthys*, Quart. J. micr. Sci. 98: 179–188 5 figs.

Nicolae, I. & Cautis, I. Observations sur le comportement du chinchar (Trachurus trachurus mediterraneus Stdr.) au littoral Romain de la Mer Noire —en 1953–1955. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 207–226.

Nicolau, A. see Vasiliu, G. D.

Nielsen, E. Notes on Triassic Fishes from Madagascar. Medd. Dansk Geol. Foren., 12 1955: 563-578 11 text-figs.

Nielson, R. S. & others. A six-year study of the survival and vitality of hatchery reared rainbow trout of catchable size in Convict Creek, California, Calif. Fish Game 43 No. 1: 5-42.

Nieuwenhuizen, A. van den. Remarkable behaviour of *Aphyosemion bivittatum*. Aquarium Philad. **26**:312-3158 figs.

Nigrelli, R. F., Jakowska, S. & Baker, H. A bacterial disease of cichlids in captivity. Aquarium Philad. 26:233-235 3 figs.

(Nikolski, G. V.) (1). [The biological significance of fish schooling.] Trud. Soresh. Ikhtiol. Komm 5 1953 [1955]: 104–116. [In Russian.]

(Nikolski, G. V.) (2). [Some facts on the marine period of life of the Pacific lamprey (*Lampetra japonica*). Zool. Zh., Moscow 35 1956:588-591 Pl. [In Russian.]

(Nikolski, G. V.) (3). [Ichthyological investigations in Chinese Republic.] Zool. Zh., Moscow 35 1956: 1747-1765 8 figs. [In Russian.]

(Nikonorov, I. V.) [Observations on behaviour of Caspian sprat.). Ichth. Probl., Moscow 6 1956: 21-36 10 figs. [In Russian.]

Nilsson, N.-A. On the feeding habits of trout in a stream of Northern Sweden. Rep. Inst. Freshwater Res. Drottningholm, No. 38:154-166 6 figs.

Nishikawa, S. Histological observation on the spermatogenetic cycle of a conger eel, *Muraenesox cinereus* (Forskal). J. Shimonoseki Coll. Fish. 6:281-284, 8 figs. [English summary.]

Noble, E. R. Seasonal variations in host parasite relations between fish and their Protozoa. J. mar. biol. Ass. U.K. 36: 143-1551 fig.

Nogusa, S. [A study of chromosomes in two species of hagfishes (Eptatredidae-Cyclostomata)]. Zool. Mag. Tokyo 66: 307–309. 4 figs. [English summary.]

Norris, K. S. Second record of the green sturgeon in southern California. Calif. Fish Game 43:317.

(Novikova, N. S.) [Behaviour of roach on its feeding grounds in the northern Caspian Sea.] Ichth. Probl., Moscow 7 1956: 36-51 3 figs. [In Russian.]

Nybelin, O. Deep-sea bottom fishes. Rep. Swed. Deep-Sea Exped. 1947–48 2 Zool. 3: 247–345 7 pls., 50 figs. [Obruchev, D. V.] (1). In Atlas of the guide forms of the fossil faunas of the USSR 3 1947: 1-245, 56 pls. figs. Leningrad.

Obruchev, D. V. (2). [Review of L. S. Berg's "Classification of Agnatha and Fishes, recent and fossil," 2nd Edit.] Zool. Zh. Acad. Sci. U.R.S.S. 36: 300-305. [In Russian.]

Ochiai, A. Zoogeographical studies on the soleoid fishes found in Japan and its neighbouring regions. I, II and III. Bull. Jap. Soc. sci. Fish. 22: 522-535 1 fig. [English summary.]

Odiorne, J. M. see Brown, M. E.

Oguri, M., & Hibiya, T. (1). [Studies on the adrenal glands of teleosts—I. Some observations from the viewpoint of comparative histology.] Bull. Jap. Soc. sci. Fish. 22: 621–625 2 figs. 1 pl. [English summary.]

Oguri, M. and Hibiya, T. (2). Studies on the adrenal gland of Teleosts. II. On the adrenal tissues in 15 species of fishes. Bull. Jap. Soc. sci. Fish. 23: 144-149 3 figs.

Ohno, S. see Saiki, M.

Okada, Y. [Introduction of Ophiocephalus to Japan.] Jap. J. Ichthyol. 5:174. [In Japanese.]

Okada, Y. & Kubota, S. S. (1). Studies on the fish fauna at the southern vicinity of Tsu City, Mie Prefecture. Rep. Fac. Fish. Univ. Mie 2 1956: 205-216 2 pls. 1 fig.

Okada, Y., & Kubota, S. S. (2). Description of a new fresh-water cat-fish, Coreobagrus ichikawai, with an emendation of the genus Coreobagrus Mori. Jap. J. Ichthyol. 5:143-1451 fig.

Okada, Y. & Mori, K. [Ecological study of Salangichthys microdon in breeding season.] Jap. J. Ichthyol. 5:99-106 2 figs. [English summary.]

Okada, Y & Suzuki, K. Ecological studies of the Ma-anago, Astroconger myriaster (Brevoort). I. Notes on the composition of total length, the relationship between total length and body weight and the condition factor in Ise Bay, Mie Prefecture. Rep. Fac. Fish. Univ. Mie 2:217-226 4 figs.

(Olifan, V. I.) [On the 24 hour rhythm of Baikal grayling fry and on the 24 hour rhythm in fish fry in general]. C.R. Acad. Sci. URSS 114:669-672 fig. [In Russian.]

Oliva, O. (1). Příspévek k posnání rybi fauny Bečvy. Přírod. Sborn. Ostravsk. Kraje 13 1952:193-203.

Oliva, O. (2). Some experiments with the use of fish poisons during fish population studies. Acta Soc. zool. Bohemoslovenicae 21:363-375.

Oliva, O. & Frank, S. Species composition, abundance and growth of fish population in the reinforced concrete basin in Prague. Univ. Carolina Biologica (Prague) 3: 1-6.

Olivereau, M. (1). Hypophyse et gland thyroide chez les poissons. Année biol. (3) 30 1954: 63-80.

Olivereau, M. (2). Radiothyroï-dectomie chez l'anguille (Anguilla anguilla L.). Arch. Anat. micr. Morph. exp. 46: 39-59 9 figs.

Olson, P. R. see Donaldson, L. R.

Olsson, R. An experimental breakage of Reissner's fibre in the central canal of the pike (Esox lucius). Z. Zellforsch. 46:12-17.

Olsson, R. see Afzelius, B. A.

Oosten, J. van see Brown, M. E.

Opalatenko, L. K. see Balabai, P. P.

Oren, O. H. Changes in temperature of the Eastern Mediterranean Sea in relation to the catch of the Israel. Trawl Fishery during the years 1954-55 and 1955-56. Bull. Inst. océanogr. Monaco No. 1102:1-15 tables.

Orska, J. The influence of temperature on the development of the skeleton in teleosts. Zool. Polon. 7:272-325 3 pls.

Orton, G. L. Embryology and evolution of the pelagic fish egg. Copeia **1957** : 56–58.

ørvig, T. (1). Remarks on the vertebrate fauna of the Lower Upper Devonian of Escuminac Bay, P.Q., Canada, with special reference to the Porolepiform Crossopterygians. Zool. 10: 367-426 13 figs.

Ørvig, T. (2). Paleohistological notes. 1. On the structure of the bone tissue in the scales of certain Palaeonisciformes. Ark. Zool. 10:481-490 4 figs.

Ørvig, T. (3). Notes on some Paleozoic lower vertebrates from Spitsbergen and North America. geol. Tidsskr. 37: 285-353, 5 pls. figs.

Oshima, M. (1). Notes on the bottomfishes obtained at off-shores of Niigata and Yamagata Prefectures, with descriptions of five new species. Jap. J. Ichthyol. 6: 1-85 figs.

Oshima, M. (2). List of the freshwater fishes obtained in the southeastern Kiushiu. Jap. J. Ichthyol. 6:

Oshima, M. (3). Notes on the fishes collected in the down stream of Tama River, with descriptions of two new species. Jap. J. Ichthyol. 6:14-19 2 figs.

Oshima, M. (4). Studies on the distribution of freshwater salmon Oncorhynchus masou (Brevoort) and Oncorhynchus rhodurus Jordan & McGregor, found in the streams of Kiushiu. Zool. Mag. Tokyo 66:21-24 2 figs. [English summary.]

Otsu, T. & Yoshida, H. O. Another record of the rare shark, Euprotomicrus bispinatus, from the Pacific Ocean. Copeia 1957: 58.

Ovsyannikov, N. S. see Borisov, P. G.

Owen, D. M. see Johnson, H. R.

Padoa, E. & others. Uova, larve e stadi giovanili di Teleostei. Fauna e Flora Neapel 38 Pt. 3 1956: 457–1064 16 pls.

Papp, A. & Thenius, E. Vösendorf ein Lebensbild aus dem Pannon des Wiener Beckens. Mitt. Geol. Gesell. Wien 46 Sonderband 1954:1-109 14 pls.

Parker, R.-R. see Larkin, P. A.

Parrott, A. W. Sea anglers' fishes of New Zealand. London (Hodder & Stoughton) pp. 176 8 pls. text-figs.

Paulian, P. La pêche autour des îles St. Paul et Amsterdam. Terre et la Vie 104: 267-282 3 figs.

(Pavlov, P. I.) [Comparison between Danube and Dnieper bream]. Zool. Zh., Moscow **35** 1956 : 891–896 fig. [Russian with English summary.]

Pera, L. Contributo alla morfologia ed allo sviluppo della ghiandola digitiforme dei plagiostomi. Marit. Zool. Ital., Suppl. **65**: 385–386.

Perruche, L. La pêche électrique en eau douce. Nature, Paris No. 3263 : 88-90 4 figs.

Peters, A. J. Bibliography of published work bearing on the natural history of the Seychelles and neighbouring archipelagoes. J. Soc. bibl. nat. Hist. 3:238-262.

Peters, H. M. Über die Regulation der Gelegegrösse bei Fischen. Z. Naturf. 12B: 255-261 1 pl., 5 figs.

(Petrova, V. G.) [A cytochemical study of egg-cells in the fish]. C.R. Acad. Sci. URSS 110 1956:674-676 2 fig. [In Russian.]

Pew, P. Occurrence of young dolphin, Coryphaena hippurus in a Texas bay. Copeia **1957**: 300.

Peyer, B. Über die morphologische Deutung der Flossenstachelm einiger Haifische. Mitt. Naturf. Ges. Bern N.F. 14:159-176 10 pls. 7 text figs.

Phillips, C. A South Florida record for the stingray, *Dasyatis centroura*. Copeia 1956 No. 4 [1957]: 249.

Pickford, G. E. (1). The response of hypophysectomized male killifish to purified fish growth hormone, as compared with the response to purified beef growth hormone. Endocrinology 55 1954: 274–287.

Pickford, G. E. (2). The response of hypophysectomized male killifish to prolonged treatment with small doses of thyrotropin. Endocrinology 55 1954: 589–592.

Pickford, G. E. & Kosto, B. Hormonal induction of melanogenesis in hypophysectomized killifish (Fundulus heteroclitus). Endocrinology 61:177-196.

Pickford, G. E. see Atz, E. H.

Pickford, G. E. see Bondy, P. K.

Picos, C. A. Recherches sur l'action dynamique spécifique des protéines chez la carpe. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 373–384.

Picos, C. A. & Todicescu, S. L'Influence de la temperature sur le metabolisme energique global de la carpe. Ann. Univ. "C. J. Parhon" Bucuresți, Ser. Sti. Nat. No. 9 1956: 153–158. [French summary.]

(Pidoplichko I. G. & Kurazhskovski, Y. N.) [Palaeogeographic importance of Dipnoi. Trav. Mus. zool. Acad. Sci. Ukr. 27 1956:144-153. [In Ukrainian with Russian summary.]

Pihlgren, A. M. see Suomalainen, P.

Pillsbury, R. W. Avoidance of poisonous eggs of the marine fish. Scorpaenichthys marmoratus by predators. Copeia, 1957: 251–252.

Pinc, R. D. see Prosser, C. L.

Pinter, H. Grossflosser. Aquar. Terrar. Z. 10:169-172, 1 colour-plate and text-figs. 1-5.

Pinto, S. Y. (1). Um novo Clinidae da Ilha da Trindade, Brasil. Bol. Mus. nac. Rio de J., (Zool.) No. 163: 1-15 12 figs.

Pinto, S. Y. (2). Sobre a validade do gênero Parablennius Ribeiro, 1915 (Blenniidae, Perciformes, Actinopterigii). Rev. Bras. biol. 17: 413-420 3 figs. Pinto, S. Y. see Travassos, H.

Pitt, T. K. & others. Temperature selection of the carp (Cyprinus carpio Linn.). Canad. J. Zool. 34 1956: 555-557.

Pliszka, F. Importance of the aquatic animals as a food base of fish after Polish investigation. Polsk. Arch. Hydrobiol. 3 1956: 429-458.

Poczopko, P. (1). An outline of fishery conditions in the Vistula river near Torun in 1952–1954. Rocz. Nauk. Rolnic. 70B 1955: 107–120 1 fig. [English summary.]

Poczopko, P. (2). The population of bream (Abramis brama L.) in the Vistula in the neighbourhood of Torun. Zeoz. Nauk. Univ. M. Kopernika. Biol. No. 1 1956: 165–185 5 figs. [English summary.]

Poersch, L. Der kleine (gelbe) Kongosalmler (*Petersius caudalis*). Aquar. Terrar. Z. 10:172-175.

(Polenov, A. L.) [Content and distribution of nucleic acids in neurosecretory cells of the diencephalon in Teleostei]. C.R. Acad. Sci. U.R.S.S. 112:1122-1125 Pl. [In Russian.]

Politis, J. (1). Recherches cytologiques sur la formation du pigment mélanique chez certains poissons. C.R. Acad. Sci. Paris 243 1956: 1559–1560.

Politis, J. (2). Recherches cytologiques sur la formation des carotinoïdes chez certains poissons. C.R. Acad. Sci. Paris 244: 511-513 2 figs.

Poll, M. (1). L'état actuel de nos connaissances sur la faune ichthyologique du Congo Belge. C.R. IIIe Congr. nat. Sci. Brux. 8 [1950]: 43-46.

Poll, M. (2). Ecologie des poissons du Lac Tanganika. Int. Congr. Zool. 14 1953 (1956): 465–468.

Poll, M. (3). Redescription du Gymnallabes tihoni Poll 1944, Clariidae microphthalme du Stanley-Pool (Congo Belge). Rev. Zool. Bot. afr. 55: 237–248 10 figs.

Poll, M. see Brien, P.

Pollingher, U. see Santa, V.

Popescu, E. see Popescu-Gorj, A.

Popescu, L. see Teodorescu-Leonte, R.

Popescu-Gorj, A. & others. Recherches hydrobiologiques-piscicoles dans le complexe des lacs Borcea-de-Jos. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 123-166.

Popescu-Gorj, A. see Busnita, T.

Pora, E. A. & others. La nourriture du chinchard (*Trachurus trachurus* mediterraneus Stdr.) durant une saison de pêche (Mai.-Octobre 1953). Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 187-206.

Porumb, I. & Fl. see Pora, E. A.

Postel, E. (1). Note succincte sur quelques Espadons tunisiens. Bull. Mus. Hist. nat. Paris (2) 28:509-511.

Postel, E. (2). Une nouvelle station a Hemichromis bimaculatus en Afrique du Nord: La Mare de Kébili en Tunisie. Bull. Mus. Hist. nat. Paris (2) 29:143-144 map.

(Povlovski, E. N.) [Structure of fish brain in connection with their mode of life.] Trud. Sovesh. Ikhtiol. Komm. 5 1953 [1955] 76-77. [In Russian.]

(Privolnev, T. I.) [Reaction of fishes to light]. Iehth. Probl., Moscow 6: 1956: 3-20 5 figs. [In Russian.]

Prokhorov, V. G. see Mednikov, B. M.

Prosser, C. L. & others. Acclimation of goldfish to low concentrations of oxygen. Physiol. Zool. 30: 137-141.

Pynnönen, A. Über die Reichlichkeit der Fische und Reptilien im Kirchspiel Konnoveu einst und jetzt. Arch. Soc. zool.-bot. Vanamo 11:171-182.

Qasim, S. Z. The biology of *Blennius* pholis L. (Teleostei). Proc. zool. Soc. Lond. 128: 161–208 13 figs 1 pl.

Quaghebeur, M. Onderzoek over de reductie van de ogen, van de oogkas en van het tectum opticum bij blinde grotvissen. Catholic Univ. Louvain: 1–120 27 figs. [In Flemish.]

Quartier, A. Sur la présence de Leucaspius delineatus Heckel dans le lac de Neuchâtel. Bull. Soc. neuchâtel. Sci. nat. (3) 78 1955: 31–32.

Querner, H. Über einen Zusammenhang zwischen der Beeinflussbarkeit der Gonaden durch Sexualhormone und dem Modus der Geschlechtsbestimmung. Verh. deuts. zool. Ges. 1956: 194–197.

Querrière, F. de la see Fontaine, M.

Rabanal, H. R., Montalban, H. R. & Villaluz, D. K. The preparation and management of Bañgos fishpond nursery in the Philippines. Philipp. J. Fish. 1 1951: 1-34 10 pls. 8 figs.

Rabeu, V. von. Vom Leben und Wandern des Aales. Kosmos, Stuttgart 48 1952 : 506-510 3 figs.

Radulescu, I. & Vasiliu-Suceveanu, N. Contributions à la connaissance des parasites des poissons du complexe lagunaire Razelm-Sinoe. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 309–333.

Rae, B. B. & Wilson, E. Rare and exotic fishes recorded in Scotland during 1954 and 1955. Scot. Nat. 68 1956: 23–38; 92–109.

Rae, B. B. see Stephen, A. C.

Raffy, A. see Fontaine, M.

Rageau, J. see Gail, R.

Rahn, J. Beitrag zur Altersbestimmung an Aalen. Z. Fisch. 6:393–396 2 figs.

Ramaswami, L. S. & Sundararaj, B. I. (1). Some aspects of induced spawning in the catfish *Heteropneustes*. Naturwissenschaften 44:46.

Ramaswami, L. S. & Sundararaj, B. I. (2). Induced spawning in the catfish *Clarias*. Naturwissenschaften 44: 384. Raney, E. C. see Robins, C. R.

Ranzi, S. see Citterio, P.

Ranzi, S. see Maldacea, L.

Rasalan, S. B. & Villadolid, D. V. The basnig, a bag net for pelagic fishing in the Philippines. Philipp. J. Fish. 3 1955: 1-29 1 pl. 9 figs.

Raut. H. Acanthophthalmus cuneovirgatus spec. nov. und Acanthophthalmus robiginosus spec. nov., mit einer übersicht über die in Deutschland eingeführten Acanthophthalmus-Arten und einem Bericht über ihre Haltung im Aquarium. Aquar. Terrar. Z. 10: 29-33 8 figs.

Raynaud, P. see Serfaty, A.

Rechnitzer, A. B. The use of paper partition chromatography in taxonomic studies. Int. Congr. Zool. 14 1953 (1956): 333–335.

Reed, R. J. (1). The prolonged spawning of the rosyface shiner, *Notropis rubellus* (Agassiz) in northwestern Pennsylvania. Copeia 1957: 250.

Reed, R. J. (2). Phases of the life history of the rosyface shiner, *Notropis rubellus*, in Northwestern Pennsylvania. Copeia **1957**: 286–290 1 fig.

Reichenbach-Klinke, H. H. Augenschäden bei Meeresfischen durch den Pilz Ichthyosporidium hoferi (Plehn et Mulsow) und Bemerkungen zu seiner Verbreitung bei Mittelmeerfischen. Pubbl. Staz. zool. Napoli 29: 22–32 1 fig. 2 pls. [English summary.]

Reid, G. K. (1). External morphology of an embryo whale shark, *Rhincodon typus* Smith. Copeia 1957:157-158 1 fig.

Reid, G. K. (2). Biologic and hydrographic adjustment in a disturbed Gulf Coast estuary. Limnol. Oceanogr. 2:198-2124 figs.

Reimers, N. see Nielson, R. S.

Reinboth, R. Sur la sexualité du Téléostéen Coris julis (L.). C.R. Acad. Sci. Paris 245 : 1662–1665.

Reynolds, R. A. see Hubbs, C.

Ribeiro, P. de M. (1). On a new genus and a new species of South American fishes. Int. Congr. Zool. 14 1953 (1956): 546–547 1 fig.

Ribeiro, P. de M. (2). Notas para o estudo dos Pygidiidae Brasileiros (Pisces Pygidiidae) VI. Pap. Rep. Zool. Sec. Agric. S. Paulo 13:71-73.

Richard, L. see Devillers, C.

Riggs, C. D. Mugil cephalus in Oklahoma and Northern Texas. Copeia 1957: 158-159.

Robertson, G. M. (1). Paleoecology of the Agnatha (cf. Bibliography). Nat. Res. Council Rep. Comm. on Treatise on Marine Ecology and Paleoecology 1948–1949. Div. Geol. & Geogr. Dec. 1949 pp. 111–116.

Robertson, G. M. (2). Agnatha in Treatise on marine ecology and paleoecology, vol. 2. Mem. geol. Soc. Amer. 67:995–998.

Robertson, J. D. The habitat of the early vertebrates. Biol. Reviews 32: 156-187.

Robertson, O. H. Survival of precociously mature king salmon male parr (Oncorhynchus tshawytscha Juv.) after spawning. Calif. Fish Game 43: 119–129 4 figs.

Robins, C. R. (1). Effects of storms on the shallow-water fish fauna of southern Florida with new records of fishes from Florida. Bull. mar. Sei. Gulf & Caribbean 7: 266–275 1 fig.

Robins, C. R. (2). A study of the bluefin tuna populations from the eastern and western North Atlantic based on an analysis of fin-ray and gillraker counts. Charles F. Johnson Oceanic Gamefish Invest., Univ. Miami, Fla. 57-13 (1957) (mimeographed). pp. 1-7.

Robins, C. R. & Miller, R. R. Classification, variation, and distribution of the sculpins, genus *Cottus*, inhabiting Pacific Slope waters in California and Southern Oregon, with a key to the species. Calif. Fish Game 43: 213–233.

Robins, C. R. & Raney, E. C. (1). Distributional and nomenclatorial notes on the suckers of the genus *Moxostoma*. Copeia 1957: 154-155.

Robins, C. R. & Raney, E. C. (2). The systematic status of the suckers of the genus *Moxostoma* from Texas, New Mexico and Mexico. Tulane Stud. Zool. 5: 291-318 12 figs.

Robins, C. R. see Voss, G. L.

Roche, G. (1). Ray's Bream, Brama raia Bloch, from the mouth of the river Moy. Irish Nats'. J. 12 1956: 109.

Roche, G. (2). Sea horse, Hippocampus hippocampus (L.), from Co. Clare, Irish Nats'. J. 12 1956: 110.

Rogers, K. T. Optokinetic testing of cyclopean and synophthalmic fish hatchlings. Biol. Bull. Woods Hole 112: 241–248 8 figs.

Roi, P. E. see Ivasik, V. M.

Roland, J. see Dieuzeide, R.

Rosca, I. D. see Pora, E. A.

Rose, E. T. Notes on the age and growth of Spirit Lake yellow pikeperch (Stizostedion v. vitreum). Proc. Iowa Acad. Sci. 58 1951: 517-525.

Rosen, W. G. see Brown, D. G.

Ross, R. J. Ordovician Fossils from wells in the Williston Basin, Eastern Montana. Bull. U.S. geol. Surv. 1021-M: 439-510 8 pls.

Roussow, G. Some considerations concerning sturgeon spawning periodicity. J. Fish. Res. Bd. Can. 14: 553-572 11 figs. 1 pl.

Roy, H. Fischpankreas als Rohstoffquelle für die Insulingewinnung unter besonderer Berücksichtigung des Dorsches (Gadus morrhua). Z. Fisch. 6:469–487 10 figs.

Rübenberg, L. see Meyer-Waarden, P. F.

Runnström, S. Migration, age and growth of the brown trout (Salmo trutta L.) in Lake Rensjön. Rep. Inst. Freshwater Res. Drottningholm No. 38: 194–246. 17 figs.

Rushton, W. (1). Fish diseases due to bacteria. Salm. Trout Mag. 1956: 289–297 1 pl. 2 figs.

Rushton, W. (2). Fish diseases due to bacteria (2). Salm. Trout Mag. No. 150: 165-171 2 figs.

Rutte, E. Einführung in die geologie von Unterfranken. Würzburg: 1–168 8 pls. figs.

Rychlicki, Z. & Zarnecki, St. Die Zatorer Karpfenaufzuchtmethode und deren Einfluss auf die Beseitigung der Bauchwassersucht. Z. Fisch. 5:423-442.

Sadoğlu, P. A Mendelian gene for albinism in natural cave fish. Experientia 13:394.

(Sadov, I. A.) (1). [Development of perivitelline space in the oocytes of sturgeons.] C.R. Acad. Sci. URSS 112:174-176 3 figs. [In Russian.]

(Sadov, I. A.) (2). [Changes occurring in the size of nuclei during growth and maturation of the oocytes in sturgeon.] C.R. Acad. Sci. URSS 112:359-362 2 figs. [In Russian.]

Sahrhage, D. Heringswanderungen. Ergebnisse der Markierung von Heringen. Kosmos, Stuttgart 53:44–49 10 figs.

Saiki, M., et al. [Studies on the radioelements in the contaminating radioactive fish. II. On skipjacks caught in the Pacific Ocean in 1956.] Bull. Jap. Soc. Sci. Fish. 22:645-653 10 figs. [English summary.]

Saint-Seine, P. de. L'Evolution de Actinoptérygiens. Colloques Internal., Centre nat. Recherches Sci., Paris 60: 1956: 27–34.

Salzen, E. A. A trawling survey off the Gold Coast. J. Cons. Int. explor. Mer 23: 72-82 2 figs.

Sandon, H. An abnormal specimen of Synodontis membranaceus (Teleostei, Siluroidea), with a discussion on the evolutionary history of the adipose fin in fish. Proc. zool. Soc. Lond. 127 1956: 453-459 Pl. 1.

San Juan, R. C. see Sulet, J. I.

Santa, V. & Pollingher, U. L'anatomie macro- et microscopique des glandes annexes du tube digestif chez la carpe. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 335-344 figs.

Santos, R. S. Ocorrência de folhelho fossiléfero Cretácico no Município de Presidente Olegário, Minas Gerais. Bol. Div. Geol. min. Rio de J. 155 1955: 1–273 pls.

Santos Pinto, J. dos (1). Parasitic castration in males of Sardina pilchardus (Walb.) due to testicular infestation by the Coccidia Eimeria sardinae (Thélohan). Rev. Fac. Cienc. Lisbon (2C) 5: 209-224 3 pls. 1 fig.

Santos Pinto, J. dos (2). Deux cas de tumeurs sexuelles dans la sardine (Sardina pilchardus Walb.). Rec. Fac. Cienc. Lisbon (2C) 5:257-260 2 pls.

Sanzo, L. see Padoa, E.

Šapkarev, J. (1). La teneur en eau de la truite *Salmo letnica* Kar. au cours du développement. Rec. Trav. Sta. Hydrobiol. Ohrid 2 1954: 1–27. [French summary.]

Šapkarev, J. (2). Variations saisonnières de la teneur en graisse chez la truite d'Ohrid (*Salmo letnica*, Karaman). Rec. Trav. Sta. Hydrobiol. Ohrid **3** 1955: 1–21 3 figs.

Sarkar, H. L. Composition of *Hilsa* catch in the Sundarbans of west Bengal during the winter months. Indian J. Fish. 4:340-343 1 fig.

Sarkar, H. L. & Kapoor, B. G. Deformities in some Indian cat fishes. J. Zool. Soc. India 8 1956:157-164 5 figs.

Sarma, K. S. On the palaeontology of the Limestone Band of the Baripada Beds, Mayurbhanj, Orissa. Quart. J. Geol. Soc. India 29: 103-105 1 pl.

Sato, M. On the pit organ system of Pseudobagrus aurantiacus, Astroconger myriaster and Gobies. Zool. Mag., Tokyo 66: 332–336 4 figs.

Sauerteig, E. & Krebs, W. Eine neue methode zur Sektion kleiner Fische. Aquar. Terrar. Z. 10:128-130 5 figs.

(Savina, N. O.) [Biology of Schizothorax argentatus in the lake Balkhash.] Izv. Fish. Inst. Fishery resources in the lake Balkhash 37 1956: 129–157.

Sawyer, P. J. Laboratory care and feeding of larval lampreys. Copeia 1957: 244.

Scattergood, L. W. & Coffin, G. W. Records of some Gulf of Maine fishes. Copeia 1957: 155-156.

Schaefer, M. B. Some considerations of population dynamics and economics in relation to the management of the commercial marine fisheries. J. Fish. Res. Bd. Can., 14:669-681.

Scheier, A. see Cairns, J.

Schiemenz, F. Ersatz des instinktmassigen Wanderns der Fische in Fischtreppen durch das reflektorische Wandern als Grundlage zum Bau von Fischtreppen, sowie die Orientierung und das Verhalten der Fische in Fischtreppen. Z. Fisch. 6:61–84 19 figs.

Schindler, O. (1). Neues von den lebenden "Urweltfischen". Kosmos, Stuttgart 51 1955:581-588 11 figs.

Schindler, O. (2). Freshwater fishes ... translated and edited by P. A. Orkin. London & New York, (Thames & Hudson) pp. 243 28 pls. (7 col.) 50 figs.

Schindowski, E. & Tesch, F. W. Methodisches zur Wachstumsrückberechnung. erläutert am Beispiel von Lucioperca sandra Cuv. u. Val., Perca fluviatilis L. und Salmo trutta fario L. Z. Fisch. 5: 247–267.

Scholander, P. F., van Dam, L. & Enns, T. The source of oxygen secreted into the swimbladder of cod. J. Cell. comp. Physiol. 48 1956: 517-522 1 fig.

Scholander, S. Cold fish problem. Nat. Hist. N.Y. 66: 456-466 figs.

Schreiter, T. Über eine eigenartige Schwimmblasenverlagerung bei Karpfen. Z. Fisch. 6:301-313 12 figs.

Schroeder, W. C. see Bigelow, H. B.

Schroll, F. Zur Ernährungsphysiologie der Ammocoeteslarven der Cyclostomen. Zool. Anz. 159: 75–78 2 figs.

Schubert, P. see Bauer, K.

Schultz, L. P. (1). A new pinecone fish, *Monocentris reedi*, from Chile, a new family record for the Eastern Pacific. Proc. U.S. nat. Mus. 106 1956: 237-239 1 fig.

Schultz, L. P. (2). The frogfishes of the family Antennariidae. Proc. U.S. Nat. Mus. 107:47-105 14 pls. 8 figs.

Schultz, L. P. (3). The generic names Barbus and Puntius. Trop. Fish, Hobbyist 5: 14-15; 29-31

Schwartz, F. J. First record of infestation and death in the ictalurid catfish, Schilbeodes miurus, by the parasite Clinostomium marginatum. Copeia 1956 No. 4 [1957]: 250-251.

Scott, J. S. Fish identification photographs. Angling in Malaya 4 1956: 60.

Scott, T. D. A new blenny (Triptery-giidae) and pipefish (Syngnathidae) from Kangaroo Island, South Australia. Trans. Roy. Soc. S. Austr. 80: 180–1832 figs.

Scott, W. B. Distributional records of fishes in Western Canada. Copeia 1957: 160–161.

Sedlár, J. Beitrag zur Eiermenge beim Ameiurus nebulosus (Le Suer, 1819) aus dem Flussame des Flusses Žitava. Biológia, Bratislava 12:708-710.

Segaar, J. Brain and instinct with Gasterosteus aculeatus. Proc. Akad. Wet. Amst. 59C 1956: 738-749 11 figs.

(Semenov, K. I.) [On the influence of different light conditions upon the development of sturgeon-larvae within the period from hatching to active feeding under artificial breeding. C.R. Acad. Sci. URSS 113: 937-940 2 figs. [In Russian.]

(Semenov-Tyan-Shanski, D. I.) [On the spawning of lamprey]. Zool. Zh., Moscow 35 1956: 937-938. [In Russian.]

(Semko, R. S.) [New data on Oncorhynchus masu in Western Kamchatka]. Zool. Zh., Moscow 1956: 1017-1022. [Russian with English summary.]

Serfaty, A., & Raynaud, P. Quelques données sur l'électro-cardiogramme de la tanche (*Tinca tinca* L.) en eau douce, Bull. Soc. zool. France 82:49-56.

(Shamardine, I. P.) [The stages in development of pike]. Trud. Inst. Morf. Zhiv. 16: 237-298 27 figs. [In Russian.]

(Shchetinina, L. A.) [On the reproduction of the Prussian carp in Veselovsk water-reservoir]. Zool. Zh., Moscow 35 1956:1517-1520. [Russian with English summary.]

Shelbourne, J. E. The feeding and condition of plaice larvae in good and bad plankton catches. J. Mar. Biol. Ass. U.K. 36: 539-552 1 pl. 3 figs.

Shetter, D. S. see Whalls, M. J.

Shimada, B. M. see Alverson, F. G.

Shirai, K. see Saiki, M.

Shiraishi, Y., & Uchida, I. [On the morphological changes of the salmon fry with the absorption of yolk.] Jap. J. Ichthyol. 5:85-92 3 figs. 4 figs.

Shirkova, A. P. [On the biological heterogeneity of Coregonus lavaretus maraenoides Poljakow in connection with its introduction into other water reservoirs.] Zool. Zh., Moscow 36: 951-953. [English summary.]

Shkorbatov, G. L. Experimental basis of fish acclimatization. Zool. Zh., Moscow 36:230-237. [English summary.]

(Shmalganzen, O. I.) [Ecological and morphological peculiarities in the development of the gill apparatus in Volga sturgeon larvae]. C.R. Acad. Sci. URSS 100 1955: 837–839 2 figs. [In Russian].

(Shturbina, M. A.) [The quality of the sturgeon fry in the rivers, related to the individual peculiarity of the eggs.] St. Petersb. Acad. Imp. Sci. Ichthyol. Com. 4 1955:105-113 4 graphs. [In Russian.]

(Shulman, G. E.) [On cutaneous respiration in Gobioids]. Zool. Zh., Moscow 35 1956: 314-316. [In Russian.]

(Shulman, G. E. & others). [Gas exchange in Neogobius melanostomus and N. syrman related to habitat in Azov Sea.] Ichth. Probl., Moscow 8: 77-80. [In Russian.]

Sibakin, K. see Kleerekoper, H.

Siddiqui, A. A. see Bhuiya, M. A. H.

Siegfried, P. Der stratigraphische Wert der Fischfauna des westfalischen Ober-Campan. Paläont. Zeitschr. 30 1956: 37–38.

Silas, E. G. The Ceylonese cyprinid genus *Eustira* Günther considered a synonym of *Danio* Hamilton. Copeia 1957: 61–62 1 fig.

Simmons, E. G. An ecological survey of the Upper Laguna Madre of Texas. Publ. Inst. Mar. Sci. Univ. Texas. 4 No. 2:156-200.

Singer, B. Living silver. An impression of the British fishing industry. London (Secker & Warburg) pp. [vi] + 232 text illust.

Sivalingam, S. & Medcof, J. C. General features and productivity of the Wadge Bank trawl fishery. Bull. Fish. Res. Sta. Ceylon No. 6:1-23 6 figs.

Slastenenko, E. P. A list of natural fish hybrids of the world. Hidrobiologi, Istanbul 4: 76-97.

Smigielska, T. Stratigraphical importance of the ichtiofauna from the Carpathian menilite Shales. Przeglad Geologiczny 1957: 567–570 figs.

(Smirnova, E. N.) (1). [Embryonic and larval peculiarities in development of *Vimba vimba* natio *carinata* (Pall.)]. Trud. Inst. Morf. Zhiv. 20:71-94 6 figs. [In Russian.]

(Smirnova, E. N.) (2). [Morphological and ecological peculiarities in the development of *Rutilus frisii* (Nordm.)]. Trud. Inst. Morf. Zhiv. 20 · 95–120 7 figs. [In Russian.]

(Smirnova, L. I.) [The physiological role of leucocytes in digestion of fishes]. Ichth. Probl., Moscow 7 1956:107–118. [In Russian.]

Smith, C. L. Two new clinid blennies (*Malacoctenus*) from Puerto Rico. Occ. Pap. Mus. Zool. Univ. Mich. No. 585: 1-15 1 pl.

Smith, I. C. New restorations of the heads of *Pharyngolepis oblongus* Kiaer and *Pharyngolepis kiaeri* sp. nov., with a note on their lateral-line systems. Norsk geol. Tidsskr. 37: 373–402 6 pls. figs.

Smith, I. W. The occurrence of damaged sea trout on the east coast of Scotland. Salm. Trout Mag. No. 150: 148-149 2 figs.

Smith, J. L. B. (1). A remarkable new unicorn fish from East Africa. Ann. Mag. nat. Hist. (12) 9 1956 [1957]: 686-688 1 fig.

Smith, J. L. B. (2). The fishes of Aldabra-Part V. Ann. Mag. nat. Hist. (12) 9 1956 [1957]: 721-729 3 figs.

Smith, J. L. B. (3). The fishes of Aldabra. Part VI. Ann. Mag. nat. Hist. (12) 9 1956 [1957]: 817–829 5 figs.

Smith, J. L. B. (4). The fishes of Aldabra—Part VII. Ann. Mag. nat. Hist. (12) 9:888-892 1 fig.

Smith, J. L. B. (5). Deep-line fishing in Northern Mozambique, with the description of a new pentapodid fish. Ann. Mag. nat. Hist. (12) 10: 121-124 1 pl.

Smith, J. L. B. (6). Dasyatid rays new to South Africa. Ann. Mag. nat. Hist. (12) 10:429-431 2 pls.

Smith, J. L. B. (7). The genus Luzonichthys Herre. Copeia 1956 [1957]: 251.

Smith, J. L. B. (8). Two rapid fatalities from stonefish stabs. Copeia, 1957: 249.

Smith, J. L. B. (9). The fishes of the family Scorpaenidae in the Western Indian Ocean. Pt. 1. Ichthyol. Bull. Rhodes Univ. No. 4:49-69 4 pls. 5 figs.

Smith, J. L. B. (10). The fishes of the family Scorpaenidae in the western Indian Ocean. Pt. II. The subfamilies Pteroinae, Apistinae, Setarchinae and Sebastinae. Ichthyol. Bull., Rhodes Univ. No. 5:75–87 2 pls. 4 figs.

Smith, J. L. B. (11). Sharks of the genus *Isurus* Rafinesque, 1810. Ichthyol. Bull., Rhodes Univ. No. 6: 91-96 l pl. l fig.

Smith, J. L. B. (12). List of the family Labridae in the Western Indian Ocean with new records and five new species. Ichthyol. Bull., Rhodes Univ. No. 7:99-114 2 pls. 4 figs.

Smith, J. L. B. (13). The labrid fishes of the subgenus Julis Cuvier, 1814 (in Coris Lacepede, 1802) from South and East Africa. Ichthyol. Bull., Rhodes Univ. No. 8:117-120 2 pls. 1 fig.

Smith, J. L. B. (14). Four interesting new fishes from South Africa. S. Afr. J. Sci. 53: 219–222 3 figs.

Smith, J. L. B. (15). A new shark from South Africa. S. Afr. J. Sci. 53: 261-264 2 figs.

Smith, J. L. B. (16). A preliminary survey of the scylliogaliid dogfishes of South Africa. S. Afr. J. Sci. 53: 353–359 2 figs.

Smith, S. see Brown, M. E.

Smith, S. H. Evolution and distribution of the coregonids. J. Fish. Res. Bd. Can. 14: 599-604 1 fig.

Smith, W. C. Gonad condition, age and size of Manx herrings, 1945-53, and comparison with earlier years. Rep. Mar. Biol. Sta., Pt. Erin. 69 1956: 21-28 2 figs.

Smyly, W. J. P. The life-history of the Bullhead or Miller's Thumb (*Cottus gobio* L.). Proc. zool. Soc. Lond. 128: 431–453 5 figs.

Sochurek, E. Proterohinus marmoratus Pallas—eine Meergrundel vom Neusiedler See. Natur u Land 43:49.

(Soin, S. G.). [On the respiratory importance of carotinoid pigment in the eggs of salmonoid fishes and of other forms belonging to the order Clupeiformes]. Zool. Zh., Moscow 35 1956: 1362–1369 2 plts. [Russian with English summary.]

Sparta, A. see Padoa, E.

Sperry, R. W. see Arora, H. L.

Spiecker, D. Die Galvanotaxis der Fische und ihre Auslösung auf Grund unmittelbarer Irritabilität der Rumpfmuskulatur. Zool. Jb. (Allg. Zool.) 67: 229-260 21 figs. Spillman, J. see Guibé, J.

Springer, S. (1). An outline for a Trinidad shark fishery. Proc. Gulf & Carib. Fish. Inst. 1949: [1-10].

Springer, S. (2). Some observations on the behaviour of schools of fishes in the Gulf of Mexico and adjacent waters. Ecology 38:166-171.

Springer, S. (3). Tuna resources of the tropical and sub-tropical Western Atlantic. Trans. Amer. Fish. Soc. 85 (1955):13-17.

Springer, S. & Bullis, H. R. Collections by the 'Oregon' in the Gulf of Mexico. Sp. Sci. Rep. Fish. No. 196 1956: 1-134.

Springer, S. & Thompson, J. R. Night sharks, *Hypoprion*, from the Gulf of Mexico and the Straits of Florida. Copeia 1957: 160.

Springer, V. G. A new genus and species of Elopid fish (*Laminospondylus transversus*) from the Upper Cretaceous of Texas. Copeia 1957:135–140 1 pl. 1 fig.

Springer, V. G. see Hubbs, C.

Stanley-Jones, D. Axial rotation and the origin of vertebrates. Amer. Nat., 90 1956: 361-368.

Starmach, K. Characteristic of rivers from biological and fishery point of view. Polsk. Arch. Hydrobiol. 3 1956: 307–332 4 figs. [English summary.]

Starr, T. J. & Fosberg, W. Filter paper electrophoresis of serum proteins from sharks. Copeia 1957 292-295 2 figs.

Stefan, M. La consommation d'oxygène individuelle et en groupe chez quelques poissons marins et d'eau douce. Bul. Inst. Cercet. Pisc. Roman. 16 (1): 73-81.

Steffens, W. see Menzel, H. V.

Steinitz, H. & Ben-Tuvia, A. The hybrid of Barbus longiceps C.V. and Varicorhinus damascinus C.V. (Cyprinidae, Teleostei). Bull. Res. Counc. Israel 6B: 176-188 19 figs.

Steinmetz, C. see Lagler, K. F.

Steopol, I. see Busnita, T.

Stephen, A. C., Rae, B. B. & Wilson, E. Rare marine invertebrates recently found in the Scottish area. Scott. Nat. 69 No. 3:1-4.

Stephenson, W. & Grant, W. M. Experiments upon impounded callop or yellowbelly (*Plectroplites ambiguus* (Richardson)) at Somerset Dam. Lehthyol. Notes Qd. 1:73-116 1 pl. 10 figs.

Sterba, G. Über die Struktur der Eihüllen bei einige Knockenfischen. Naturwissenschaften 44:187 2 figs.

Stewart, R. N. The migratory char of Iceland. Salm. Trout Mag. No. 151: 239-244.

Stinton, F. C. (1). Fish otoliths from the London Clay of Bognor Regis, Sussex. Proc. Geol. Ass. Lond. 67 1 & 2:15-31 pl. 2 text-fig. 1.

Stinton, F. C. (2). Teleostean otoliths from the Tertiary of New Zealand. Trans. roy. Soc. N.Z. 84:513-517 l pl.

Stokely, P. S. see Dineen, C. F.

Stolk, A. (1). Changes in the pituitary gland of the cichlid *Cichlasoma biocellatum* (Regan) with thyroidal tumour. Proc. Akad. Wet. Amst. **59C** 1956: 494–505 11 figs.

Stolk, A. (2). Polycistic kidneys in the characid *Moenkhausia pittieri* Eigenmann, I–II. Proc. Akad. Wet. Amst. **59C** 1956: 506–519 12 figs.

Stolk, A. (3). Thyroid and pituitary gland in a goldfish (*Carassius auratus*) with acromegaly. Proc. Akad. Wet. Amst. **59C** 1956: 611–623 8 figs.

Stolk, A. (4). Tumours of fishes, XI. Carcinoma of the skin in the anabantid Colisa labiosa (Day). Proc. Akad. Wet. Amst. 596 1956: 624-633 9 figs.

Stolk, A. (5). Tumours of fishes. XII–XIII. Proc. Akad. Wet. Amst. 60C: 31–52 21 figs.

Stolk, A. (6). Oval cells with clubshaped inclusions in the connective tissue of the cichlid *Pterophyllum scalare* Cuvier. Proc. Akad. Wet. Amst. 60C: 53-60 8 figs.

Stolk, A. (7). Tumours of fishes. XIV-XV. Proc. Akad. Wet. Amst. 60C: 185-211 23 figs.

Stolk, A. (8). Multiple cysts in the central nervous system and the optic nerves of the cichlid *Cichlasoma facetum* (Jenyns). Proc. Akad. Wet. Amst. 60C: 338-348 01 figs.

Stolk, A. (9). Cerebral cysts, abnormal pituitary gland and changes of the thyroid gland in the viviparous cyprinodont *Lebistes reticulatus* (Peters). I-II. Proc. Akad. Wet. Amst. 60C: 349–363 14 figs.

Stolk, A. (10). Tumours of fishes. XVI. Fibroma of the intestine in the viviparous cyprinodont *Lebistes reticulatus* (Peters). Proc. Akad. Wet. Amst. 60C: 364-375 13 figs.

Stolk, A. (11). Pharyngeal glands in the cichlid *Haplochromis multicolor* (Hilgendorf). Proc. Akad. Wet. Amst. 60C: 567-577 15 figs.

Stolk, A. (12). Tumours in fishes. XVII-XIX. Proc. Akad. Wet. Amst. 60C: 640-665 25 figs.

Storey, M. H. see Myers, G. S.

Strasburg, D. W. Notes on the respiration of small Remora remora. Copeia 1957: 58-60.

Strawn, K. see Hubbs, C.

Stutinsky, F. La neurosécrétion chez les vertébrés. Année biol. (3) 29 1953 : 487-516.

Sulit, J. I., Navarro, O. B., San Juan, R. C. & Caldito, E. B. Proximate chemical composition of various species of Philippine market fishes. Philipp. J. Fish. 2 1954: 109–123 1 fig.

Sullivan, B. J. see Musacchia, X. J.

Sundararaj, B. I. see Ramaswami,

Suomalainen, P. & Pihlgren, A. On the thiaminase activity of fish and some other animals and on the preservation of thiaminase in silage made from fish. Acta Agral. fenn. 83 1955: 221-229.

Suttkus, R. D. (1). Age and growth of a small stream population of "stunted" smallmouth black bass, *Micropterus dolomieu dolomieu* (Lacépède). N.Y. Fish Game J. 2 1955: 83-94 1 fig.

Suttkus, R. D. (2). First record of the mountain mullet, Agonostomus monticola (Bancroft) in Louisiana. Proc. La Acad. Sci. 19 1956: 43-46.

Suttkus, R. D. (3). Early life history of the Gulf menhaden, Brevoortia patronus, in Louisiana. Trans. 21st N. Amer. Wldl. Conf. 1956: 390–407 6 pls. 8 figs.

Suzuki, H. see Hayashi, S.

Suzuki, K. see Okada, Y.

Suzuki, R. (1). [Development of carp eggs with some foreign sperms (preliminary report).] Jap. J. Ichthyol. 5:114-117 1 fig. [English summary.]

Suzuki, R. (2). Development of goldfish eggs inseminated with some foreign sperms. (A preliminary report). Zool. Mag., Tokyo 66: 34-37.

Svärdson, G. (1). The Coregonid problem. VI. The Palearctic species and their intergrades. Rep. Inst. Freshwater Res. Drottningholm No. 38:267-356.

Svärdson, G. (2). The salmon and the climate. Rep. Inst. Freshwater Res. Drottningholm No. 38: 357–384. [English summary.]

Svetovidov, A. N. (1). On the correlation between the number of pyloric caeca and the type of food in fishes. Notes on general problems of ichthyology 19 1952: 282–289. [In Russian.]

Svetovidov, A. N. (2). Some features of resemblance and difference in the distribution, ecology and other characters between *Gadus morrhua* and *Clupea harengus*. Notes on general problems of ichthyology 20 1953: 122—133 2 figs. [In Russian.]

Svetovidov, A. N. (3). The Atlantic origin of some groups of fishes. Int. Congr. Zool. 14 1953 (1956):118-124.

Svetovidov, A. N. (4). Morphological principles of the classification of the Gadidae. Int. Congr. Zool. 14 1953 (1956): 535-540.

Svetovidov, A. N. see Kirpichnikov, V. S.

Svichenskaya, A. A. [New species of mullet from the Lower Miocene of the Caucasus.] Mater. Osnovam. Paleont., 1:61-62.

Swan, M. A. (1). Brown trout of Loch Kernsary. Salm. Trout Mag. No. 150: 121-127 1 pl.

Swan, M. A. (2). Specimens of Char from Shetland and the Faroes. Scot. Nat. 69:67-70.

Swineford, A. see Miller, H. W.

Syazuki, K. & Hayama, K. Studies on nutrition of carp-fry. 3. Consumption of oxygen. J. Shimonoseki Coll. Fish. 6:255-260. [English summary.]

Szabo, S. see Gyurko, S.

Szabo, T. (1). Un noyan particulier dans la formation réticulée bulbaire de certains poissons électriques (Mormyridés). Arch. Anat. micr. Morph. exp. 46:81-91 6 figs.

Szabo, T. (2). Un noyeau particulier dans la formation réticulée bulbaire de quatre poissons électriques appartenant à la famille des Gymnotidae. C.R. Acad. Sci. Paris 245: 1957–1959 1 pl.

Taguchi, K. [Seasonal variation of the good fishing area of salmon and the movements of water masses in the waters of Western North Pacific. III. Relation between the distribution of salmon population and oceanographical conditions mainly in the southern waters of 48° N lat.] Bull. Jap. Soc. Sci. Fish. 22: 609–617 8 figs. [English summary.]

Taguchi, S. see Ishida, J.

Tait, J. S. Nitrogen and argon in salmonoid swimbladders. Canad. J. Zool. 34 1956: 58-62 1 fig.

Takahasi, N. On the so-called accessory respiratory organ "gill-helix" found in some clupeiform fishes, with special reference to its function and its genealogy. Jap. J. Ichthyol. 5: 71-77 l pl.

Talbot, F. H. The fishes of the genus *Lutianus* of the East African coast. Ann. Mag. nat. Hist. (12) 10: 241-258 9 pls.

(Tambs-Lyuke, H.) [Development of the sea-fishery investigations in Norway.] Ichth. Probl. Moscow 8:19-24. [In Russian.]

Tamura, T. (1). A study of visual perception in fish, especially on resolving power and accommodation. Bull. Jap. Soc. sci. Fish. 22: 536-557 12 figs.

Tamura, T. (2). On the relation between the intensity of illumination and the shifting of cones in the fish retina. Bull. Jap. Soc. sci. Fish. 22: 742-746 6 figs. [English summary.]

Tanonaka, G. K. The occurrence of the square-tail, *Tetragonurus cuvieri*, on the high seas south of the Aleutian islands. Copeia 1957: 53-54.

Tarlo, L. B. A preliminary note on new ostracoderms from the lower Devonian (Emsian) of Central Poland. Acta palaeont. Polon. 2:225-234 1 pl. Teichmann, H. Das Reichvermögen des Aales (Anguilla anguilla L.). Naturwissenschaften 44: 242.

Teodorescu-Leonte, R. & others. Quelques données sur la reproduction de Caspialosa pontica Eichw. dans le Danube. Bul. Inst. Cercet. Pisc. Roman. 16 (1):37-46. [French summary.]

Terpenning, J. G. see Larkin, P. A.

Tesch, F. W. Percidenwachstum in eutrophen norddeutschen Flachseen. Z. Fisch. 5:593-606 11 figs.

Tesch, F. W. see Schindowski, E.

Thenius, E. see Papp, A.

Thienemann, A. Die Fische der Deutschen Limnologischen Sunda-Expedition. Arch. Hydrobiol. Suppl. 23:471-477.

Thines, G. & Kähling, J. Untersuchungen über die Farbempfindlichkeit des Höhlenfisches *Anoptichthys jordani* Hubbs und Innes (Characidae). Zeitschr. Biol. 109: 150–160 3 figs.

Thomopoulos, A. (1). Sur quelques oeufs planctoniques de téléostéens de la Baie de Villefranche. II. Pêches du mois de Septembre. Bull. Inst. océanogr. Monaco No. 1072 1956: 1–162 figs.

Thomopoulos, A. (2). Sur la ligne latérale de téléostéens. I. Coecula (Sphagebranchus) imberbis De la Roche et Ophidion barbatum Klein à l'éclosion. Bull. Soc. zool. France 82:102-107 3 figs.

Thomopoulos, A. (3). Influence de la lumière sur le mouvement de l'oeuf de Truite (Salmo irideus Gib.). C.R. Soc. Biol., Paris 150: 2091-2093.

Thomopoulos, A. & Bauchot, R. Sur Cottus bubalis Euphrasen. I. Reproduction en aquarium et développement embryonnaire. Bull. Soc. 2001. France 82:120-126. 2 figs.

Thompson, J. R. see Springer, S.

Tibbo, S. N. see Leim, A. H.

Tiews, K. Biologische Unterschungen am Roten Thun (*Thunnus thynnus* [Linnaeus]) in der Nordsee. Ber. Dtsch. Wiss, Komm, Meeresforsch. 14:192-220.

Tiews, K. see Lillelund, K.

(Tikhonov, V. I.) [A new commercial fish in the Black Sea]. Priroda 5: 102–104 2 figs.

(Tkacheva, K. S.) [Reproduction of Trahurus trachurus L. in Black Sea]. Ichth. Probl., Moscow 8:51-54 3 figs. [In Russian.]

Točko, M. Développement de Chondrostoma nasus ohridanus Kar. Rec. Trav. Sta. Hydrobiol. Ohrid 3 1955: 41-54 17 figs.

Todicescu, S. see Picos, C. A.

(Tokarev, A. K.) [Peculiarities on searching for food and character of feeding of shoals of pelagic fishes.] Trud. Sovesh. Ikhtiol. Komm. 5 1953 [1955]: 158-166 5 figs. [In Russian.]

Tomiyama, I. & Abe, T. Figures and descriptions of the fishes of Japan. Tokyo 56: 1141-1169 3 pls.

Toner, E. D. Movements of salmon around Ireland—VII: From Rath. Co. Kerry (1950–1953). Proc. roy. Irish Acad. 58B: 321–328 2 figs.

Tortonese, E. Spedizione subacquea Italiana nel Mar Rosso, Ricerche zoologiche. VIII. Pesci Isospondili; Apodi, Sinentognati; Eterosomi e Discocefali. Riv. Biol. colon. 15: 49–55.

Tortonese, E. see Padoa, E.

Tosko, M. Développement de Chondrostoma nasus ohridanus Kar. Fac. phil. Univ. Skopje 3 1955: 41-54 17 figs.

Travassos, H. & Pinto, S. Y. Ictio-fauna de Pirassununga. III. Familia Cichlidae (Perciformes-Actinopterygii). Bol. Mus. nac. Rio. de J. Zool. No. 169: 1-16 19 figs.

Trites, R. W. see Leim, A. H.

Trochanis, S. The incredible foureyed *Anableps dovii*. Aquarium, Philad. 26:149-152 3 figs.

(Troitzkii, S. K.) [Reproduction and growth of the yearlings of *Aspius* in Kuban river.] Ichth. Probl., Moscow 7 1956: 134–138. [In Russian.]

Troshin, A. S. see Kirpichnikov, V. S.

Trout, G. C. The Bear Island cod: migrations and movements. Fish. Invest. London (2) 21 No. 6:1-51 40 figs.

Tschörtner, U. Untersuchungen über den Einfluss einiger Milieufaktoren auf die Entwicklung des Hechtes (Esox lucius L.). Arch. Hydrobiol. Suppl. 24 1956: 123–152 3 pls. 56 figs.

Tucker, D. W. Studies on the Trichiuroid fishes.—IV. A specimen of Evoxymetopon taeniatus (Poey) Gill, from the Gulf of Mexico. Ann. Mag. nat. Hist. (12) 10:425-428 1 fig.

(Tugarina, P. Y.) [Some data on reproduction of *Thymallus arcticus baicalensis*]. Zool. Zh. Moscow **35** 1956: 938–939. [In Russian.]

Turchini, J. Recherches histochimiques sur les corps suprarénoux des sélaciens. Ann. Histochém. 1 1956: 78-80.

Turner, C. L. The breeding cycle of the South American fish, *Jenynsia lineata*, in the Northern Hemisphere. Copeia 1957: 195-203.

Tuurala, O. Über den Einfluss der osmotischen Belastung auf die Neurosekretion der Kleinfische Gasterosteus aculeatus L. und Phoxinus laevis L. aus dem Brackwasser des Finnischen Meerbusens. Ann. Acad. sci. Fenn. Helsinki (A) IV Biol. No. 36: 1–9 pl. i.

Twomey, E. Trout of Caragh Lake, Co. Kerry, Ireland. Salm. Trout Mag. 1956: 112-119 3 figs.

Twomey, E. see Went, A. E. J.

Uchida, I. see Shiraishi, Y.

Underhill, J. C. The distribution of Minnesota minnows and darters in relation to Pleistocene glaciation. Occ. Pap. Minn. Mus. nat. Hist. No. 7: 1-45 maps.

Upton, G. V. see Bondy, P. K.

Vaas, K. F. On the nutritional relationship between plankton and fish in Indonesian freshwater ponds. Symposium on marine and freshwater plankton in the Indo-Pacific (Madras 1954) 1–8 (sep.page.)

Valérien, J. L'anhydrase carbonique et la production des gaz de la vessie natatoire de Serranus cabrilla (L.) et S. scriba (L.) Vie et Milieu 8:23-57 2 pls. 11 figs.

Valimitjana, L. & Gadea, E. Estudios histológicos del atún (*Thynnus thynnus* L.). Publ. Inst. Biol. apl. Barcelona 23 1956: 5-19 14 figs. [English summary.] Vallowe, H. H. Sexual differentiation in the teleost fish *Xiphophorus hellerii*, as modified by experimental treatment. Biol. Bull. Woods Hole, 112: 422-429.

Van Bemmel, A. C. V. see Bemmel, A. C. V. van.

Van Campen, W. & Hoven, E. E. Tunas and tuna fisheries of the world: an annotated bibliography, 1930-53. Fish. Bull. U.S., 57 1956:173-249.

Van Dam, L. see Scholander, P. F.

Van de Kamer, J. C. see Kamer, J. C. van de.

Van der Heide, S. see Heide, S. van der.

Van Oosten, J. see Brown, M. E.

Vargas, F. & Concha, J. (1). Histologia y técnica de extirpación de las glándulas adrenales en el Teleósteo Sicyases sanguineus. Invest. Zool. Chil. 3:81-87 5 figs.

Vargas, F. & Concha, J. (2). Fisiologia de las glándulas adrenales en ál Teleósteo, *Sicyases sanguineus*. Invest. Zool, Chil. 3:88-95.

Vargas, F. F. see Concha, B. J.

Vasiliu, G. D. & Costea, E. Contribution à la connaissance de la biologie de la nutrition chez *Vimba vimba* natio carinata L. dans les eaux de la Roumanie. Notul. biol., București 5 1947: 229–236.

Vasiliu, G. D. & Nicolau, A. (1). Nouvelles données sur la nutrition naturelle de la Sandre (Sander lucioperca L.) dans les eaux de la Roumanie. Notul. biol. București. 5 1947 : 181–198 2 pls.

Vasiliu, G. D. & Nicolau, A. (2). Recherches sur la biologie de la nutrition naturelle chez Aspius aspius L. dans les eaux de la Roumanie. Notul. biol. București 5 1947: 250–261.

Vasiliu-Suceveanu, N. see Radulescu, I.

(Vasnetzov, V. V. & others). [The stages of development of semi-migratory fishes of the Volga and Don, Abramis brama, Cyprinus carpio, Rutilus rutilus caspicus, Rutilus r. heckeli and Lucioperca lucioperca]. Trud. Inst. morf. Zhiv. 16:7-76 5 figs. [In Russian.]

Vengrzhin, E. P. see Shulman, G. E.

Verigin, B. V. [Structure of the gill apparatus and the superbranchial organ of Hypophthalmichthys molitrix Val.] Zool. Zh., Moseow 36:595-602 4 pl. 9 figs. [English summary.]

Vernon, E. H. Morphometric comparison of three races of kokanee (Oncorhynchus nerka) within a large British Columbia lake. J. Fish. Res. Bd. Can. 14: 573-598 l fig.

Vernon, E. H. & McMynn, R. G. Scale characteristics of yearling coastal cutthroat and steelhead trout. J. Fish. Res. Bd. Can., 14: 203-212 6 figs.

Vervoort, W. Note on the pituitary cytology of Lebistes reticulatus (Peters) I-II. Proc. Akad. Wet. Amst. 60C: 486-509 2 pls. 4 figs.

Vialli, M. La quantita' di acido desosscribonucleia per nucleo negli entroati' di *Latimeria*. R.C. Ist. Lombardo Cl. Sei. Mat. Nat. 91: 680–686.

Vialli, M. see Padoa, E.

Vickers, K. U. see Went, A. E. J.

Vignais, P. see Grangaud, R.

Villadolid, D. V. see Blanco, G. J.

Villadolid, D. V. see Rasalan, S. B.

Villaluz, D. K. see Rabanal, H. R.

Villmitjana, L. Contribución al estudios e los spermatozoides de teléostes. Bol. Soc. esp. Hist. nat. Sec. Biol. 55:185-192 2 pls.

Vilter, V. Organisation remarquable de la rétine du poisson abyssal *Bathylagus benedicti*. Int. Congr. Zool. 14 1953 (1956): 290.

(Vladimirskaya, M. I.) (1). [The white fish of lake Imandra.] Ichth. Probl. Moscow 6 1956:136-148. [In Russian.]

Vladimirskaya, M. I. (2). Lake and river trout in the basin of lake Imandra. Bull. Soc. Nat. Moscou, Biol. 62 No. 4: 37-50. [English summary.]

Vladimirskaya, M. I. (3). [Grayling from the lakes of the North-Western area of the Lake Imandra basin.] Zool. Zh., Moscow 36:729-736 1 pl. [English summary.]

Vladykov, V. D. (1). Distant recaptures of shad (Alosa sapidissima) tagged in Quebec. Nat. canad. 83 1956: 235-249 4 figs.

Vladykov, V. D. (2). Les formes locales de la truite rouge du Québec (Salvelinus marstoni) (1). Nat. canad. 84: 233-248.

Vokes, H. E. Miocene fossils of Maryland. Bull. Maryland Dep. Geol. 20: VII, 1-85, 31 pls.

(Volodin, V. M.) (Embryonic development of the autumn Clupea harengus membras and the oxygen demand during this development.] Ichth. Probl., Moscou 7 1956:123-133 3 figs. [In Russian.]

Volpe, A. V. Fingerprinting fish. Bull. int. oceanogr. Fdn. 2 1956: 135–142.

Vondett, J. E. see Whalls, M. J.

Vorderwinkler, W. see Axelrod, H. R.

Voss, G. L. & Robins, C. R. Progress report No. 3. Summary of investigations for the period comprising September 1956 through April, 1957. Charles F. Johnson Oceanic Gamefish Invest. Univ. Miami 57-15 (1957) pp. 1-11 (mimeographed).

Voss, N. A. Fishes of the family Gempylidae collected by the Bermuda Oceanographic Expeditions 1929, 1930, 1931 and 1934. Copeia 1957: 304–305.

Wallin, O. On the growth structure and developmental physiology of the scale of fishes. Rep. Inst. Freshwater Res. Drottningholm No. 38:385-447 5 figs. 32 pls.

Walters, V. (1). Protolophotus, a new genus of allotriognath fish from the Oligocene of Iran. Copeia 1957: 60-61.

Walters, V. (2). Alphestes scholanderi, a new sea bass from the West Indies. Copeia 1957: 283–286 4 figs.

Warburton, B. & others. Reproductive behavior of *Gambusia heterochir*. Copeia **1957**: 299–300.

Ward, J. W. The reproduction and early development of the sea catfish, *Galeichthys felis*, in the Biloxi (Mississippi) Bay. Copeia **1957**: 295–298.

Warren, F. J. see Denton, E. J.

Weeks, L. G. Origin of carbonate concretions in Shales, Magdalena Valley, Colombia. Bull. geol. Soc. Amer. 68: 95-102 4 pls.

Weiler, W. (1). Fischreste aus dem Tertiär von Epterode südlich von Grossalmerode. Notizbl. hess. Landesamt. Bodenforsch. 85: 24–27 9 figs.

Weiler, W. (2). Zur fischfauna des Dobergs bei Bünde in Westfalen. Paläont. Z. 31: 135–138.

Weiler, W. (3). in Martin, G. P. R. & Weiler, W. Das Aldorfer Otolithen—
"Pflaster" und seine Fauna. (Mittlerer Münder Mergel, Malm.). Senckenbergiana leth. 38: 211–249 3 pls. 2 figs.

Welander, A. D., Alverson, D. L. & Bergman, P. Rare fishes from the Eastern North Pacific Ocean. Fish. Res. Pap. Wash. 2:1-72 figs.

Welander, A. D. & others. Occurrence of the boar fish, *Pseudopentaceros richardsoni*, and the zeid, *Allocyttus verrucosus* in the North Pacific. Copeia 1957: 244-245.

Went, A. E. J. (1). Salmon movements around Ireland. VI. From Portballintrae, Co. Antrim. (1953 and 1954). Proc. R. Irish Acad. 58B 2 1956: 17-21 1 fig.

Went, A. E. J. (2). Sea trout of the Cashla river with notes on the salmon. Salm. Trout Mag. 1956: 63-76 3 figs.

Went, A. E. J. (3). The pike in Ireland. Irish Nat. J. 12:177-178 2 figs.

Went, A. E. J. (4). A further specimen of dusky perch, *Epinephelus gigas* Brunnich from Irish waters. Irish Nat. J. 12: 202.

Went, A. E. J. (5). Biology of the salmon. Nature London 180: 789-790.

Went, A. E. J. (6). Sea trout of the River Ilen. Salm. Trout Mag. No. 150: 139-147.

Went, A. E. J. & Twomey, E. Notes on the Irish Char, Salvelinus colii Gunther: VI. Irish Nat. J. 12:205-206.

Went, A. E. J. & Vickers, K. U. Salmon and trout research on the river Foyle. 3rd Ann. Rep. Foyle Fish. Comm. App. III. 1955: 1–81 fig.

Whalls, M. J., Shetter, D. S. & Vondett, J. E. A simplified rotary fish screen and an automatic water gate. Trans. Amer. Fish. Soc. 86 (1956): 371–380 4 figs.

Wheeler, A. C. Syngnathus rostellatus as a British fish. Essex Nat. 30:56.

Wheeler, J. F. G. The problem of poisonous fishes. Rev. Agric. Ile Maurice 33 1954:132–139.

Whiting, H. P. Mauthner neurones in young larval lampreys (Lampetra spp.). Quart. J. micr. Sci. 98: 163-178 7 figs.

Whitley, G. P. (1). The freshwater fishes of Australia. Australasian Aqualife, Sydney 1 1956 No. 9: 9–12 no. 10: 9–12.

Whitley, G. P. (2). Ichthyological illustrations. Proc. R. Zool, Soc. N.S.W. 1955-56:56-71 12 figs.

Wickler, W. (1). Ei und Larve von Symphysodon discus Heckel. Aquar. Terrar. Z. 10: 7-12 20 figs.

Wickler, W. (2). Die Larve von Blennius fluviatilis Asso 1784. Biol. Zbl. 76: 453-466 13 figs.

Wickler, W. (3). Das Kampfverhalten von Nannostomus beckfordi aripirangensis (Pisces-Cypriniformes-Erythrinidae). Bull. aquat. Biol. 1: 13-15.

Wickler, W. (4). Das Ei und die Embryonalentwicklung bei Fischen. Mikrokosmos 46: 217–220 6 figs.

Wiepkema, P. R. Het spiel van de Bittervoorns. Levende Nat. 60: 169-177 8 figs.

Wilber, C. G. see Musacchia, X. J.

Wilimovsky, N. J. see Mohr, J. L.

Willgohs, J. F. Some notes on *Beryx decadactylus* (Cuv. & Val.) in European waters with a special account of Norwegian records. Arb. Univ. Bergen 1956 No. 15: 1–13 2 figs.

Williams, F. Preliminary survey of the pelagic fishes of East Africa. Fish. Publ. Lond. No. 8 1956: 1-68 15 pls.

Williams, F. & Newell, B. S. Notes on the biology of the dorade or dolphin fish (*Coryphaena hippurus*) in East African waters. E. Afr. Agric. J. 23: 113-118 3 figs.

Wilson, E. see Rae, B. B.

Wilson, E. see Stephen, A. C.

Winn, H. E. Egg site selection by three species of darters (Pisces: Percidae). Brit. J. Anim. Behav. 5: 25-28. Wohlschlag, D. E. Information from studies of marked fishes in the Alaskan Arctic. Copeia 1956 No. 4 [1957]: 237-242 1 fig.

Wolfsheimer, G. A spawning of porcupine puffers. Aquarium, Philad. **26**: 288-290 1 fig.

Wood, E. M. & Yasutake, W. T. Histopathology of kidney disease in fish. Amer. J. Path. 32 1956: 845-857 18 figs.

Wood, F. G. Southern extension of the known range of the basking shark, *Cetorhinus maximus* (Gunnerus). Copeia 1957: 153-154.

Woodhead, P. M. J. Reactions of Salmonid larvae to light. J. exp. Biol. 34:402-416 7 figs.

Woods, L. P. Beryx splendens Lowe in the Gulf of Mexico. Copeia 1957: 298-299.

Woods, L. P. & Inger, R. F. The cave, spring and swamp fishes of the family Amblyopsidae of Central and Eastern United States. Amer. Midl. Nat. 58: 232-256 11 figs.

Woolcott, W. S. Comparative osteology of serranid fishes of the genus *Roccus* (Mitchill). Copeia 1957: 1–10 2 pls.

Worbe, J. F. see Cordier, D.

Wunder, W. Über den Bau der Netzhaut beim Rotbarsch (Sebastes marinus L.). Verh. dtsch. zool. Ges. 1956: 95-96.

Yakovlev, V. N. [First occurrence of the genus *Scardinius* of the family Clupeidae in the Upper Tertiary deposits of Siberia.] Mater. Osnovam Paleont. 1:63-65. [In Russian.]

Yamaguchi, M. see Kuronuma, K.

Yamamoto, K. Studies on the formation of fish eggs. X. Demonstration of carbohydrates related to Da-Fanopositive bodies in the oocyte of the flounder, *Liopsetta obscura*. Annot. zool. jap. 30: 33–37 10 figs.

Yamamoto, T. S. [Histochemical study on the early development of the herring]. Zool. Mag. Tokyo 66: 289–294 5 figs. [English summary.]

Yamamoto, T. S. see Kanoh, Y.

Yanagimachi, R. (1). Studies of fertilization in Clupea pallasii. I. Extension of fertilizable life of the unfertilized eggs by means of Isotonic Ringer's solution. Zool. Mag., Tokyo 66: 218-221. [English summary.]

Yanagimachi, R. (2). Studies of fertilization in Clupea pallasii. II. Structure and activity of spermatozoa. Zool. Mag., Tokyo 66: 222-225 2 figs.

Yanagimachi, R. (3). Studies of fertilization in Clupea pallasii. III. Manner of sperm entrance into the egg. Zool. Mag., Tokyo 66: 226-233. 9 figs. [English summary.]

Yanagishima, S., & Mori, S. [Studies on the variation and adaptation of fishes. I. Adaptation of killifish (Oryzias latipes T. & S. 1. Field survey. 3. Experimental study]. Zool. Mag., Tokyo 66:351-366 9 figs. [English summary.]

Yarmoshenko, N. P. see Movchan, V. A.

Yasutake, W. T. see Wood, E. M.

Yeh, S. see Liu, H.

Yoshida, H. & Ito, T. Fish fauna of the Japan Sea. J. Shimonoseki Coll. Fish. 6: 261-270.

Yoshida, H. O. see Otsu, T.

Yoshihara, T. & Fujioka, J. [The toxicity of chemicals on fish. II. Estimation of the effectiveness of insecticides on the young carp, Cyprinus carpio Linné.] Bull. Jap. Soc. Sci. Fish. 22:641-644 3 figs. [English summary.]

Yoshizaki, M. [Morphological change of intestine in the larval stage of wild goldfish (*Carassius auratus*). I. Observation on variety "Gengorobuna".] Jap. J. Ichthyol. 5: 78–82 1 pl. 5 figs. [English summary.]

(Yudanov, I. G.) [New data on the spawning places of Atlantic-Scandinavian herrings.] C.R. Acad. Sci. URSS 112:1139-1141. [In Russian.]

Yuen, H. S. H. & June, F. C. Yellowfin tuna spawning in the central equatorial Pacific. Fish. Bull. U.S. 57: 251-264.

Zahnd, J. P. see Clavert, J.

(Zaitzev, Y. P.) [Diagrammatic representation of the occurrence of pelagic eggs of some Black Sea fishes.] C.R. Acad. Sci. U.R.S.S. 111 1956: 193-194 1 fig. [In Russian.]

Zambriborshch, F. S. [Structure and function of the superbranchial organ of *Hypophthalmichthys molitrix* Val.] Zool. Zh., Moscow 36: 587-594. [English summary.]

Zanandrea, G. Neoteny in a lamprey. Nature, London 179: 925-926.

Zarnecki, S. see Rychlicki, Z.

Zelenin, A. M. see Irkimovich, A. I.

Zenkevitch, L. Neue Vertreter der Mittelmeer-fauna im Kaspischen Meer. Int. Congr. Zool. 14 1953 (1956): 113– 118.

(Zheltenkova, M. V.) (1). [Critical consideration of modern methods of fish feeding in natural environment.] Trud. Sovesh. Ikhtiol. Komm. 6 1954 (1955): 22-39 7 figs. [In Russian.]

(Zheltenkova, M. V.) (2). [Prognosis on fish feeding conditions in Azov Sea.] Ichth. Probl., Moscou 8:102–106. [In Russian.]

(Zhukov, L. A.) [Tests in the pond breeding of Vioksinks *Coregonus* for its acclimatisation in small lakes.] Trud. Lab. Ozerov. 5:270-275. [In Russian.]

(Zhuravlev, M.) [Influence of "black water" on fish in the river Kura in the Mingechaur region.] C.R. Acad. Sci. Azerbaidj 10 1954: 585-588. [In Russian.]

(Zusser, S. G.) [Critical remarks on the theory of tropisms as applied to the study of fish behaviour.] Trud. Sovest. Ikhtiol. Komm. 5 1953 (1955): 154– 155. [In Russian.]

II.—SUBJECT INDEX

GENERAL

Text-books.—Physiology of fishes, Brown, M. & others; determination of fossils, Chavan & Cailleux.

Popular books.—Freshwater fishes, Schindler (2); fishes of the world, Le Danois & others; natural history of Israeli seashore, Carmin; tropical fish breeding, Axelrod & Vorderwinkler. Biography and bibliography.—Bibliography of natural history works of Seychelles, Peters; of Philippine fishes, Blanco & Montalban; of tunas and their fishery, Van Campen & Hoven; method of documentation of fishery publications, Lillelund & Tiews; an early article on the swim bladder, Nelson; biographical notes on the Russian pisciculturist, O. A. Grimm, Movchan & Yarmoshenko; obituary of V. Pietschmann, Kähsbauer.

Historical.—Notes on some drawings ascribed to J. P. Kirtland, Moulton; iehthyology in China, Nikolski (3).

Expeditions and Collections.—Gempylids from the Bermuda Oceanographic Exped. 1929-31 and 1934, Voss, N. A.; "Oregon" collections from Gulf of Mexico, Springer, S. & Bullis; Carapidae of Monaco Oceanographic Inst., Arnold; method of filing specimens at Stanford Univ., Myers & Storey.

Techniques.—Method of making fish prints, Goodwin; use and construction of large fish traps, Hallock & others; use of tricaine as a fish tranquillizer, Gossington; method of sectioning small fish, Sauerteig & Krebs; methods of measuring external characters, Amosov; of fish tagging, Volpe; a subcutaneous plastic tag, Butler; method of studying feeding habits, Borutzki; Fortunatova (1); method of studying food requirements and nutrition, Karzinkin & Krivobok: method of calculating back growth by correlation curve, Schindowski & Tesch; sampling methods for determining size composition of catches, Hennemuth; method of removing egg membrane, Kanoh & Yamamoto; technique for adrenalectomy, Vargas & Concha (1); of collecting fish pituitaries, Pickford (1); use of chromatography in taxonomy, Rechnitzer.

ECONOMICS

Fishery research.—Izhevski; in Norway, Tambs-Lyuke; in Barents Sea, Lee; on Iceland-Faroe Ridge, Magnússon (2); tuna resources of Western Atlantic, Springer, S. (3); survey of pelagic fishes of E. Africa, Williams; population dynamics and economics, Schaefer.

Overfishing.—Of Neothunnus population, Broadhead; of Cetengraulis as tuna baitfish, Alverson, F. G. & Shimada; effect of heavy fishing on Abramis populations, Bervald.

Fishes and control of disease.— Haplochromis used for molluse control, Bont (2) & (3).

Effects on fishes of industrial effluents.

—Effect of mill effluent on young Oncorhynchus, Alderdice & Brett; influence of 'black water' on fishes in Kura river, Zhuravlev.

Dams and other obstructions affecting fisheries.—Effects of hydro-electric development on migrating Oncorhynchus, Brett.

Pisciculture and fish management.— In Africa, Lemasson; fish culture in rice-fields, Bont (1); fish production in the Vistula, Poczopko (1); stocking Angare river with Thymallus, Egorov; methods of studying fish productivity of lakes, Hayes; problems of rearing marine fishes from eggs, Morris, R. W.; effect of electric shocks on developing Salmo eggs, Godfrey; fish feeding, Karpevich; biological approach to fish feeding, Gaevskaya; modern methods of feeding, Zheltenkova (1); estimation of feeding conditions in Azov Sea, Zheltenkova (2); feeding of medicated diet to prevent pathogens, Grassl (2); use of pelleted dry food in trout hatcheries, Grassl (1); importance of artificial food for young Cyprinus, Bodrova & others; artificial propagation of Acipenser, Manea & Mihai; of Coregonus in ponds, Zhukov; of Leuciscus and Barbus, Gyurko & others; varying quality of Acipenser fry, Shturbina; results of artificial hybridization in Cyprinus, Lieder; maintenance of Salmo gairdneri stock by selective breeding, Donaldson & Olson; rearing Esox masquinongy in Nogies Creek, Crossman; management of Chanos nurseries in Philippines, Rabanal & others; causes and prevention of mortality during transport, Ahmad; study of survival and vitality of hatchery bred Salmo irideus, Nielson & others; use of PMA in controlling gill disease, Allison: resistance to dropsy obtained by Zator carpbreeding methods, Rychlicki & Zarnecki; control of dropsy in Cyprinus, Ivasik & Roi; experiments on Plectroplites, Stephenson & Grant; factors to be considered in acclimatization, Shkorbatov; behaviour in a fish pass, Schiemenz; poisons used for population study, Oliva (2); a simplified rotary fish screen, Whalls, Shetter & Vondett.

Fishes useful or harmful to man.—Copper acetate used as shark repellent, Hill; injuries caused by garfish, Haneveld.

Poisonous and venomous fishes.—The problem of poisonous fishes, Wheeler, J. F. G.; toxicity of fish, Kawabata, Halstead & Judefind; Trachinus stings and their treatment, Halstead; two fatalities from Synanceja wounds, Smith, J. L. B. (8); observations on Synanceja, Gail & Rageau.

Fishes as test animals.—Effect of radioactive chemicals on *Cyprinus*, Kirpichnikov & others.

Methods of fishing.—Electrical fishing, Denzer; electrical fishing in fresh water, Perruche; gear used for E. African pelagic fishes, Williams; gear for pelagic fishing in Philippines, Rasalan & Villadolid; goby fishing in Philippines, Capeo & Manacop.

Fisheries of.—Barents Sea, Mironova (2); Sweden (Esox and Perca), Alm; Roumania (marine), Busnita & Arachelian; Spain, Doumenge; L. Victoria, Garrod; inland Africa, Lemasson; Ceylon, Canagaratnam & Medcof: Silvalin gam & Medcof; St. Paul and Amsterdam Isl., Paulian; Trinidad (shark), Springer, S. (1); Barbados (flying fish), Hall.

Oceanography.—Of Uruguayan coast, Buen (1); effect of sound waves on vertical distribution of fishes, Beklemishev; interpretation of echo traces, Cushing (1); midwater sound scattering, Johnson, H. R. & others; luminescence in a sonic-scattering layer, Kampa & Boden.

STRUCTURE

General.—External morphology of embryo, Rhincodon, Reid (1).

Tegumentary system.—Fish scales, Grave; growth structure of scales, Wallin; scales of Ophicephalidae, Bhatti & Aminul Huq; scalation in Cyprinus, Dürr; Histology of palaeoniscoid scales, Ørvig (2); crystal orientation in Permian scales, Konta.

Fins.—Dorsal fin of Salmo, François; anal rays of Lebistes, Mohsen; bony rays in adipose fin of Synodontis, Sandon; finlets detached from main fin during ontogeny in Thunnus, Robins (2); significance of selachian fin-spines, Peyer.

Skeleton.—Of Archoplites, Dineen & Stokely; of Nystactes, Böhlke (3); of Roccus, Woolcott; skull of Balistes, Garnaud (1); chondrocranium of Eleutheronema, Marathe & Bal; architecture of head in Pleuronectes, Blok; microstructure of dermal bone in arthrodires, Bystrow; endoskeletal tissues of certain Arthrodires, Ørvig (3); exoskeletal microstructure of certain Holoptychiids, Ørvig (1).

Dentition.—Study of dentine, James; dentition of Mugilidae, Ebeling; microstructure of teeth, etc. of certain Devonian Acanthodians, Arthrodires and Rhipidistia, Gross.

Nervous system.—Brain of fishes related to mode of life, Povlovski; form of synapses in teleosts, Horstmann; Mauthner's fibre in fishes, Leghissa; Mauthner neurones in larval Lampetra, Whiting; subcommissural organ in Lampetra and Myxine, Adam; subcommissural cells and Reissners' fibre of Myxine, Afzelius & Olsson; neurosecretory cells of diencephalon in teleosts, Polenov; torus longitudinalis in some teleosts, Banarescu (1); innervation of membranous labyrinth, Casasco; presence of a paraphysis in Selachians, Kappers; optic nerves of Gambusia, Hamon; a plexus in digestive tract of Perca, Milokhin (2); innervation of some Selachian fin muscles, Barets: brain cast of triassic palaeoniscid, Beltan.

Sense Organs.—Eye of fishes, Baburina; cone arrangement in teleost retinae, Lyall (2); density of cones and resolving power of retina, Tamura (1); retinal pigments in deep-sea fishes, Denton & Warren; retina of Bathylagus, Vilter; retina of Sebastes, Wunder; sensory canals of head in Wallagonia, Kapoor, A. S.; Cutaneous sense organs of Rita, Bhatti (2); barbels of Rita, Bhatti (1).

Luminous organs.—Pre-caudal photo phores of Lampadena, Anadon; photophores of Porichthys, Nicol.

Otoliths.—Of Pliocene teleosts, Papp & Thenius; from London Clay of Bognor Regis, Stinton (1); from Miocene New Zealand, Stinton (2); from Jurassic Western Germany, Weiler (3).

Lateral line.—Pit organ system of some fishes, Satô.

Muscles.—Myomeres of fishes, Marshall, J. R.; lateral muscles of Trachurus, Hasan; muscles of Rutilus, Lobintzev; muscles of skull in Raja, Gottenbos; muscular tissue of auricle in Anguilla, Couteaux & Laurent.

Digestive system.—Alimentary canal of Basilichthys, Carlo & Lopez; pharyngeal pockets in Gadusia, Kapoor, B. G. (3); callous pad of pharyngeal apparatus in Indian Cyprinids, Kapoor, B. G. (4); gastric epithelium in young Leuciscus, Bernard.

Swimbladder.—Of Arapaima, Migdalski; of Ophicephalus, George; unusual swimbladders in Cyprinus, Schreiter.

Glands.—Endocrine organs of Anabantids, Forselius; pineal organ in fishes, Kamer; paired pineal organs, Edinger; hypophysis of some teleosts, Honma (3); of Lebistes, Vervoort; of Stichaeus, Honma (1); adrenal tissues of some fishes, Oguri & Hibiya (2); adrenals of Sicyases, Vargas & Concha (1); pancreas, spleen and liver of Thunnus, Vallmitjana & Gadea; histology of Hippocampus kidney, Gabe; pharyngeal glands of Haplochromis, Stolk (11); Stannius bodies of teleosts, Oguri & Hibiya (1); glands associated with digestive system of Cyprinus, Santa & Pollingher; hatching gland of Huso, Ignatyeva (1).

Electric organs.—Bulbous structure in Gymnotids, Szabo (2); electric discharge in Mormyrids controlled by paraseptal nucleus, Szabo (1).

Respiratory system.—Oral valves of some teleosts, Kapoor, B. G. (2); peculiarities of branchial apparatus in Acipenser, Shmalganzen; gill apparatus of Hypophthalmichthys, Verigin; suprabranchial of Hypophthalmichthys, Zambriborshch; gills of Lampetra, Morris, R.; secretory cells in gills of Mastacembelus, Kapoor, B. G. (1).

Vascular system.—Of liver of Cyprinus, Friedrich; afferent branchials of Ophicephalus, Bhuiya & Siddiqui; intestinal blood system of Petromyzonidae, Baxter.

Reproductive system,—Spermatozoa of teleosts, Villmitjana; gonads of Oncorhynchus, Ivanova, S. A.; urogenital apparatus of 3 and \$\times\$ Sebastes marinus, Magnusson (1); egg membrane of fishes, Sterba.

Teratology and pathology.—Teratological example of Raja, Barraclough; abnormal vertebral column in Leuciscus, Greven; deformation of pectoral fin of Acipenser, Huzum; Esox with no pelvic fins, Herbst: Janus monstrosity in Oncorhynchus, Honma (4); teratological examples of Wallago and Mystus, Sarkar & Kapoor; histopathology of kidney disease, Wood, E. M. & Yasutake; multiple cysts in nervous system of Cichlasoma, Stolk (8); cerebral cysts and abnormal glands in Lebistes, Stolk (9); polycystic kidneys in Moenkhausia, Stolk (2); coarse granulocytes in Pterophyllum, Stolk (6); changes in pituitary of Cichlasoma due to thyroidal tumour, Stolk (1); gonadal tumour in Sardina, Santos Pinto (2); fibroma of heart in Lebistes, Stolk (7); rectal fibroma in Lebistes, Stolk (10); a sarcoma in Pseudocorynopoma, Duijn; renal adenocarcinoma in Barbus, Stolk (7); adenoma of pharyngeal glands in Haplochromis, Stolk (12); adenoma of swim bladder in Lebistes, Stolk (12): carcinoma in Colisa, Stolk (4); carcinoma of kidney in Thayeria; skin fibromas in Malapterurus, Stolk (5); glands of an aeromegalic Carassius, Stolk (3).

Comparative morphology.—Of midbrain of *Protopterus*, **Dorn**; significance of accessory denticles in teeth of Lamnid and Scapanorhynchid sharks, **Ghlikman** (1).

Fossilization.-Breder.

Restorations.—Pharyngolepis, Smith, I. C.; Semionotus capensis, Guffroy.

DEVELOPMENT

General.—Early development of Bregmaceros, Clancey; of Chondrostoma, Točko, Tosko; of Clupea in water of high salinity, Krizhanovski (1); of Esox, Shamardina; of Galeichthys, Ward; of Psenes, Legaspi; of Wallagonia, Mazumdar; influence of light on developmental stages, Lyubitskaya; developmental physiology of scales, Wallin.

Eggs.—Of some teleosts, Wickler (4); Symphysodon, Wickler (1); gynogenesis in Carassius, Busnita & others (2); respiratory importance of pigment in eggs of Salmonoids, Soin; effect of light on movement of Salmo egg, Thomopoulos (3); development of Salmo egg in absence of oxygen, Devillers & others; development of perivitelline

space in oocytes of Acipenseridae, Sadov (1); differentiation of blastoderm after removal of covering layer, Devillers & others; use of Ringer's solution in prolonging life of unfertilized eggs, Yanagimachi (1).

Embryos.—Of Clupea, Volodin; of Cobitis, Kojhanova; of Cottus bubalis, Thomopoulos & Bauchot; of Aral Sea Rutilus, Gosteeva; peculiarities in Vimba, Smirnova, E. N. (1); lactic acid production in Oryzias embryos, Ishida & Taguchi; optokinetic testing of cyclopean Fundulus embryos, Rogers.

Larvae and young.—Of some Volga and Don river fishes, Vasnetzov & others; of Abramis, Rutilus and Lucioperca, Dmitrieva (1); of Alosa kessleri, Braginskaya; of Blennius, Wickler (2); of Odontamblyopus, Dôtu (4); of Pomatomichthys constanciae, Cavaliere; of Symphysodon, Wickler (1); changes in Oncorhynchus fry during absorption of yolk, Shiraishi & Uchida; sexual maturity in ammocoete larvae, Zanandrea.

Organogeny and histogenesis-Origin of vascular couples in Petromyzonidæ, Baxter; of digitiform gland in Plagiostomes, Pera; of scales in Rutilus and Leucaspius, Balon (1); histogenesis of dorsal fin components in Salmo, Français, François; of caudal fin in teleosts, Bhatti; of lateral line in Caecula and Ophidion, Thomopoulos (2); origin of acoustico-facialis complex in Lampetra, Fisk; cell migration in embryonic brain of Torpedo, Bergquist; growth of retina in Salmo, Lyall (1); organogenesis of Salmo trutta, Bernard; gonad differentiation in Amia, Ancona (2); in Salmo trutta, Ashby (1); in Teleosts, Ancona (3); postembryonic development of pronephros in Clupea, Holstvoogd; changes occurring in liver during vitellogenesis, Clavert & Zahnd (1) & (2).

Effect of external factors.—On development of Esox, Tschörtner; influence of light conditions no development of Acipenser larvae, Semenov; influence of temperature on skeleton, Orska; low temperature effects on stimulation of growth of oocytes, Aleev (1).

Regeneration and degeneration.—Regeneration of teleost scales, Aoyama; of caudal fin of *Oryzias*, Mori, H.; of liver in *Salmo*, Faktorovich (1); of

melanomas, Ermin & Gordon; reduction of eye and associated structures in blind cave fishes, Quaghebeur.

Abnormal development.—Transplantation of melanomas, Humm, Clark & Humm; development of Salmo, Esox and Rutilus in paraffin oil, Domurat.

PHYSIOLOGY

Senses.—Visual perception, Tamura (1); adjustment of cones to intensity of illumination, Tomura (2); absorption of light by crystallin, Denton; Motais (2); colour sensitivity in Anoptichthys, Thines & Kähling; experiments on behaviour of blind and blinded fishes, John; scent perception by Anguilla, Teichmann; temperature perception, Bardach & Bjorklund; sensitivity of Carassius to heat stimulation, Bardach; function of ampullae of Lorenzini, Murray.

Nervous system.—Instinct and brain function in Gasterosteus, Segaar; experiments on neural crest of Lampetra, Newth; innervation of photophores in Porichthys, Nicol; two motor systems in lateral muscle of Tinca, Barets & Le Touzé; results of breakage of Reissner's fibre, Olsson; regeneration of cranial nerve after severing, with re-innervation of muscles, Arora & Sperry.

Digestion.—Relation between food and type of nervous system in digestive tract in some Cyprinids, Milokhin (1); nutrition in Cyclostome larvae, Schroll; relation between pyloric caeca and type of food, Svetovidov (1); rôle of leucocytes in digestion, Smirnova, L. I.

Metabolism.—Of Carassius acclimatised to low oxygen concentrations, Prosser & others; in young Cyprinus, Picos; effect of temperature on metabolism in Cyprinus, Picos & Todicescu; metabolism and growth of Salmonid fry at varying temperatures, Mekhanik; metabolism of fishes, Brown, M. & others.

Respiration.—Oxygen consumption of Cyprinus fry, Syazuki & Hayama; dependence of breathing frequency on temperature in Cyprinus, Meuwis & Heuts; cutaneous respiration in Neogobius, Shulman; respiration of Remora, Strasburg; oxygen consumption of some fishes, Stefan; survival at differing concentrations of dissolved O and CO₂, Alabaster, Herbert & Hemens.

Blood. — An isoagglutinin system found in *Ictalurus*, Cushing & Durall; results of an electro-cardiogram of *Tinca*, Serfaty & Raynaud; alterations in lactic acid level following muscular activity, Black (1).

Osmotic relations.—Osmoregulation in Lampetra, Morris, R.; in egg and early stages of Lampetra, Hardisty; permeability of chorion of Fundulus egg, Kao & Chambers, R. & E. L.; effect on Gobius of alteration of salinity, Cordier, Barnoud & Brandon; gas exchange in Neogobius in Azov Sea, Shulman & others; adaptation of Oryzias to salt water, Yanagishima & Mori; osmoregulation in Salvelinus, Gordon, M. S.; effect of hormones on osmoregulation, Lockley; osmotic behaviour of cortical alveoli, Kusa; role of pituitary, Tuurala.

Locomotion and equilibration.—Reaction of blind Carassius to rotation, Jones, F. R. H.; functional importance of lateral body orientation in Heterosomata, Aleev (2); characteristics and functions of fins, Aleev (5); musculature and swimming, Marshall, J. R.

Swim bladder.—Source of oxygen in swim bladder of Gadus, Scholander & others; gas composition in bathypelagic fishes, Kanwisher & Ebeling; analysis of gas content, Tait; metabolism of gases, Valérien.

Influence of temperature.—On hatching gland of *Acipenser*, Ignatyeva (2); on survival in low tensions of oxygen, Downing & Merkens.

Colour.—Formation of carotenoids, Politis (2); formation of melanic pigment, Politis (1); genesis of melanophores and melanin, Pickford & Kosto; effect on chromatophores of induced epilepsy, Branzei & Jitariu.

Endocrinology.—Effect of hormones on euryhalinity of young Anguilla, Motais (1); influence of hormone on halotropism in young Anguilla, Fontaine & Motais; effect of hypophysectomy on respiratory metabolism in Anguilla, Fontaine, Querrière & Raffy; effect of radio-active iodine on thyroid tissue of Anguilla, Olivereau (2); changes occurring in pituitary of Cyprinus during gonad maturation, Irkimovich & Zelenin; response of hypophysectomized Fundulus to thyrotropin, Pickford (2); sexual maturity in presence of unmetamorphosed endostyle in Lampetra

ammocoetes, Zanandrea; neurosecretion in Salmo, Arvy, Fontaine & Gabe; effect of thyroxine on chloride exchange in Salmo, Baraduc; effect of adrenal ectomy on Sicyases, Vargus & Concha (2); neurosecretion in fishes, Stutinsky; hypophysis and thyroid glands of fishes, Olivereau (1); response of hypophysectomized fishes to mammalian and fish growth-hormone, Pickford (1); spawning reflex factor of pituitary, Atz & Pickford; pituitary, nucleus praeopticus, role in osmoregulation, Tuurala; presence of cortisol in adrenal, Bondy, Upton & Pickford; hormones influencing melanogenesis, Pickford & Kosto.

Electric organs.—Properties of electroplax membrane of *Electrophorus*, Altamirans.

Electrical properties.—Effect of galvanic current on heart of Esox, Acolat; 'spike' potentials produced by Petromyson, Kleerekoper & Sibakin; cardiac electric activity of Sicyases, Concha & Vargas; measurement of electrical activity of cardiac tissues, Gargouil & Coraboeuf; galvanotaxis in fishes, Spiecker.

Reactions to chemicals.—Effect of insecticide on young Cyprinus, Yoshihara & Fujioka; factors affecting toxicity of zinc to Lepomis, Cairns & Scheier; effect of agricultural chemicals on Misgurnus, Kubota & others; resistance of skin to natural and artificial attack, Acolat & Muller; toxicity of un-ionized ammonia, Merkens & Downing.

Effect of X-rays etc.—Survival of Carassius at low temperatures after irradiation, Gros & Bloch (1) & (2); effect of radioactive chemicals on Cyprinus, Kirpichnikov & others; radioactivity in Katsuwonus, Saiki & others; damage by X-rays to gonads of Lebistes, Kessler; effect of radiation on behaviour of egg cells, Neifakh (2).

Tropisms.—Photo-orthokinesis in larval Salmonids, Woodhead; reaction to light in migrating salmon and factors influencing it, general survey, Fields; reactions to light, Privolnev.

BIOCHEMISTRY

Chemical composition of some Philippine fishes, **Sulet & others**; protein content of some fish muscle, **Alexander**; amino acids of pyloric caeca of *Katsu*-

wonus, Kakimoto & others; nucleic acids of Latimeria, Vialli; synthesis of vitamin A by Gambusia, Grangaud & others; Vitamin A content of fish flesh, Junker; B-vitamins in pyloric caeca of Katsuwonus, Kakimoto; presence of thiaminase in freshwater fish organs, Suomalainen & Pihlgren; effect of raised temperature on tissue chemistry of Myoxocephalus, Musacchia & Clark; fat content of Pleuronectes flesus, Meyer-Waarden & Rübenberg; distribution of fat in some flatfish, Morawa; seasonal variation of fat content in Salmo letnica, Sapkarev (2); acetylcholine cycle in electric organs, Kuwabara; lipids of oocytes of Acipenser, Ginzburg; liver and muscle lipids in Arctic fishes, Musacchia & others; changes occurring in distribution of lipids in Esox testis, Lofts & Marshall; enzymes of hatching glands of Vimba, Baburina & Buznikov; skin carotenoids of Crenilabrus, Abolins & Abolins-Krogis; electrophoretic study of fish albumin, Drilhon & Fine; electrophoresis of shark serum, Starr & Fosberg; constituents of plasma of Scyliorhinus, Brull; constituents of blood serum of Anguilla, Drilhon & others; changes in blood levels of lactic acid in Salvelinus, Black (2); presence of myosin and actin in electric organ of Torpedo, Citterio & others; Maldacea & others; intestinal sugar absorption of Anguilla in fresh and salt water, Cordier & Maurice (1); intestinal absorption of sugars in Esox, Cordier & Maurice (2); pH of intestine of Scorpaena and its effect on absorption of glucose, Cordier & Worbe: suprarenal of Selachians, Turchini: proteolytic activity of thyroid in Scyliorhinus, Clements; histochemical study of developing Clupea egg, Yamamoto, T. S.; cytochemical study of fish eggs, Petrova; chemical differentiation of proteins during spermatogenesis, Alfert; constituents of oocytes, Yamamoto, K.

REPRODUCTION AND SEX

Age and size at sexual maturity.—Of Capoetobrama, Maxunov; length of Neothunnus, Yuen & June; "neoteny" (paedogenesis) in Lampetra, Zanandrea.

Differences between the sexes.— Sexual dimorphism in Rutilus pigus virgo, Misik.

Sex ratio—in Notropis, Reed (2).

Spawning season and place.—Spawning areas of Abramis in Vistula lagoon, Dziekonska; spawning places of Atlantic Clupea, Yudanov; of Sundsvall Bay Coregonus, Lindroth (1); egg site selection by Etheostoma, Winn; spawning season of Galeichthys, Ward; Hippoglossoides, Bagenal (1); of Istiophorus, Gehringer; of Neothunnus, Yuen & June; spawning grounds of Labeo in Katanga Province, Bont & Maes; breeding place of Rastrelliger, Kuthalingam; of Portugues Sardina, Cruz (1).

Factors influencing spawning.—Problems of winter spawning Acipenser, Barannikova; reproductive rate of Etheostoma controlled by temperature, Hubbs, C. & Strawn (1).

Breeding habits.—Pre-nuptial behaviour of Alosa, Medcof; breeding behaviour of Anableps, Trochanis; nest building of Arapaima, Migdalski; spawning habits of Dascyllus, Garnaud (2); reproductive behaviour of Gambusia, Warburton & others; breeding colouration of female Gasterosteus, Drapkin; breeding habits of Macropodus, Pinter; spawning behaviour of Notropis, Reed (1); of Petersius in aquarium, Poersch; spawning behaviour of Rhodeus, Wiepkema; of Wallagonia, Mazumdar.

Reproductive cycle. — In annual fishes, Myers; in Ditrema, Ishii; in Jenynsia, Turner; in & and & Sebastes marinus, Magnusson (1); spawning periodicity in Acipenser, Roussow; sexual photoperiodicity in Notropis, Harrington; factors influencing sexual cycle in Perca, Morozova, T. E.

Oogenesis and spermatogenesis.—Size of ovarian eggs in *Hemichromis*, **Peters**; spermatogenesis in *Muraenesox*, **Nishikawa**; size of nuclei during development of oocytes, **Sadov** (2).

Reproduction. — Of Acanthorhinus, Bjerkan; of Cottus bubalis, Thomopoulos & Bauchot; ecological adaptations in Comephoridae, Koryakov; function of testis in various types of spawning, Butzkaya.

Fertilization.—Results of insemination of Carassius eggs with sperm from other genera, Suzuki (2); structure and activity of spermatozoa in Clupea, Yanagimachi (2); manner of sperm entry into egg in Clupea, Yanagimachi (3); cytological and cytochemical investigations on fertilization in Lampetra, Chubareva; results of cross fertilization with different genera, Suzuki (1); change in X-ray sensitivity during fertilization, Neifakh (1).

Parthenogenesis.—In Sakhalin Clupea, Krizhanovski (2).

Hermaphroditism.—Sex reversal in teleosts, Ancona (1); in Coris, Reinboth.

Hormones and sex.—Effect of hormone injection on gonads of Anguilla, Ancona (5); induced spawning in Clarias, Ramaswami & Sundararaj (2); induced spawning in Heteropneustes, Ramaswami & Sundararaj (1); effect of hormone on vas deferens of male Monacanthus, Ishii & Egami; inhibition of male characteristics in female Oryzias, Egami; effect of steroid hormones on Salmo alevins, Ashby (2); effect of pituitary extract on ovary, Kirshenblat; effect of hormones on sexual differentiation, Vallowe.

Number of eggs produced. — By Ameiurus, Sedlár; by Arapaima, Migdalski; by Dascyllus, Garnaud (2); by Hippoglossoides, Bagenal (1); by Hirundichthys, Hall; annual variations in fecundity, Bagenal (2).

Parental care.—Oral gestation in Galeichthys, Ward; oral incubation in Opisthognathus, Böhlke & Chaplin; parental care in Nannacara, Kuenzer.

EVOLUTION, VARIATION AND GENETICS

Evolution. — Of actinopterygians, Saint-Seine; of Carangidae, Aleev (7); Of Dipnoi, Lehman (1); of vertebrate teeth, Buistrov; of pelagic eggs, Orton.

Hybrids.—Survival rate of Etheostominae hybrids, Hubbs & Strawn (3); \mathbf{F}_1 hybrids of Lepomis gibbosus \times L. macrochirus, Lagler & Steinmetz; Rutilus rutilus \times Chalcalburnus chalcoides, Nalbandoglu; evidence of introgressive hybridization in Salmo, Miller, R. B. (2); intraspecific hybrids and hybrid vigour, Donaldson, Hansler & Buckridge; sperm viability as a factor in hybridization, Hubbs, C. (3); list of natural hybrids, Slastenenko.

Genetics.—Influence of environment on sex in Anguilla, Ancona (4); hereditary differences in populations of Xiphophorus, Gordon, M.; genetical colour-varieties in aquarium fishes, Ladiges. Chromosome studies.—In Eptatretus, Nogusa.

Raciation and speciation.—Speciation in Coregonids, Svärdson (1); Intraspecific relationships in fishes, Klyuchareva; evidence of phenotypic or genotypic effects of interference by man on fish populations, Miller, R. B. (2).

Albinism, etc.—Gene for albinism present in Anoptichthys, Sadoğlu.

Biometric studies.—On Merluccius, Franca (1); on Rutilus pigus virgo, Misik; on Sardina and Sardinella in Castiglione Bay, Dieuzeide & Roland (3); on Angolan Sardinella, Monteiro; on Thunnus thynnus, Genovese.

Phylogeny.—Origin of Vertebrates, Berrill; Lehman (3); axial rotation and origin of vertebrates, Stanley-Jones; first tetrapods and Crossopterygii, Jarvik; phylogeny of the Agnatha, Balabai (1).

ECOLOGY AND HABITS

General bionomics.—Of Blennius pholis, Qasim; of Bregmaceros, Clancey; of Chaenobryttus gulosus, Larimore; of Cottus gobio, Smyly; of Lagodon, Caldwell; of Lepomis in an Iowa lake, Mayhew (1); of Luciogobius, Dôtu (3); of Psenes, Legaspi; of Salmo and Salvelinus in Murmansk region, Vladimirskaya (2); of Sardinops ocellata, Davies (3); of Schizothorax argentatus in Lake Balkhash, Savina; of Squalius, Hoffmann; treatise on marine ecology, Hedgpeth.

Age and growth.—Age estimation of Anguilla, Rahn; maximum size and weight of Arapaima, Migdalski; growth of Aspius, Troitzkii; Length/weight relationship in Astroconger, Okada & Suzuki; growth of Carpiodes, Buchholz; condition factor of Chrysophrys, Cassie; age composition of Manx Clupea, Bowers (1), (2), (3); age and size of Manx Clupea, Smith, W. C.; growth of Cyprinus, Menzel & Steffens; growth rhythm of young Cyprinus, Caraiman; growth of Engraulis post larvae, using vertebral counts, Hayashi & Suzuki; age and growth of Irish Esox, Austin; growth rates of Esox and Scardinius, Healy (2); size related to feeding of young Gadus, Mironova (1); growth rate and size groups of Hilsa, Sarkar; growth of Ictalurus, Harrison; of Istiophorus

americanus, de Sylva; of Lampanyctus, Bolin; size differences in Merluccius, Matta; growth of Micropogon, Haven; of a 'stunted' population of Micropterus, Suttkus (1); study on Adriatic Mugilidae, Moravic; age and growth of Mullus barbatus in Israel waters, Gottlieb; growth rate of Notropis, Reed (2); growth of Perca, Lucioperca and Acerina in north German waters, Tesch; growth of exoskeleton in Pteraspis dunensis, Fahlbusch.; Growth of Salmo trutta in Lake Rensjön, Runnström; of Loch Kernsary Salmo, Swan (1); some large Irish Salmo salar, 'Avondhu'; growth of Irish Salmo, Healy (4), Twomey, Went (2), Went (6), Went & Vickers; growth of Salmo before and after raising of lake level, Frost (1); growth of Salmonid fry at differing temperatures, Mekhanik; of Portuguese Sardina, Cruz (1); of Sardina in Jugoslav waters, Muzinic; of Stizostedion, Rose: length/weight relationship in Xiphias, Postel (1); relation between growth-rate and density of population, Oliva & Frank; influence of size on growth-rate, Larkin, Terpenning Parker; relation between food supply and condition coefficient, Ellis Gowing; principles of fish growth, Aleev (3); Lee's phenomenon, Kuznetzov.

Fish marking experiments.—Recaptures of tagged Alosa, Vladykov (1); experiments on Clupea, Sahrhage; results of marking Gadus in Faroes, Joensen; of Faxa Bay Gadus, Jónsson; tagging study on Micropterus, Kimsey (2); tagged Microstomus recaptured after six years, Best (2); results of Oncorhynchus tagging, Milne; results of Salmo tagging in Labrador, Blair (1) & (2); experiments on fishes in Arctic Alaska, Wohlschlag.

Scale reading etc.—Of north German Percidae, Tesch; validity of scale-reading in *Cirrhina*, Jhingram.

Population studies.—Abramis brama of the Vistula, Poczopko (2); Clupea of southern North Sea, Cushing & Burd; of Atlantic Clupea, Biryukov; Leim & others; Coregonus of Sundsvall Bay, Lindroth (1); estimation of Lepomis population in Iowa, Lawrence; German Bight Pleuronectes, Kotthaus; Oncorhynchus keta in Amur basin, Birman (2); comparison of three races of Oncorhynchus nerka, Vernon; North Sea

Thunnus, Tiews; Thymallus in Murmansk lakes, Vladimirskaya (3); populations of Iowa ponds, Carlander & Moorman.

Abundance.—Estimate of number of Sardina pilchardus in Channel, Cushing (2).

Fluctuations of stock.—Swedish Esox and Perca, Alm; Oncorhynchus in Kamchatka waters, Krogius & Krokhin; self regulation of Osmerus stock in Zhizhitzk Lake, Abrosov & Agapov; fluctuations of Parophrys vetulus stock, Ketchen; causes of fluctuation in Baltic Salmo, Lindroth (2); Svärdson (2); Portuguese Sardina, Cruz (2); fluctuations due to environmental changes, Moiseev (1).

Ecological studies.—In Polish waters, Pliszka; some Hungarian waters, Jaszfalusi; Carpathian rivers, Starmach; Borcea-de-Jos lakes complex, Danube, Popescu-Gorj & others; fishes of a German reservoir, Miegel (1); fishes of Niger Delta, Daget; fisheries study of a Florida lake, Moody; brackish water systems in Mie Prefecture, Okada & Kubota (1); adjustment in a disturbed estuary, Reid (2); Palaeoecology of Agnatha (bibliography), Robertson, G. M. (1) & (2); Palaeoecology of Haskell Limestone (with palaeonisids), Miller & Swineford; Palaeoecology of fishes, bibliography, David.

Ecological Surveys.—Of Baffin and Alazan Bays, Texas, Breuer; Upper Laguna Madre, Texas, Simmons; trawling survey off Algerian coast, Dieuzeide & Roland (1); off the Gold Coast, Salzen.

Relations with physical environment. —Of Blennius, Lecal; temperature selection by Cyprinus, Pitt & others; effects of storms on shallow water fauna, Robins (1); effect on growth and breeding of repeated variations in river level, Moskalenko; survival of fish in water at low temperatures, Scholander; size of marine fishes invading fresh water, Gunter; poisoning from red tide and its causes, Anon. (2).

Special Habitats.—Emergence of fishes from Californian subterranean waters, Herald; blind fishes, Lüling; annual fishes, Myers.

Introduction and acclimatization.— Introduction of Abramis into Khristoforov reservoir, Melnikov; Ophiocephalus in Japan, Okada; Carassius into Veselovsk reservoir, Shchetinina; results of acclimatising Coregonus in Roumania, Busnita & others (1).

Aestivation and hibernation.—Drought survival of eggs and conditions of life in Nothobranchius, Hansen.

Food and feeding habits.—Feeding habits of Acanthogobius, Therapon, Tribolodon and Oryzias, Imamura & Hashitani; food of Anguilla in Baltic, Murina; Ameiurus, Frank; Aplodinotis, Moen; Arapaima, Migdalski; Aspius, Vasiliu & Nicolau (2); influence of light on feeding of Blicca, Dudek; food of Coryphaena, Williams & Newell: young Esox, Makkoveeva; young Istiophorus, Gehringer; Mugilidae, Ebeling; Notropis, Reed (2); Oncorhynchus masu, Semko; O. tshawytscha, Merkel; seizing of food from ground by Osteoglossum, Heck; food of Parathunnus and Neothunnus, King & Ikehara; Perca in Volga delta, Ivanova, M. N.; feeding habits of Percarina, Kanaeva; Pleuronectes larvae, Shelbourne; Rastrelliger, Kuthalingam; Caspian Rutilus, Novi-kova; Salmo trutta, Nilsson; food of Sander, Vasiliu & Nicolau (1); Sardinops ocellata, Davies (2); Scyliorhinus feeding on the nudibranch, Tritonia, Stephen & others; food of Sebastes, Magnusson (3); Trachurus, Pora & others; feeding habits of Vimba, Vasiliu & Costea; of some pelagic fishes, Tokarev; of fishes from an Iowa lake, Kutkuhn; of Veselovsk reservoir fishes, Kruglova; of Kuyu-Mazar reservoir fishes, Abdullaev; food of Gruzian fishes, Kokhiya (1) & (2); role of plankton in nutrition of oriental pond-fishes, Vaas; feeding rhythm of fry, Olifan; use of sense organs in feeding, Andriyashev; adaptations to feeding on large, armed prey, Fedorov.

Migrations and movements.—Of Clupea, Biryukov; Coryphaena off E. Africa, Williams & Newell; Bear Island Gadus, Trout; Oncorhynchus, Birman (3); Oncorhynchus in western North Pacific, Taguchi; diurnal activity of Perca flavescens, Brown & Rosen; migrations of Salmo, Azbelev & Lagunov; L. Rensjön Salmo trutta, Runnström; movements of Salmo off N.E. coast of England, Hartley & Brayshaw; Irish

Salmo salar, Toner; Went (1); Went & Vickers: Salmo in home territory, Miller, R. B. (1); factors determining age and season of seaward migration of smolts of Salmo, Went (5); Seriola along Pacific coast of Japan, Mitani; Trachurus off the Roumanian coast of Black Sea, Nicolae & Cautis; daily rhythmic behaviour of some marine fishes, Hempel; migratory instinct in fishes, Kozlovski; movements of spawning population of Eopsetta, Alverson, D. L. & Chatwin; spawning migrations of Oncorhynchus gorbuscha affecting size, Birman (1); tenacity of age of spawning migration in a population or fraction of a population through several generations in Salmo, Went (5).

Behaviour.—Of Clupeonella in captivity, Nikonorov; Electrophorus, in captivity, Boursot; Black Sea fish in winter, Aslanova; study of fish behaviour, Zusser.

Social behaviour.—Aggregational behaviour of Cottus, Greenbank; method of fighting by Nannostomus, Wickler (3); biological significance of fish shoaling, Nikolski (1); schooling behaviour of some fishes, Springer, S. (2).

Relations with other organisms.— Effect of predators on size of population, Fortunatova (2).

Symbiosis and other associations.— Cleaning of one species by another; association for concealment of predator with harmless fish (repetition of 1955 report), Eibl-Eibesfeldt.

Methods of offence and defence.— Avoidance of Scorpaenichthys eggs by predators, Pillsbury.

Diseases and parasites.—Cauliflower disease of Anguilla, Lühmann & Mann; diseases of hatchery Salmo fed on artificial food, Faktorovich (2); a bacterial infection of Cichlids in captivity, Nigrelli & others; diseases caused by bacteria, Rushton (1) & (2); protozoan parasites of Gadus, Callionymus and Microstomus, Noble; parasites of a population of Phoxinus, Dalenius; Sardina parasitized by Peroderma, Dieuzeide & Roland (4); parasitic infestation of gonad of Sardina, Santos Pinto (1); Clinostomium parasitising Schilbeodes, Schwartz; presence of the fungus Ichthyosporidium in Serranus and Polyprion, Reichenbach-Klinke;

parasites of Xiphias, Postel (1); of some freshwater fishes of Roumania, Radulescu & Vasiliu-Suceveanu.

Mortality.—Causes of mass mortality in Oka river, Kozlov; possible cause of mass mortality, Bonte.

DISTRIBUTION

a. GENERAL

Habitat of early vertebrates, Robertson, J. D.; Atlantic origin of some fish groups, Svetovidov (3); diagram of distribution of fish eggs, Zaitzev.

b. MARINE

General. — Effect of temperature changes on distribution in E. Mediterranean, Oren; Zooplancton (including fishes) of western Mediterranean, Alvarino (3); Zooplancton of Spanish waters, Alvarino (4); indicator species and fish eggs and larvae from Newfoundland waters, Alvarino (1) & (2); Zoogeography of Labrador fishes, Backus (1); Zoogeographic study on Japanese Soleidae, Ochiai; vertical distribution of post-larval fishes of Florida current, Bsharah.

Arctic.—Alaska, Mohr, Wilimovsky & Dawson; Barents Sea, Barsukov (3).

North temperate and subarctic.-Aleutians, Welander, Alverson & Bergman; Behring Sea, Mednikov & Prokhorov; Labrador, Backus (1), Gordon & Backus; Iceland and Greenland, Kotthaus & Krefft; Sweden, Curry-Lindahl; Julin; Scotland, Rae & Wilson; Bristol Channel, Bassindale & Barrett; British coast, Wheeler, A. C.; Gironde, France, Bauchot & Lubet; Portugal, Albuquerque; Corsica, Houdemer (1) & (2); Algerian waters, Dieuzeide & Roland (2); Mediterranean, Padoa & others; Madeira, Maul; North Atlantic, Nybelin; Western North Atlantic, Caldwell & Briggs; Bermuda, Schultz (2); British Columbia, Welander, Alverson & Bergman; Gulf of Maine, Scattergood & Coffin; New England and Massachusetts, Mather & Gibbs; Long Island Sound, Morrow (1);Alabama coast, Boschung; Chesapeake Bay, Massmann; Monterey Bay, California, Fast; Southern California, Norris; Texas, Kemp; Leary (1) & (2); Florida, Bigelow & Schroeder; Phillips; Robins (1); Wood, F. G.; North Pacific,

Welander & others; China, Fowler (1); Palau Isl., Aoyagi; Japan, Abe (1); Dôtu (1); Kamohara (2); Kuroda (2); Kuroda (3); Kuronuma & Yamaguchi; Maruyama; Oshima (1); Tomiyama & Abe; Yoshida & Ito; Riu-Kiu Islands, Kamohara (3).

Tropical.—Red Sea, Tortonese; Senegal and Gold Coast, Briggs; East Africa, Smith, J. L. B. (1); Williams; South and East Africa, Smith, J. L. B. (13); Mozambique, Smith, J. L. B. (3), (5) & (12); Aldabra, Smith, J. L. B. (2), (3) & (4); Seychelles, Smith, J. L. B. (4); W. Indian Ocean, Smith, J. L. B. (9), (10) & (12); Eastern Pacific, Clemens; Pacific, Otsu & Yoshida; North Borneo, Inger: Hawaii, Schultz (2); Gulf of Mexico, Briggs & Caldwell; Grey; Kanazawa; Mead (2); Schultz (2); Springer, S. & Bullis; Springer, S. & Thompson; Woods; West Indies, Walters (2); Bahamas, Böhlke (1), (2), (3) & (4); Krumholz (1); Cuba, Mead (1); Puerto Rico, Erdman; Smith, C. L.; Trindade Isl., Brazil, Pinto (1).

South Temperate.—South Africa, Jubb; Smith, J. L. B. (6), (11), (14), (15) & (16); Chile, Buen (3); Schultz (1); Queensland, Haysom; Munro (2); N.S. Wales, Schultz (2); Australia, Fowler (2); Marshall, T. C.; Munro (1); Scott, T. D.; Whitley (2); New Zealand, Garrick; Parrott.

Antarctic.—Abe (3).

c. Freshwater

General.—Biogeography of African lakes, Blanc; hydrobiology of L. Tanganyika, Poll (2); effect of water diversions on distribution of British Columbia fishes, Lindsey; distributional patterns of Texas fishes, Hubbs, C. (1); pleistocene glaciation and distribution of Minnesota Cyprinoids, Underhill.

Palaearctic. — Finland, Pynnönen; Shetland, Swan (2); Yorkshire, Hazelwood; Camargue, France, Lévèque; Holland, Bemmel; Austria, Bauer & Schubert; Rumania, Balon (2); Nalbant (1) & (2); Czechoslovakia, Oliva (1); Liguria, Italy, Arbocco; Portugal, Albuquerque; Iran, Kozhin; Israel, Steinitz & Ben-Tuvia; Pechora river, Russia, Kurilov; Sari-Su basin, Russia, Ereshchenko; Gruzia, Demetrashvih; Elanidge (1) & (2); East Asia, Boeseman

(1); North China, Mori, T.; Japan, Hamada; Okada & Kubota (2); Oshima (2) & (3).

Oriental.—Ceylon, Meinken (2); Johore and Java, Raut; Thailand, Benl (1) & (2); Klausewitz (8), (9) & (10); Sunda Sea, Thienemann.

Ethiopian.—French Guinea, Lambert; Belgian Congo, Boeseman (2); Poll (1); Uganda, Greenwood (4); Lake Victoria, Greenwood (1); Lake Nyasa, Fryer; East Africa, Klausewitz (11).

Nearctic.—Labrador, Backus (1); British Columbia, Lindsey; Saskatchewan, Scott, W. B.; Washington, Fitzgerald; Dakota, Robins & Raney (1); Iowa, Carlander; Cleary; Texas, Hubbs, C. (2) & (4); Hubbs & Brown; Hubbs & Springer; Robins & Raney (2); Louisiana, Suttkus (2); Lower Mississippi drainage, Hemphill; Alabama, Cooper & Hemphill; Florida, Kilby & Caldwell.

Central America.—Mexico, Hubbs & Springer; Miller, R. R. (2); New Mexico, Koster.

South America.—British Guiana, Meinken (1); Surinam, Hoedeman (4); Brazil, Hoedeman (2); Ribeiro (1) & (2); Argentina, Aramburu (3).

Australian region and Polynesia.—Good; Whitley (1).

d. Fossil

General.—Origin of Vertebrates, Berrill; Lehman (3); origin of first tetrapods, Jarvik; habitat of early vertebrates, Robertson, J. D.; palaeoecology of Agnatha, Robertson, G. M. (1); bibliography of palaeoecology of Agnatha, Robertson, G. M. (2); bibliography of palaeoecology of fishes, David; evolution of Dipnoi, Lehman (1); palaeogeographic importance of Dipnoi, Pidoplichko Kurazhskovski; evolution actinopterygians, Saint-Seine; evolution of teeth, Buistrov; fish in carbonate concretions, Weeks; guide fossils (Chordata) of USSR Devonian, Obruchev (1); paired pineal organs, Edinger; determination, Chavan & Cailleux; fossilization, Breder.

Palaeozoic.—Ordovician, Ross; Devonian, Tarlo; Balabai (2); Balabai & Opalatenko; Bystrow; Chi; Fahlbusch; Gross; Kulczycki; Obruchev (1); Orvig (1), (2) & (3); Smith, I. C.; Carboni-

ferous, Baird; Dorr; Heide; Khabakov (1) & (2); Miller, H. W.; Miller & Swineford; Peyer; Permian, Hoehne; Konta.

Mesozoic.—Trias, Beltan; Triassic, Guffroy; Lehman (2); Nielsen; Rutte; Jurassic, Khabakov (2); Lapparent; Liu & Yeh; Orvig (2); Peyer; Weiler (3); Cretaceous, Avnimelech; Ghlikman (1) & (2); Liu, T. S.; Santos; Siegfried; Springer, V. G.

Tertiary.—Eocene, Stinton (1); Oligocene, Danil'chenko; Ghorbach; Smigrelska; Walters (1); Weiler (1) & (2); Miocene, Henriques da Silva; Kruckow; Kudrin; Liu (2); Stinton (2); Svichenskaya; Vokes; Pliocene, Liu (1); Papp & Thenius; Sarma; Yakovlev.

Quaternary.—Munthe; Pleistocene, Huang; Holocene, Greenwood (2) & (3).

III.—SYSTEMATIC INDEX

GENERAL

Classification.—Evolutionary aspects of, Marshall, N. B. (1); Review of Berg's classification, Obruchev (2); utilization of X-rays, Miller, R. R. (3).

Nomenclature.—Status of Bagre Cuvier (Oken), Bagrus Val. and Porcus Geoffroy St. Hilaire, Jayaram; addition of Cubiceps Lowe 1843 to the Official List of Generic Names, Opinion 461; designation of type species for Heterandria, Opinion 375; Neoceratodus valid in Dipnoi, Miller, R. R. (1); validation of Osmerus, Hemming; Direction 69; additions to Official List of Generic Names, Directions 33-34; correction of entries in Official List of Generic Names, Direction 56; additions to the Official List of Specific Names, Direction 57; determination of gender, Direction 30; Opinions 438, 485 & 487.

Faunistic works.—Fishes of Portuguese waters, with keys, Albuquerque; keys to German freshwater fishes, Döderlein; freshwater fishes of New Mexico, Koster; fishes of Uganda, Greenwood (4); Freshwater fishes of Australia, Whitley (1); New Zealand marine fishes, Parrott; fishes of China, with keys, Fowler (1); Keys to economic fishes of U.S.S.R., Borisov & Ovsyannikov; Ostracoderms and fishes of the Devonian in USSR, Obruchev (1).

MARSIPOBRANCHII

Phylogeny, Balabai (1).

Bibliography of palaeoecology, Robertson, G. M. (1) & (2).

†EUPHANERIDA

Jamoytius and origin of vertebrates, Berrill.

†HETEROSTRACI

New psammosteids from Emsian of C. Poland, Tarlo.

Astraspis desiderata and Eriptychius americanus figd. from Ordovician E. Montana, Ross.

Ganosteus stellatus figd., Obruchev (1).

Psammolepis paradoxa figd., Obruchev (1).

Psammosteus marginalis, P. maeandrinus figd., Obruchev (1).

Pteraspis dunensis deser. and figd., Fahlbusch; P. lerichei, P. major, P. elongata figd. and range in Podolian Old Red Sandstone discussed, Balabai (2).

Pycnosteus palaeformis, P. tuberculatus figd., Obruchev (1).

Rhinopteraspis Jackel 1919 validated, type Archaeoteuthis dunensis Roemer 1855, and Palaeoteuthis Roemer 1855 and Archaeoteuthis Roemer 1855 suppressed, Opinion 438.

Schizosteus striatus figd., Obruchev (1).

†THELODONTI

Scales figd. from West Spitsbergen, ørvig (3).

†OSTEOSTRACI

Alaspis gen. nov. type A. macrotuberculata sp. nov. U. Devonian Escuminac Bay, Canada, ørvig (1) p. 372 fig. 1D.

Benneviaspis holtedahli, endoskeletal hard tissues figd., Ørvig (3).

Escuminaspis gen. nov. type Cephalaspis laticeps Traquair, ørvig (1) p. 372 fig. 1C.

†ANASPIDA

Pharyngolepis kiaeri sp. nov. Downtonian Norway p. 380 text-figs. 2-4, 8, 10 pls. iv, v; P. oblongus figd., Smith, I. C.

CYCLOSTOMATA

PETROMYZONIDAE

Of Australia, Whitley (1).

Distribution of Eurasian forms, Lanzing.

List of natural hybrids, Slastenenko.

Development of intestinal blood system and vascular couples, Baxter.

Lampetra experiments on neural crest, Newth; Mauthner neurones of larvae, Whiting; L. aepyptera recorded from Alabama, Cooper & Hemphill; L. fluviatilis of the Baltic, Abakumov; structure of gills, Morris, R.; L. japonica bionomics and systematics, Morozova, T. A.; ecological notes, Nikolski (2); L. japonica kessleri notes on spawning, Semenov-Tyan-Shanski; L. planeri osmotic conditions in egg and early stages, Hardisty; origin of acousticofacialis complex, Fisk; L. zanandreai, sexual maturity in larvae, Zanandrea,

Petromyzon marinus laboratory care and feeding of larvae, Sawyer; 'spike' potentials produced in head region, Kleerekoper & Sibakin.

MYXINIDAE

Eptatretus burgeri, E. okinoseanus chromosome study, Nogusa.

Myxine glutinosa occurrence in Mediterranean, Dieuzeide.

†PLACODERMI

ACANTHODII

Teeth and spines of Gomphodus and Nostolepis descr. and figd., Gross.

Acanthodes ef. bronni scales figd., Hoehne; A. lopatini figd., Khabakov (1).

Atopacanthus jaws descr. and figd., ørvig (3).

Onchus marginalis, O. concinnus figd., Obruchev (1).

ARTHRODIRA

Microstructure of dermal bone, Bystrow.

Paired pineal organs, Edinger.

Actinolepis? sp. skull-roof figd., Ørvig (3); A. tuberculata, Plourdosteus livonicus, Phlyctaenaspis acadica microstructure of tubercles, Gross.

Anomalichthys ingens figd. from U. Devonian Poland, Kulczycki.

Chelyophorus pskovensis sp. nov. Devonian Russia p. 201 pl. liii fig. 4; C. verneuili figd., Obruchev (1).

Coccosteus livonicus, minor, mironovi, trautscholdi figd., Obruchev (1).

Deveonema gen. nov. type D. obrucevi sp. nov. U. Devonian Poland, Kulczycki p. 331.

Dinichthys denisoni p. 313 text-fig. 10 pl. vi fig. 4; vii; D. ceterus p. 314 pl. viii fig. 1, spp. nov. U. Devonian Poland; D. pustulosus, D. cf. tuberculatus figd., Kulczycki.

Dunkleosteus endoskeletal tissues descr. and figd., Ørvig (3).

Ellopetalichthys gen. nov. type Macropetalichthys scheii Kiaer Ørvig (3) p. 300 text-figs. 5F, 7A pl. i.

Heterostius sp. figd., Obruchev (1).

Holonema radiatum figd., Obruchev (1); figd. from U. Devonian Poland, Kulczycki.

Homostius sp. figd., Obruchev (1).

Luetkeichthys gen. nov. type L. borealis sp. nov. M. Devonian North Land USSR, Bystrow p. 250 figs. 9-14.

Lunaspis sp. figd. from U.S.S.R., Obruchev (1).

Malerosteus gen. nov. type M. gorizdroae sp. nov. U. Devonian Poland, Kulczycki p. 291 text-figs. 3-6, pl. i, figs. 4-7; ii; iii figs. 1-3.

Operchallosteus gen. nov. type O. vialowi sp. nov. U. Devonian Poland, Kulczycki p. 335 pl. xii fig. 3.

Pachyosteus bulla figd. from U. Devonian Poland, Kulczycki.

Palaeomylus and Ptyctodus microstructure of tooth-plates descr. and figd., Ørvig (3).

Phyllolepis sp. figd., Obruchev (1).

Ptyctodus sp. figd., Obruchev (1).

Rhynchodus and Ptyctodus microstructure of jaws, Gross.

Sedowichthys gen. nov. type S. terraeboreae sp. nov. M. Devonian North Land USSR, Bystrow p. 240 figs. 1-8.

Titanichthys kozlowskii sp. nov. U. Devonian Poland, Kulczycki. Tollichthys gen. nov. type T. polaris sp. nov. M. Devonian North Land U.S.S.R., Bystrow p. 257 figs. 15-19.

Tomaiosteus gen. nov. type T. grossi sp. nov. U. Devonian Poland, Kulczycki p. 298 text-fig. 7 pl. iii figs. 4, 5.

ANTIARCHA

Asterolepis ornata, radiata figd., Obruchev (1).

Bothriolepis-beds in Yunnan, age, Chi; B. ornata, maxima, cellulosa panderi figd., Obruchev (1).

SELACHII

Presence of a paraphysis, Kappers.

? Egg capsules, *Palaeoxyris tamica* sp. nov. Carboniferous Donetz Basin p. 86 pl. ix fig. 8; *Crookallia czernyshevi* figd., Jurassic Crimea, Khabakov (2).

†ICHTHYODORULITES

Alienacanthus gen. nov. type A, malkowskii sp. nov. U. Devonian Poland, Kulczycki p. 357 pl. xiii fig. 1.

Eczematolepis, Gamphocanthus, Oëstophorus systematic position discussed, Ørvig (3).

Sentacanthus gen. nov. type S. zelichowskae sp. nov. U. Devonian Poland, Kulczycki p. 356 pl. xiii fig. 3.

†ICHTHYOTOMI

Pleuracanthus cf. arcuatus figd. from Pennsylvanian Michigan, **Dorr**.

EUSELACHII

Inc. sed.

†Dicrenodus okensis teeth figd., Khabakov (1).

Significance of fin-spines, Peyer.

Morphology of hypophysis, Honma (2).

†CTENACANTHIDAE

Ctenacanthus varians var. russakovi spine figd., Khabakov (1).

Nemacanthus monilifer, Ctenacanthus heterogyrus, C. brevis fin-spines figured, **Peyer**.

PLEUROTREMATA

HEXANCHIDAE

Hexanchus griseus from western North Atlantic; stomach contents, Backus (2).

ODONTASPIDAE

Odontaspis †cuspidata teeth figd., Miocene Maryland, Vokes.

†Pseudoisurus gen. nov. type P. tomošus sp. nov. U. Cretaceous? Russia, Ghlikman (2) p. 116 pl. i figs. 16-20.

LAMNIDAE

Relationship to Odontaspidae, Ghlikman (2).

Carcharodon carcharias juvenile female taken in Cape Cod Bay, **Backus** (2).

Cetorhinus maximus southern extension of range, Wood, F. G.; notes on juvenile female, Backus (2).

Isurus key to species, Smith, J. L. B. (11); I. †denticulatus sp. nov. Cenomanian, U.S.S.R., Ghlikman (1) p. 103 figs. 1-17; I. oxyrinchus measurements of an adult, Krumholz (2); I. tigris africanus subsp. nov. South Africa, Smith, J. L. B. (11) p. 96.

†Otodus obliquus teeth figd. Miocene Maryland, Vokes.

Oxyrhina †desori tooth figd. Miocene Maryland, Vokes.

†Paraisurus gen. nov., type Oxyrhina macrorhiza Pietet and Campiche, Ghlikman (2) p. 116 pl. i figs. 14, 15.

†ORTHACODIDAE fam. nov.

Euchlaodus gen. nov. type Oxyrhina lundgreni Davis, Ghlikman (2) p. 116 pl. 1 f. 16.

Paraorthacodus gen. nov., type Synechodus recurvus Traut., Ghlikman (2) p. 115 pl. i figs. 6-13.

RHINCODONTIDAE

Rhincodon typus external morphology of embryo, Reid (1).

SCYLIORHINIDAE

Scyliorhinus caniculus feeding on the nudibranch, Tritonia, Stephen, Rae & Wilson; S. stellaris comparison of lampbrush chromosomes with those in Sepia and Anilocra, Callan.

CARCHARHINIDAE

Aprionodon isodon from Puerto Rico, Erdman.

Carcharhinus acronotus from Puerto Rico, Erdman; C. albimarginatus recorded from Mozambique, Smith, J. L. B. (5).

Eulamia falciformis stomach contents and colour notes, Backus (2).

Galeocerdo †aduncus tooth figd. Miocene Maryland, Vokes.

Hemipristis †serra figd. from Orissa, Sarma; teeth figd. Miocene Maryland, Vokes.

Hypoprion signatus recorded from Bahamas, Krumholz (1); from Gulf of Mexico, Springer, S. & Thompson; from New England, Mather & Gibbs.

Mustelus, included with Scylliogaleus as separate family, key to S. African spp. p. 357, M. palumbes sp. nov., S. Africa, pp. 357 and 358 text-fig. 1, E and F, Smith, J. L. B. (16); M. canis from Puerto Rico, Erdman.

Neotriakis sinuans gen. et sp. nov. S. Africa, Smith, J. L. B. (15) p. 262 figd.

Prionace glauca data on embryos, Backus (2).

Pterolamiops longimanus from Puerto Rico, Erdman.

Scoliodon terrae-novae taken in deep water, Backus (2).

Scylliogaleus made type genus of separate family with Mustelus, p. 353 S. quecketti descr. and figd., Smith, J. L. B. (16).

†Edestidae

Physonemus cf. acinaciformis figd. from Pennsylvanian of W. Virginia, Baird.

†HYBODONTIDAE

Acrodus anningiae fin-spines figured, Peyer.

SQUALIDAE

Taxonomic review, with keys to genera and species, Bigelow & Schroeder.

Acanthorhinus carcharias description of uterus and foetus, Koefoed; reproduction, Bjerkan.

Etmopterus baxteri p. 172 figd., E. abernethyi p. 181 figd. spp. nov. New Zealand, Garrick; E. bullisi sp. nov. Florida, Bigelow & Schroeder p. 50 figd.

Euprotomicrus bispinatus female captured north of Midway Isl., Pacific, Otsu & Yoshida.

Squalus acanthias acanthias, S. a. suckleyi, a comparison, Lindberg & Legeza.

HYPOTREMATA

RAJIDAE

Raja batis correlation between muscles and skull, Gottenbos; R. clavata, function of ampullae of Lorenzini, Murray; R. rhina an abnormal example, Barraclough; R. spinicauda extension of range and description of young stages, Kreftt.

DASYATIDAE

Dasyatis centroura from South Florida, Phillips; D. favus, D. sephen recorded from E. Africa, Smith, J. L. B. (6); D. lubricus sp. nov. Algoa Bay, Smith, J. L. B. (6) p. 429 figd.

MYLIOBATIDAE

Aetobatus narinari from Chesapeake Bay, Massmann.

Myliobatis †pachyodon tooth-plate figd. Miocene Maryland, Vokes.

†BRADYODONTI

Petalodus jewetti sp. nov. Pennsylvanian Kansas, Miller, H. W. p. 82 figs.

Psammodus rugosus teeth figd., Khabakov (1).

OSTEICHTHYES

Evolution of actinopterygians, Saint-Seine.

Sexual differentiation, Ancona (3).

PALAEOPTERYGII †ARCHISTIA

PALAEONISCIDAE

Birgeria sp. figd. from Triassic S. Madagascar, Guffroy.

Boreosomus merlei (sic) brain-cast figd., Beltan.

Elonichthys aitkeni scales figd. from L. Carboniferous Egypt, Heide.

Rhadinichthys laevis, R. canobiensis scales figd. from L. Carboniferous Egypt, Heide.

Scanilepis dubia, Orvikuina vardiaensis, Gyrolepis albertii histology of scales, Ørvig (2).

Sunolepis gen. nov. type S. yumenensis sp. nov. Cretaceous, Western Karsu, China, Liu, T. S. p. 108 2 text-figs. pls. i-vi.

BOBASATRANIIDAE

Ecrinesomus dixoni and Bobasatrania mahavavica figd. and redescr., Lehman (2).

Lambeichthys gen. nov. type Platysomus canadensis Lambe, Lehman (2) p. 16.

CHONDROSTEI

Otol. (inc. sed.) malmensis sp. nov. Jurassic W. Germany p. 229 pl. iii figs. 1, 2; Otol. (inc. sed.) altus, dorsolobatus, brandi, dorsoarcuatus figd., Weiler (3).

†ERROLICHTHYIDAE

 $\it Errolichthys\ mirabilis\ deser.\ and\ figd.,$ Nielsen.

ACIPENSERIDAE

List of natural hybrids, Slastenenko.

Development of perivitelline space in occytes, Sadov (1).

Acipenser results of artificial propagation, Manea & Mihai; breeding migrations in Roumanian Danube, Leonte; A. fulvescens spawning periodicity, Roussow; A. medirostris taken at Point Vicente, S. California, Norris; A. stellatus lipids in oocytes and eggs, Ginzburg; effect of different temperatures on hatching gland, Ignatyeva (2); A. stellatus, A. güldenstaedti deformation of pectoral fin, Huzum.

Huso huso hatching gland, Ignatyeva (1).

NEOPTERYGII

PROTOSPONDYLI

†SEMIONOTIDAE

Lepidotes sp. rec. Bathonian Morocco, Lapparent.

Semionotus bergeri figd., Rutte; S. capensis figd. from Basutoland, Guffroy.

AMIIDAE

Amia calva gonad differentiation, Ancona (2).

ISOSPONDYLI

Respiratory importance of pigment in eggs, Soin.

†LEPTOLEPIDAE

Otol. (Leptolepidarum) modestus, aff. simplex, aldorfensis, malloides martini, probabilis, obliquus, similis figd., Weiler (3).

ELOPIDAE

Elops hawaiensis post larvae descr. and figd. from Luzon, Blanco & Villadolid.

†Laminospondylus gen. nov. type L. transversus sp. nov. U. Cretaceous Texas, Springer, V. G. p. 135, 1 pl. 1 text-fig.

Otol. (Elopoidarum?) postconvexus, ellipticus figd., Weiler (3).

ALBULIDAE

Albula vulpes third Texas record, Leary (2).

Pterothrissus †umbonatus otoliths figd. from London Clay of Bognor, Stinton (7).

Otol. (Pterothrissidarum) cf. galtinus figd., Weiler (3).

ALEPOCEPHALIDAE

Alepocephalus rostratus from Algerian coast, Dieuzeide & Roland (2).

Xenodermichthys copei from Icelandic waters, Kotthaus & Krefft.

SEARSIDAE

Holtbyrnia macrops sp. nov. Madeira, Maul p. 11 figd.

Searsia schnakenbecki first E. Greenland record, Kotthaus & Krefft.

CLUPEIDAE

List of natural hybrids, Slastenenko.

Baltic herring, development in late Quaternary times, Munthe.

Genealogy of gill-helix, Takahasi.

Alosa alosa, A. fallax killarniensis Irish records, Gibson (1); A. kessleri kessleri, A. k. volgensis development of larvae, Braginskaya; A. pseudoharengus distribution in Great Lakes, Miller, R. R. (4); A. sapidissima recaptures of tagged specimens, Vladykov (1); prenuptial behaviour, Medcof.

Brevoortia patronus bionomics, description of larvae and young, Suttkus (3).

Caspialosa pontica reproduction in the Danube, Teodorescu-Leonte & others.

Clupea fishery of, Hodgson; C. harengus age composition of Manx population, Bowers (1), (2), (3); southern

North Sea populations, Cushing & Burd; research on Atlantic populations, Leim & others; spawning places, Yudanov; development of pronephros, Holstvoogd; C. harengus membras development in water of high salinity, Krizhanovski (1); C. pallasii structure and activity of spermatozoa, Yanagimachi (2); entrance of sperm into egg, Yanagimachi (3); use of Ringer's solution in experimental work on fertilization and development of eggs, Yanagimachi (1); C. †sardinites figd. from Oligocene Hesse, Weiler (1); C. †trolli sp. nov. otoliths Pannonian Vienna Basin, Papp & Thenius p. 34 pl. vi figs. 45, 46.

[1957]

Clupeonella delicatula caspia behaviour in captivity, Nikonorov.

Dastilbe †moraesi sp. nov. Cretaceous Brazil, Santos p. 19 pls. i, ii.

Dorosoma cepedianum distribution in Great Lakes, Miller, R. R. (4).

Engraulis japonica vertebral counts on post-larvae correlated with growth, Hayashi & Suzuki.

Fluvialosa erebi figd., Whitley (2).
Gadusia chapra pharyngeal pockets,
Kapoor, B. G. (3).

Opisthonema oglinum aggregation in rough water, Springer, S. (2).

Sardina pilchardus population study, Cruz (1); distribution and number of in Channel, Cushing (2); growth in Jugoslav waters, Muzinic; distribution of eggs and larvae off Balearie Isl., Massuti; parasitized by Peroderma, Dieuzeide & Roland (4); parasitic castration, Santos Pinto (1); gonadal tumours, Santos Pinto (2); S. pilchardus, Sardinella aurita biometric study, Dieuzeide & Roland (3).

Sardinella aurita biometric study, Monteiro; S. leiogaster new Borneo record, Inger.

Sardinops occillata bionomics, Davies (3); feeding habits, Davies (2); size composition of commercial catches, Davies (1).

CHANIDAE

Chanos chanos post larvae descr. and figd. from Luzon, Blanco & Villadolid; culture in Philippines, Rabanal & others.

PHRACTOLAEMIDAE

Phractolaemus ansorgii recorded from Stanley Pool, Boeseman (2).

SALMONIDAE

Natural hybrids, Slastenenko.

Reactions to light in different conditions, Fields.

Evolution and distribution of Coregonus, Prosopium and Leucichthys, Smith, S. H.

Coregonus of Lake Imandra, Russia, Vladimirskaya (1); results of acclimatising in Roumania, Busnita & others (1); speciation and population studies, Svärdson (1); C. lavaretus, C. nasus bionomics of Sundsvall Bay populations, Lindroth (1); C. lavaretus maraenoides biological heterogeneity of introduced specimens, Shirkova; C. lavaretus maraenoides × C. lavaretus cross breeding and feeding habits of hybrids, Gasowska; C. sardinella bannti taxonomic position, Anpilova.

Oncorhynchus gorbuscha variation in size of year classes in relation to spawning migration, Birman (1); O. keta development in Amur river, Disler; a Janus monstrosity, Honma (4); population studies in Amur basin, Birman (2); O. masu ecological notes, Semko; O. masu morpha formosanus a non-migratory form in Sedanka river, Moiseev (2); O. masou, O. rhodurus distribution in rivers of Kiushiu, Japan, Oshima (4): O. nerka distribution in two British Columbia lakes, Johnson, W. E.; population study, Vernon; fluctuations of stock in Kamtchatka waters, Krogius & Krokhin; O. tshawytscha feeding habits, Merkel; survival of mature male parr after spawning, Robertson, O. H.

Salmo structure and morphogenesis of dorsal fin, François; S. clarkii, interhybrids, Donaldson racial permanence and size of home territory, Miller, R. B. (1); S. clarki, S. gairdneri scale characteristics of yearlings compared, Vernon & McMynn; S. gairdneri alterations in blood constituents following muscular activity, Black (1); growth, Larkin, Terpenning & Parker; S. irideus liver regeneration, Faktorovich (1); S. letnica, Sapkarev (1); variation in fat content, Sapkarev (2); S. salar age, growth and movements of Foyle river specimens, Went & Vickers; causes of fluctuations of Baltic stocks, Lindroth (2); migrations, Azbelev & Lagunov; differential characters of populations of tributaries of the same

river-system, age and season of smolt migrations, Went (5): from the Mersey. Anon. (1); investigation on 'late running' specimens in Taw and Torridge estuaries, Cameron: large Irish examples. 'Avondhu'; movements around Ireland, Toner; Went (1); results of tagging in Labrador, Blair (1) & (2); movements off N.E. coast of England, Hartley & Brayshaw; disappearance from the Elbe, Bauch; S. trutta, S. salar, comparison of growth rates in Cashla river, Went (2); S. trutta growth rate before and after raising of lake level, Frost (1); of Lough Rea, Co. Galway, Healy (3); growth rate of in Fergus river system, Co. Clare, Healy (4); age and growth of Ilen river examples, Went (6); age and growth of Loch Kernsary specimens, Swan (1); feeding habits in a Swedish stream, Nilsson; migration, age and growth in Lake Rensjön, Runnström; notes on wounded specimens, Smith, I. W.; of R. Baracca, Liguria, Arbocco; organogeny, Bernard; retinal growth, Lyall (1); S. trutta, Salvelinus alpinus, bionomics of in Murmansk region, Vladimirskaya (2).

Salvelinus, spawning in Llyn Bodlyn, N. Wales, in November, Jones, R. W.; forms occurring in southern Saghalien, Ishida; S. alpinus notes on Icelandic examples, Stewart; osmoregulation, Gordon, M. S.; S. colii Lough Corrib specimens, Went & Twomey; S. gracillimus from Shetland, Swan (2); S. namayoush changes in blood levels of lactic acid following muscular activity, Black (2).

Thymallus bionomics of in Murmansk lakes, Vladimirskaya (3); T. arcticus baicalensis brevipinnis reproduction, Tugarina.

MICROSTOMIDAE

Bathylagus benedicti description of retina, ∇ ilter.

Microstoma microstoma recorded from Gulf of Mexico, Grey; M. pacificus a tagged example recaptured after six years, Best (2).

OSMERIDAE

Hypomesus sakhalinus sp. nov. L. Taraika, Sakhalin, Hamada p. 137 figd.

Spirinchus dilatus new Californian record, Jensen.

SALANGIDAE

Salangichthys microdon ecological observations during breeding season, Okada & Mori.

GONOSTOMATIDAE

Cyclothone braueri from E. Greenland waters, Kotthaus & Krefft; C. microdon Japanese examples discussed, Tomiyama & Abe.

Valenciennellus tripunctulatus from Gulf of Mexico, Grey.

STERNOPTYCHIDAE

Argyropelecus gigas from Gulf of Mexico, Grey; A. hemigymnus from Greenland, Kotthaus & Krefft.

OSTEOGLOSSIDAE

Arapaima gigas bionomics, Mig-dalski.

Osteoglossum bicirrhosum, behaviour in aquarium, Heck.

Scleropages, abundance and size in Malaya, fig., Locke.

NOTOPTERIDAE

Notopterus sp., photograph, Scott, J. S.

MORMYRIDAE

Control of electric discharge by 'paraseptal nucleus,' Szabo (1).

HAPLOMI

ESOCIDAE

Natural hybrids, Slastenenko.

Esox lucius systematic and biological notes, Čihăr; specimen lacking pelvic fins, Herbst; introduction into Irish waters, Went (3); growth rate in Lough Rea, Galway, Healy (2); age and growth of Irish specimens, Austin; developmental stages, Shamardina; effect of external factors on development, Tschörtner; food of young, Makkoveeva; E. masquinongy growth, mortality and movements of a population, Crossman.

UMBRIDAE

Novumbra hubbsi extension of range, Fitzgerald.

INIOMI

Larval and juvenile forms from the Mediterranean, Padoa & others.

SUDIDAE

[1957]

Key to genera and species, Kamohara (1).

Bathypterois dubius from Algerian coast, Dieuzeide & Roland (2).

Bathytyphlops gen. nov. type Bathymicrops sewelli Norman, Nybelin p. 258.

Chlorophthalmus borealis sp. nov. Chosi, Japan, Kuronuma & Yamaguchi p. 272.

Мусторнідав

Ontogenetic differences in Mediterranean forms, Padoa & others.

Lampadena nitida structure of precaudal photophores, Anadon.

Lampanyctus leucopsarus age and growth, Bolin; L. †serratus sp. nov. otoliths Miocene New Zealand, Stinton (2) p. 514 pl. xxxii figs. 1-4.

Myctophum nitidulum, a distinct species, Gibbs (2); M. (Myctophum) punctatum from Greenland waters, Kotthaus & Krefft.

"Scopelus" †ovatus sp. nov. otoliths Miocene New Zealand, Stinton (2) p. 514 pl. xxxii figs. 10, 11.

ANOTOPTERIDAE

Anotopterus pharao from Kiska Isl., Aleutians, Welander, Alverson & Bergman.

PARALEPIDIDAE

Paralepis key to postlarvae of Mediterranean species, Padoa & others.

ATELEOPIDAE

Ijimaia antillarum from Gulf of Mexico, Kanazawa.

OSTARIOPHYSI CYPRINOIDEA

Parallel evolution in some Cyprinids and Catostomids, Kafuku.

CHARACIDAE

Natural hybrids, Slastenenko.

Anoptichthys presence of a gene for albinism, Sadoğlu; A. hubbsi, Astyanax mexicanus observations on behaviour, John; A. jordani, general account, Lüling; colour sensitivity, Thines & Kähling.

Cheirodon axelrodi Schultz and Hyphessobrycon cardinalis Myers & Weitzman, relative priority, Opinion 485.

Cyrtocharax squamosus, Aramburu (1).

Hemigrammus armstrongi perhaps = Cheirodon piaba, Ladiges.

Hydrocyon vittatus, Groenewald.

Hyphessobrycon griemi sp. nov. Gojas, Brazil, Hoedeman (2) p. 87 figd.

Moenkhausia pittieri polycystic kidneys, Stolk (2); M. sanctae filomenae, life-colours and description, Meinken (3).

Moojenichthys myersi gen. et sp. nov. Brazil, Ribeiro (1) p. 546 figd.

Petersius caudalis, breeding in aquarium, Poersch.

Thayeria obliqua carcinoma of kidneys, Stolk (5).

HEMIODONTIDAE

Nannostomus beckfordi aripirangensis method of fighting, Wickler (3).

STERNARCHIDAE

Porotergus ellisi sp. nov. Parana river, Argentina, Aramburu (3) p. 154 figd.

GYMNOTIDAE

Parallelism with Mormyridae, Atz & Coates.

Structure of electric organs, Szabo (2).

ELECTROPHORIDAE

Electrophorus electricus behaviour in captivity, Boursot.

CATOSTOMIDAE

Natural hybrids, Slastenenko.

Distribution in Minnesota, Underhill.

.Carpiodes carpio carpio age and growth, Buchholz.

Minytrema melanops discussion of Chesapeake Bay record, Robins & Raney (1).

Moxostoma (Scartomyzon) key to species of Texas and Mexico; relationships, Robins & Raney (2); M. austrinum milleri subsp. nov. Mexico, Robins & Raney (2) p. 307 figd.; M. carinatum notes on southern range, Robins & Raney (1); M. valenciennesi recorded from North Dakota, Robins & Raney (1).

Pantosteus jordani recorded from Alberta waters, Scott, W. B.

Ptychostomus haydeni Girard a synonym of Moxostoma aureolum, Robins & Raney (1).

CYPRINIDAE

Natural hybrids, Slastenenko.

Distribution in Minnesota, Underhill.

Adaptive significance of horn teeth during development, Eremeeva.

Abramis description of an unnamed form from Germany, Miegel (2); A. brama larval development, Dmitrieva (1); Vasnetzov & others; spawning and early development in Vistula lagoon, Dziekonska; population study in Vistula region, Poczopko (2); introduction into Khristoforov reservoir, Melnikov; comparison of Danube and Dnieper river populations, Pavlov; A. brama orientalis effect on stock of heavy fishing, Bervald.

Aspius aspius growth and reproduction, Troitzkii; feeding habits, Vasiliu & Nicolau (2).

Barbodes, type and definition, Schultz (3); Meinken (4).

Barbus Cuvier & Cloquet, notes on status, Hoedeman (3); restriction of genus, Schultz (3); Meinken (4); B. longiceps × Varicorhinus damascinus description of hybrid, Steinitz & Ben-Tuvia; B. schneemanni sp. nov. E. Africa, Klausewitz (11) p. 279 fig.; B. tetrazona renal adenocarcinoma, Stolk (7); B. ef. yunnanensis, B. †brevicephalus, B. szechuanensis figd. from L. Pliocene Choukoutien, Liu (1).

Biwia tama sp. nov. Tama river, Japan, Oshima (3) p. 16 figd.

Blicca bjoerkna influence of light on feeding, Dudek.

Caecobarbus geertsi, general account, Lüling.

Capoeta, type and definition, Schultz (3).

Capoetobrama kuschakewitschi bionomics, Maxunov.

Carassius ecology and taxonomy of two species, **Dmitrieva** (2); C. auratus ecology of during introduction into Veselovsk reservoir, **Shchetinina**; coiling of intestine during development, **Yoshi**zaki. Chondrostoma nasus ohridanus development, Točko.

Chrosomus eos recorded from Saskatchewan, Scott, W. B.; C. erythrogaster erythrogaster recorded from Lower Mississippi drainage, Hemphill.

Cirrhina mrigala, age-determination, Jhingram.

Cyprinella status of subgenus, with key to species, Gibbs (1).

Cyprinus morphological changes in specimens acclimatised in Lake Balkhash, Burmakin; oxygen consumption of fry, Syazuki & Hayama; C. carpio different forms of scales, Dürr; growth of mirror and leather carp, Mengel & Steffens: some unusual swimbladders, Schreiter; temperature selection, Pitt & others; results of artificial hybridization, Lieder; changes in pituitary during maturation of gonads, Irkimovich & Zelenin; larval development, Vasnetzov & others; vascular system of liver, Friedrich; structure of glands associated with digestive system, Santa & Pollingher; main, accessory and emergency sources of food, Vaas; C. carpio, Mylopharyngodon piceus, Ctenopharyngodon idellus rec. from Pleistocene Honan, Huang; C. †maomingensis sp. nov. Miocene Kwantung China, Liu (2) p. 151 2 figs.

Dionda diaboli sp. nov. Texas, Hubbs & Brown p. 69 figd.

Eustira Gunther a synonym of Danio Hamilton, E. ceylonensis a synonym of D. malabaricus, Silas.

Gobio gobio, G. uranoscopus, G. kessleri notes on Roumanian specimens, Nalbant (1).

Hybognathus hankinsoni recorded from Saskatchewan, Scott, W. B.

Hypophthalmichthys molitrix structure and function of suprabranchial, Zambriborshch; structure of gill apparatus, Verigin.

Labeo altivelis spawning grounds in Katanga Province, Bont & Maes; L. bicolor colour variation, Klausewitz (8).

Leucaspius delineatus development and growth of scales, Balon (1); from Lake Neuchatel, Switzerland, Quartier.

Leuciscus presbyter C. et V. a synonym of Barbus (Puntius) amphibius (C. et V.), Guibé & Spillmann; L. rutilus

gastric epithelium, Bernard; L. waleckii suiyuani subsp. nov. Pao-Tou, N. China, Mori, T. p. 183 figd. See also Squalius.

Matsya †hsichihi sp. nov. Lower Pliocene, Choukoutien, China, Liu (1) pp. 7, 19 text-figs. 4, 5 pl. iii.

Notropis ortenburgeri, N. maculatus, N. blennius added to E. Texas fauna, Hubbs, C. & Kemp; N. rubellus spawning behaviour, Reed (1); feeding habits and growth rate, Reed (2); N. venustus redescribed and subspecies characterized, Gibbs (3).

Osteochilus hasseltii (C. et V.) a synonym of O. vittatus (C. et V.), Guibé & Spillmann; main, accessory and emergency sources of food, Vaas.

Pfrille neogaea recorded from Sas-katchewan, Scott, W. B.

Pungtungia shiraii sp. nov. Tama river, Japan, Oshima (3) p. 15 figd.

Puntius, type and definition, Schultz (3); Meinken (4); P. javanicus, main, accessory and emergency sources of food, Vaas; P. roloffi sp. nov. Thailand, Klausewitz (9) p. 193 figd., Klausewitz (2).

Rasbora nigromarginata sp. nov. Ceylon, Meinken (2) p. 66 figd.

Rhodeus amarus spawning behaviour, Wiepkema.

Rutilus frisii development, Smirnova, E. N. (2); R. pigus virgo biometric study and sexual dimorphism, Misik; R. rutilus development and growth of scales, Balon (1); development of Aral Sea examples, Gosteeva; R. rutilus caspicus behaviour on feeding grounds, Novikova; larval development, Vasnetzov & others; R. rutilus heckeli larval development, Dmitrieva (1); Vasnetzov & others; R. rutilus lacustris histology of muscles, Lobintzev; R. rutilus × Chalcalburnus chalcoides, Nalbandoglu. See also Leuciscus.

Scardinius erythrophthalmus growth rate in Lough Rea, Galway, Healy (2); S. erythrophthalmus †brevicaudus subsp. nov. U. Tertiary Siberia, Yakovlev p. 63.

Schizothorax argentatus bionomics, Savina.

Squalius leuciscus distribution, bionomics in Luxemburg, Hoffmann.

Tinca tinca from Lough Lein, Killarney, Healy (1); results obtained from an electro-cardiogram, Serfaty & Raynaud.

Toxabramis paiyantieni sp. nov. Paoting, N. China, Mori, T. p. 182 figd.

Tribolodon hakuensis feeding habits, Imamura & Hashitani.

Vimba vimba natio carinata feeding habits, Vasiliu & Costea; enzymes of the hatching glands, Baburina & Buznikov; peculiarities in development, Smirnova, E. N. (1).

COBITIDAE

Natural hybrids, Slastenenko.

Acanthophthalmus critical review, Moldenhauer; A. cuneovirgatus p. 30 figd. Johore, Singapore, A. robiginosus p. 31 figd., Java spp. nov., Raut.

Botia horae, B. modesta, B. lecontei Klausewitz (1), (4), (8); B. lohachata, Klausewitz (5); behaviour in aquarium, Bader.

Cobitis aurata vallachica subsp. nov. Rumania, Nalbant (2) p. 209 figd.; C. elongata relationships, Banarescu & Nalbant; C. taenia embryology, Kokhanova; C. taenia, C. aurata notes on Roumanian specimens, Nalbant (1).

Noemacheilus phuketensis sp. nov. Thailand, Klausewitz (9) p. 195 figd.

SILUROIDEA

ARIIDAE

Galeichthys felis reproduction and early development, Ward.

DORADIDAE

Auchenipterus nuchalis, Aramburu (4).

PLOTOSIDAE

Copidoglanis ater recorded from Queensland, Marshall, T. C.

SILURIDAE

Remains from Wadi Hoff, Egypt, Moustafa.

Centromochlus creutzbergi recorded from Surinam, Hoedeman (4).

Wallagonia attu spawning habits and early development Mazumdar; sensory canals of head, Kapoor, A. S.

Wallago attu a teratological example, Sarkar & Kapoor.

BAGRIDAE

Status of Bagre, Bagrus and Porcus, Jayaram.

Coreobagrus generic description emended, Okada & Kubota (2); C. ichikawai sp. nov. Miya river, Mie Prefecture, Japan, Okada & Kubota (2) p. 144 figd.

Mystus tengara teratological examples, Sarkar & Kapoor.

Pseudobagrus aurantiacus pit organ system, Satô.

Rita rita structure of barbels, Bhatti (1); cutaneous sense organs, Bhatti (2).

AMEIURIDAE

Natural hybrids, Slastenenko.

Distribution in Minnesota, Underhill.

Ameiurus nebulosus food, Frank; number of eggs produced, Sedlár.

Ictalurus melas occurrence of blind specimens, Mayhew (2); I. punctatus growth, Harrison.

Schilbeodes miurus infestation by parasite Clinostomium, Schwartz.

SCHILBEIDAE

Eutropius debauwi recorded from Stanley Pool, Boeseman (2).

CLARIIDAE

Clarias palaeolithic remains from Mumba Cave, E. Africa, Greenwood (2); C. batrachus induced spawning, Ramaswami & Sundararaj (2).

Gymnallabes tihoni redescribed, Poll (3).

Heterobranchus longifilis, in R. Kilombero, Tanganyika Territory, Ford.

Heteropneustes fossilis induced spawning, Ramaswami & Sundararaj (1).

MOCHOCIDAE

Microsynodontis batesii recorded from Stanley Pool, Boeseman (2).

Synodontis membranaceus consideration of bony rays in adipose fin, Sandon; S. soloni recorded from Stanley Pool, Boeseman (2); S. werneri sp. nov. Stanley Pool, Belgian Congo, Boeseman (2) p. 146 figd.

MALAPTERURIDAE

Malapterurus electricus fibromas of skin, Stolk (5).

PYGIDIIDAE

Eremophilus camposi sp. nov. rio Juquia, Brazil, Ribeiro (2) p. 72 figd.

CALLICHTHYIDAE

Cascadura Ellis based on juvenile example of Hoplosternum thoracatum, Hoedeman (5).

Corydoras griseus de weyeri subsp. nov. British Guiana, Meinken (1) p. 5 figd.

Hoplosternum, Callichthys notes on Surinam species, Hoedeman (4).

APODES

Colours in life, Kuroda (1).

ANGUILLIDAE

Anguilla spawning areas of African species, Jubb; notes on species occurring in Zambesi river system, Frost (2); A. anguilla distribution and migrations, Raben; food of in Baltic, Murina; estimation of age, using scales and otoliths, Rahn; scent perception, Teichmann; effect of hormone injection on sex, Ancona (5); structure of auricular muscular tissue, Couteaux & Laurent; cause of cauliflower disease, Lühmann & Mann; A. obscura recorded from Southern Africa, Jubb.

XENOCONGRIDAE

Ariosoma impressa from Puerto Rico, Erdman.

Kaupichthys atlanticus from Puerto Rico, Erdman.

MURAENIDAE

Channomuraena vittata new Puerto Rican record, Erdman.

Echidna delicatula, E. leucotaenia, E. nebulosa, E. polyzona, E. zebra, new Borneo records, Inger.

Enchelycore nigricans from Puerto Rico, Erdman.

Gymnothorax buroensis, G. chilospilus, G. flavimarginatus, G. hepaticus, G. javanicus, new Borneo records, Inger.

HETERENCHELIDAE

Heterenchelys biaggii from Puerto Rico, Erdman; H. †crossus sp. nov. otolith Miocene New Zealand, Stinton (2) p. 514 pl. axxii fig. 6.

NEMICHTHYIDAE

Nemichthys scolopaceus from Algerian coast, Dieuzeide & Roland (2).

MURAENESOCIDAE

Muraenesox cinereus spermatogenesis, Nishikawa.

CONGRIDAE

Heterocongrinae, key to genera, Böhlke (3).

Astroconger myriaster length/weight relationship, Okada & Suzuki; pit organ system, Satô.

Coloconger meadi sp. nov. Gulf of Mexico, Kanazawa p. 234 figd.

Conger †acutus sp. nov. otolith London Clay Bognor, Stinton (1) p. 18 pl. ii fig. 3.

Congermuraena mystax description of eggs, Thomopoulos (1).

Nystactes halis gen. et sp. nov. Green Cay, Bahamas, Böhlke (3) p. 68 figd.

Ophisoma balearicum description of eggs, Thomopoulos (1).

Uroconger †minimus figd. from London Clay of Bognor, Stinton (1).

ECHELIDAE

Myrophis egmontis from Puerto Rico, Erdman.

OPHICHTHIDAE

Aplatophis chauliodus from Puerto Rico, Erdman.

Caecula imberbis description of eggs, Thomopoulos (1).

Leiuranus semicinctus new Borneo record, Inger.

? Mystriophis tobliquum sp. nov. otolith Miocene New Zealand, Stinton (2) p. 513 pl. xxxii fig. 5.

SYNAPHOBRANCHIDAE

Synaphobranchus kaupii recorded from Nabuto, Japan, Kuroda (3).

HETEROMI

†DERCETIDAE

Stratodus sp. rec. U. Senonian Israel, Avnimelech.

SYNENTOGNATHI

BELONIDAE

Key to genera, Fernandez Yepez.

Belone depressa Poey a valid species, Fernandez Yepez.

Tylosurus leiurus new Japanese record, Kamohara (2).

SCOMBRESOCIDAE

Scomberesox saurus Argentine specimens described, Lopez.

HEMIRAMPHIDAE

Euleptorhamphus velox from Puerto Rico, Erdman.

Hemiramphus, Zenarchopterus, postlarvae described and figd. from Luzon, Blanco & Villadolid; H. gaimardi C. & V. a synonym of Hyporhamphus melanurus (Val.), Inger.

EXOCOETIDAE

Bionomics and fishery, Mohr.

Cypselurus rondeletii new Red Sea record, Tortonese.

Exocoetus volitans Japanese examples discussed, Tomiyama & Abe.

Fodiator acutus redefinition of subspecies, Morrow (2); F. a. pacificus Bruun a synonym of F. a. rostratus (Gunther), Morrow (2).

Hemiexocoetus caudimaculatus Fowler a synonym of Fodiator acutus rostratus (Gunther), Morrow (2).

Hirundichthys affinis bionomics and fishery, Hall.

Parexocoetus brachypterus post-larvae descr. and figd. from Luzon, Blanco & Villadolid.

MICROCYPRINI

AMBLYOPSIDAE

Key to genera and species of United States, distribution, Woods & Inger.

CYPRINODONTIDAE

Ecology; account of African and S. American annual species, Myers.

Natural hybrids, Slastenenko.

Anableps dovii breeding habits, Tro-chanis.

Aphyosemion bivittatum account of "spawning" between two males, Nieuwenhuizen; A. cognatum, colour and behaviour in aquarium, Foersch; A. elegans recorded from Stanley Pool, Boeseman (2).

Aplocheilichthys maculatus sp. nov. E. Africa, Klausewitz (11) p. 281 figd.

Aplocheilus luzonensis post larvae descr. and figd. from Luzon, Blanco & Villadolid; A. panchax dorsomarginatus subsp. nov. Thailand, Klausewitz (9) p. 196 figd.

Austrofundulus and Cynolebias secondary sexual characters Myers.

Cynolebias (Leptolebias) subgen. nov. type Cynopoecilus marmoratus Ladiges. 1934, Myers p. 140; C. aureoguttatus sp. nov. Paranagua, Brazil, Myers p. 139 figd.; C. zingiberinus Myers a synonym of C. marmoratus (Ladiges), Myers.

Cynopoecilus Regan a synonym of Cynolebias Steindachner, Myers; C. sicheleri M. Ribeiro a synonym of Cynolebias marmoratus (Ladiges), Myers.

Cyprinodon fasciatus, C. iberus French records, Arnoult.

Epiplatys multifasciatus recorded from Stanley Pool, Boeseman (2).

Fundulus key to species of eastern U.S.A. and Canada, Brown (2); F. cinqulatus, F. chrysotus distinguishing characters, Brown (1); F. hispanicus French records, Arnoult.

Gambusia a revision of the G. nobilis species group, with key, Hubbs, C. & Springer; systematic study of specimens acclimatized in Russia, Gogol; pectoral support of gonopodium during copulation, Hubbs & Reynolds; G. geiseri sp. nov. San Marcos river, Texas, Hubbs & Hubbs in Hubbs & Springer p. 285 figd.

Gambusia heterochir sp. nov. Texas, Hubbs, C. (4) p. 3 figd.; reproductive behaviour, Warburton & others; hybridising with G. affinis, Hubbs, C. (4); G. hurtadoi p. 307 figd., G. alvarezi p. 310 figd. spp. nov. Chihuahua, Mexico, Hubbs, C. & Springer.

Jenkinsia lamprotaenia schooling behaviour, Springer, S. (2); J. lineata breeding cycle, Turner.

Lebistes reticulatus fibroma of heart, Stolk (7); rectal fibroma, Stolk (10); adenoma of swim bladder, Stolk (12); cerebral cysts and abnormal glands, Stolk (9); morphology of rays of anal fin, Mohsen; cytology of pituitary, Vervoort.

Leptolebias ${f subgen.\ nov.}$ of Cynolebias ${f q.v.}$

Nothobranchius, drought - resistant eggs and conditions of life, Hansen.

Oryzias latipes regeneration of caudal fin, Mori, H.; feeding habits, Imamura & Hashitani.

Rivulus discussion of species complex, Hoedeman (1).

SOLENICHTHYES

AULOSTOMIDAE

Aulostomus maculatus association with scarid, Eibl-Eibesfeldt.

CENTRISCIDAE

Centriscus (Amphisyle) †heinrichi figd. Snigielska.

SYNGNATHIDAE

Corythoichthys flindersi sp. nov. S. Australia, Scott, T. D. p. 182 figd.

Hippocampus guttulatus histology of kidney, Gabe; H. hippocampus from Co. Clare, Ireland, Roche (2).

Ichthyocampus kampeni new Bornean record, Inger.

Phycodurus eques col'd illustration, Hale.

Phyllopteryx foliatus, photo of ♂ with eggs, Hale.

Syngnathus rostellatus recorded from British coast, Wheeler, A. C.

ANACANTHINI

MACROURIDAE

Cetonurus microps Vaillant a nomen nudum, Nybelin.

Chalinura key to some N. Atlantic species, Nybelin; C. profundicola sp. nov. North Atlantic, Nybelin p. 268.

Coelorhynchus coelorhynchus recorded from Greenland waters, Kotthaus & Krefft; C. †elevatus sp. nov. otolith Miocene New Zealand, Stinton (2) p. 515 pl. xxxii fig. 13. Hymenocephalus? sp. figd. from Oligocene E. Carpathians, Ghorbach.

Lionurus carapinus euabyssalis subsp. nov. North Atlantic, Nybelin p. 273 figd.

Trachyrhynchus †densus sp. nov. Miocene New Zealand, Stinton (2) p. 516 pl. xxxii fig. 15.

MERLUCCHDAE

Merluccius merluccius size differences between populations, Matta; M. merluccius, M. capensis, M. polli reviewed, Franca (1).

GADIDAE

Natural hybrids, Slastenenko.

Morphological principles of classification, Syctovidov (4).

Bregmacerina gen. nov. type Motella tricirrata antiqua Smirnov, Danil'chenko p. 57 1 text-fig. Oligocene N. Caucasus.

Bregmaceros atlanticus bionomics, Clancey.

Gadus callarias migrations of Bear Island populations, Trout; distribution and ecology compared with that of Clupea harengus, Svetovidov (2); pancreas as source of insulin, Roy; G. poutassou distribution of young, Henderson; G. †subnotus otolith figd. from London Clay of Bognor, Stinton (1).

Gaidropsarus ensis from E. Greenland waters, Kotthaus & Krefft.

Melanonus unipennis from Gulf of Mexico, Grey.

Pollachius virens biology and fishery in Barents Sea, Mironova (2).

Raniceps †planus otolith figd. from London Clay of Bognor, Stinton (1).

Urophycis blennoides from Irish waters, Gibson (4).

Ot. (Gadidarum) †laevigatum sp. nov. Miocene New Zealand, Stinton (2) p. 515 pl. xxxii fig. 7. Ot. (Gadidarum) †ponticum sp. nov. Pannonian Vienna Basin, Papp & Thenius p. 39 pl. vi figs. 47, 48.

ALLOTRIOGNATHI

LOPHOTIDAE

† Protolophotus gen. nov. type Lophotus elami Arambourg, Walters (1) p. 60.

BERYCOMORPHI

POLYMIXIIDAE

Polymixia lowei from Puerto Rico, Erdman.

BERYCIDAE

Beryx decadactylus European and Norwegian records, Willgohs; B. splendens from Gulf of Mexico, Woods.

DIRETMIDAE

Diretmus argenteus from Gulf of Mexico, Grey.

CARISTIIDAE

Caristius macropus recorded from British Columbia, Welander, Alverson & Bergman.

MONOCENTRIDAE

Monocentris †lemoinei otolith figd. from London Clay of Bognor, Stinton (1); M. reedi sp. nov. Chile, Schultz (1) p. 238 figd.

HOLOCENTRIDAE

Holocentrus cornutus, H. lacteoguttatus, H. sammara new Bornean records, Inger.

Osteichthys trachypomus from Puerto Rico, Erdman.

Plectrypops retrospinis from Puerto Rico, Erdman.

MELAMPHAIDAE

Melamphaes cristiceps recorded from Japanese waters, Maruyama.

ZEOMORPHI

Larvae and juveniles from the Mediterranean, Padoa & others.

ZEIDAE

Allocyttus verrucosus recorded from North Pacific, Welander & others.

Parazen pacificus first Atlantic record, described and figd., Mead (1).

Xenocyttus gen. nov. p. 225 type X. nemotoi sp. nov. Antarctic p. 228 figd., Abe (3).

PERCOMORPHI

PERCOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

SERRANIDAE

Alphestes scholanderi sp. nov. Bahamas, Walters (2) p. 283 figd.

Anyperodon leucogrammicus new Bornean record, Inger.

Aporops japonicus sp. nov. Riu Kiu Islands, Kamohara (3) p. 21 figd.

Dermatolepis inermis from Puerto Rico, Erdman.

Ellerkeldia annulata recorded from Queensland, Marshall, T. C.

Epinephelus corallicola, E. merra, new Bornean records, Iuger; E. fusco-guttatus, E. morrhua post larvae descr. and figd. from Luzon, Blanco & Villadolid; E. gigas from Irish waters, Went (4); E. leprosus recorded from Mozambique, Smith, J. L. B. (5); E. mystacinus from Puerto Rico, Erdman; E. †ovalis sp. nov. otolith London Clay Bognor, Stinton (1) p. 23 pl. ii figd.; E. striatus, association with Elacatinus, Eibl-Eibesfeldt.

Grammistes sexlineatus new Bornean record, Inger.

Labrax (Morone) †serrata sp. nov. otoliths Pannonian Vienna Basin, Papp & Thenius p. 38 pl. vi figs. 31, 32.

Luzonichthys Herre, species included in genus, Smith, J. L. B. (7).

Mycteroperca interstitialis, M. venenosa from Puerto Rico, Erdman.

Prionodes tigrinus, P. phoebe, P. tabacaria from Puerto Rico, Erdman.

Promicrops sp., photograph, Scott, J. S.; P. itaiara from Puerto Rico, Erdman; P. lanceolatus figd., Whitley (2).

Roccus comparative osteology, Woolcott.

Therapon jarbua feeding habits, Imamura & Hashitani.

Otolithus (Serranidarum) †undulatus sp. nov. London Clay Bognor p. 22 pl. ii fig. 9; Ot. (S.) minsterensis figd., Stinton (1).

PSEUDOCHROMIDAE

Nematochromis annae new Bornean record, Inger.

PLESIOPIDAE

Plesiops coeruleolineatus new Bornean record, Inger.

Pseudoplesiops typus new Australian record, Marshall, T. C.

CENTRARCHIDAE

Natural hybrids, Slastenenko.

Archoplites interruptus osteology, Dineen & Stokely.

Chaenobryttus gulosus bionomics, Larimore.

Lepomis gibbosus \times L. macrochirus fertility and characters of F_1 hybrids, Lagler & Steinmetz; L. macrochirus bionomics, Mayhew (1).

Micropterus coosae absent from Californian waters, Kimsey (1); M. dolomieu dolomieu age and growth of a 'stunted' population, Suttkus (1); M. salmoides results of a tagging programme, Kimsey (2).

PRIACANTHIDAE

Pseudopriacanthus altus first record for Maine coast, Scattergood & Coffin.

APOGONIDAE

Apogon post larvae deser. and figd. from Luzon, Blanco & Villadolid; A. †bellovacinus otolith figd. from London Clay of Bognor, Stinton (1); A. binotatus from Puerto Rico, Erdman; A. cardinalis, A. novemfasciatus new Bornean records, Inger; A. pigmentarius, A. pseudomaculatus northern extension of range, Caldwell & Briggs.

Pomatomichthys constanciae juvenile forms described and compared with young of Pomatomus telescopium, Cavaliere.

PERCIDAE

Growth study in north German eutrophic waters, Tesch.

Natural hybrids, Slastenenko.

Etheostoma blennioides, E. caeruleum, E. spectabile egg site selection, Winn; E. lepidum factors affecting fecundity, Hubbs & Strawn (1); E. spectabile × Percina caprodes relative variability of hybrids, Hubbs & Strawn (2).

Lucioperca lucioperca larval development, Dmitrieva (1); Vasnetzov & others.

Perca flavescens diurnal activity, Brown & Rosen; P. fluviatilis, food of in Volga delta, Ivanova, M. N.; of Lough Rea, Co. Galway, Healy (3); factors influencing sexual cycle, Morozova, T. E.; discovery of a plexus in digestive tube, Milokhin (2); P. fluviatilis, P. schrenki and Lucioperca lucioperca comparison of developmental stages, Konstantinov.

[1957]

Percarina demidoffi maeotica feeding habits, Kanaeva.

Sander lucioperca food of in Roumanian waters, Vasiliu & Nicolau (1).

Stizostedion v. vitreum age and growth, Rose.

Ot. (Percidarum) †uawaensis sp. nov. Miocene New Zealand, Stinton (2) p. 516 pl. xxxii fig. 8.

LATILIDAE

Malacanthus hoedtii new Australian record, Marshall, T. C.

SILLAGINIDAE

Sillago analis new Queensland record, Haysom.

RACHYCENTRIDAE

Rachycentron canadus from Buzzards Bay, Mass., Mather & Gibbs; from Puerto Rico, Erdman.

CARANGIDAE

Evolution, Aleev (7).

Angolan species, Franca (2).

Alectis crinitus from Alabama coast, Boschung; from Puerto Rico, Erdman.

Caranx lugubris, C. guara from Puerto Rico, Erdman; C. trachurus description of eggs, Thomopoulos (1).

Decapterus macarellus from Puerto Rico, Erdman.

Elagatis bipinnulatus from Puerto Rico, Erdman.

Seriola dumerili from Puerto Rico, Erdman.

Trachurus systematics of species in USSR waters, Aleev (6); taxonomy of Black Sea species, Aleev (4); T. mediterraneus ponticus commercial use in Black Sea, Tikhonov; T. trachurus reproduction, Tkacheva; T. trachurus mediterraneus feeding habits, Pora & others; movements in Black Sea, Nicolae & Cautis; T. trachurus trachurus structure and use of lateral muscles, Hasan.

BRAMIDAE

Brama raii from Puerto Rico, Erdman; from Moy river, Ireland, Roche (1).

Centropholoides Smith a synonym of Pterycombus Fries, Caldwell & Briggs.

Centropholis Hilgendorf a synonym of Pterycombus Fries, Caldwell & Briggs.

Collybus drachme recorded from Gulf of Mexico, Mead (2).

Pterycombus goodei recorded from Gulf of Mexico, Caldwell & Briggs.

Taractes key to species, Mead (2); T. longipinnis recorded from Gulf of Mexico; growth changes, Mead (2).

CORYPHAENIDAE

Coryphaena hippurus bionomics, Williams & Newell; recorded from Aldabra, Smith, J. L. B. (2); occurrence in a Texas bay, Pew.

CENTROPOMIDAE

Ambassis urotaenia new Bornean record, Inger.

Centropomus ensiferus from Puerto Rico, Erdman; C. †superpendens otolith figd. from London Clay of Bognor, Stinton (1).

Gymnochanda filamentosa gen. et sp. nov. Singapore, Boeseman (1) p. 75 figd.

Psammoperca †sheppeyensis otolith figd. from London Clay of Bognor, Stinton (1).

EMMELICHTHYIDAE

Emmelichthys nitidus new Japanese record, Kamohara (2).

LUTIANIDAE

Aphareus rutilans recorded from Mozambique, Smith, J. L. B. (5); recorded from Riu-Kiu Islands, Kamohara (3).

Caesio †bognoriensis sp. nov. otolith London Clay Bognor, Stinton (1) p. 25 pl. ii fig. 15.

Lutianus key to E. African species, with descriptions, Talbot; L. analis first Gulf of Maine record, Mather & Gibbs; L. buccanella from Puerto Rico, Erdman; L. niger recorded from Riu-Kiu Islands, Kamohara (3); L. russelli, new Bornean record, Inger.

Pentapus caninus new Bornean record, Inger.

Pristipomoides filamentosus roseus Japanese examples discussed, Tomiyama & Abe; P. macrophthalmus from Puerto Rico, Erdman; P. typus recorded from Mozambique, Smith, J. L. B. (5).

NEMIPTERIDAE

Nemipterus tolu new Australian record, Marshall, T. C.

LEIOGNATHIDAE

Pentaprion longimanus new Australian record, Marshall, T. C.

POMADASYIDAE

Brachygenys chrysargyreus from Puerto Rico, Erdman.

Pomadasys †ovalis sp. nov. otolith London Clay Bognor, Stinton (1) p. 26 pl. ii fig. 16.

SCIAENIDAE

Aplodinotus grunniens food of, Moen• Eques punctatus recorded from Florida, Caldwell & Briggs.

Johnius carutta new Australian record, Marshall, T. C.

Micropogon undulatus distribution and growth, Haven.

Stellifer stellifer from Puerto Rico, Erdman.

MULLIDAE

Mullus barbatus age and growth in Israel waters and description of post-larvae, Gottlieb.

Parupeneus malabaricus new Bornean record, Inger.

Upeneus sundaicus new Australian record, Marshall, T. C.

LETHRINIDAE

Gnathodentex mossambicus sp. nov. Mozambique, Smith, J. L. B. (5) p. 122 figd.

Pentapodus microdon new Australian record, Marshall, T. C.

SPARIDAE

Chrysophrys auratus weight and length data, Cassie.

Dentex †pentagonalis sp. nov. otolith London Clay Bognor, Stinton (1) p. 26 pl. ii fig. 17. Lagodon rhomboides systematics and bionomics, Caldwell.

Mylio latus, M. macrocephalus ecological study on young, Kasahara.

Sargus puntazzo, Oblada melanura, description of eggs, Thomopoulos (1).

Taius tumifrons regeneration of scales, Aoyama.

†Ot. (Sparidarum) kühni sp. nov. p. 35 pl. vi figs. 39, 40; and subsp. gracilis ssp. nov. p. 35 pl. vi figs. 41, 42; Ot. (S.) brunnense sp. nov. p. 35 pl. vi figs. 43, 44 all Pannonian Vienna Basin, Papp & Thenius.

MAENIDAE

Smaris insidiator description of eggs, Thomopoulos (1).

PEMPHERIDAE

Pempheris mulleri from Puerto Rico, Erdman.

KYPHOSIDAE

Kyphosus vaigiensis post-larvae described and figd. from Luzon, Blanco & Villadolid.

GIRELLIDAE

Girella elevata new Queensland record, Marshall, T. C.

SCATOPHAGIDAE

Scatophagus argus post-larvae deser. and figd. from Luzon, Blanco & Villadolid.

CHAETODONTIDAE

Natural hybrids, Slastenenko.

Anisochaetodon pictus Forskål a valid species, Klausewitz (7).

Chaetodon ocellatus, C. striatus recorded from Florida, Caldwell & Briggs; C. rafflesi, C. vagabundus new Bornean records, Inger; C. sedentarius from Puerto Rico, Erdman; C. speculum new Australian record, Marshall, T. C.; C. vitulus sp. nov. Jervis Bay, N.S.W., Whitley (2) p. 63 figd.

Holacanthus ciliaris range extension into north eastern Gulf of Mexico, Caldwell & Briggs.

Pomacanthus arcuatus from Puerto Rico, Erdman.

HISTIOPTERIDAE

Paristiopterus gallipavo new Queensland record, Marshall, T. C.

Pseudopentaceros richardsoni recorded from Japanese waters, Abe (1); recorded from North Pacific, Welander & others.

Quinquarius Jordan a synonym of Pseudopentaceros Gill, Welander & others.

NANDIDAE

Badis badis siamensis subsp. nov. Thailand, Klausewitz (9) p. 199 figd.

CICHLIDAE

Astronotus ocellatus regeneration of cranial nerve with re-innervation of muscles after severing, Arora & Sperry.

Cichlasoma Swainson retained in preference to Cichlaurus Swainson, Bailey; C. biocellatum changes in pituitary due to thyroidal tumour, Stolk (1); C. facetum multiple cysts in nervous system, Stolk (8).

Crenicichla jaguarensis, Geophagus brasiliensis Pirassununga specimens described, Travassos & Pinto.

Gephyrochromis lawsi sp. nov. L. Nyasa, with ecological notes, Fryer p. 346 figd.

Haplochromis mellandi used for molluse control, Bont (2) & (3); Bont-Hers, A. F. & M. J. de; H. multicolor structure of pharyngeal glands, Stolk (11); adenoma of pharyngeal glands, Stolk (12); H. sauvagei, H. prodromus, H. granti synonymy, distribution and ecology, Greenwood (1); H. xenognathus sp. nov. Lake Victoria, Greenwood (1) p. 90 figd.

Hemichromis bimaculatus from Tunisia, Postel (2).

Nannacara anomala, parental care, Kuenzer.

Pterophyllum eimekei, probable mutant, Buschendorf.

Symphysodon discus, structure of egg and larva, parental care, Wickler (1).

Tilapia palaeolithic remains from Mumba Cave, E. Africa, Greenwood (2); T. leucosticta diagnosis and biology, Lowe; T. mossambica, main, accessory and emergency sources of food, Vaas.

CEPOLIDAE

Acanthocepola abbreviata new Australian record, Haysom.

Cepola rubescens from Co. Waterford, Ireland, Gibson (3).

CIRRHITIDAE

Cirrhitoidea unimacula sp. nov. Kagoshima Prefecture, Japan, Kamohara (3) p. 29 figd.

APLODACTYLIDAE

Crinodus lophodon new Queensland record, Marshall, T. C.

CHEILODACTYLIDAE

Morwong gen. nov. type Chilodactylus fuscus Castelnau, Whitley (2) p. 65.

Nemadactylus douglasii, N. macropterus, figd., Whitley (2).

EMBIOTOCIDAE

Ditrema †sheppeyensis otolith figd. from London Clay of Bognor, Stinton (1); D. temmincki reproductive cycle, Ishii.

POMACENTRIDAE

Abudefduf anabatoides, A. oxyodon, A. sexfasciatus, new Bornean records, Inger; A. azysron new Red Sea record, Baschieri-Salvadori; A. saxatilis northern extension of range, Caldwell & Briggs.

Amphiprion frenatus, A. percula new Bornean records, Inger; A. percula, sound production in aquarium, Koenig; A. xanthurus new Red Sea record, Baschieri-Salvadori.

Cheiloprion labiatus new Bornean record, Inger; new Australian record, Marshall, T. C.

Chromis bitaeniatus new Australian record, Marshall, T. C.; C. marginatus, C. cyanea from Puerto Rico, Erdman; C. ternatensis new Red Sea record, Baschieri-Salvadori.

Chrysiptera melanomaculata sp. nov. Palau Isl., Aoyagi p. 180 figd.

Dascyllus aruanus, eggs and larva, Chlupaty; D. carneus, sound-production in aquarium, Koenig; D. trimaculatus ethology, Garnaud (2).

Pomacentrus albicaudatus sp. nov. Red Sea, Baschieri-Salvadori p. 64 figd.; P. bifasciatus, P. chrysopoecilus, P. chrysurus, P. moluccensis, P. perspicilatus, P. prosopotaenia new Bornean records, Inger; P. taeniurus new Red Sea record, Baschieri-Salvadori; P. xanthurus northward extension of range, Caldwell & Briggs.

LABRIDAE

Keys to Chinese genera and species, Fowler (1).

Key to genera of Western Indian Ocean, Smith, J. L. B. (12).

Anampses twistii recorded from Pemba and Mozambique, Smith, J. L. B. (12).

Aspiurochilus subgen. nov. of Lepidaplois q.v.

Choerodon robustus recorded from Seychelles, Smith, J. L. B. (12).

Cirrhilabrus exquisitus sp. nov. Pinda, W. Indian Ocean, Smith, J. L. B. (12) p. 109 figd.

Clepticus, association with Thalassoma, Eibl-Eibesfeldt.

Coris key to W. Indian Ocean species, Smith, J. L. B. (13); C. frerei recorded from E. Africa, Smith, J. L. B. (12); C. gaimard africana subsp. nov. Seychelles, Smith, J. L. B. (13) p. 119 figd.; C. grenovii Bennett juvenile form of C. gaimard, Smith, J. L. B. (12); C. julis description of eggs, Thomopoulos (1); sex reversal in old females, Reinboth; C. multicolor Rüppell young stages of C. caudimacula, Smith, J. L. B. (12); C. philippina recorded from Riu-Kiu Islands, Kamohara (3).

Cossyphus unilineatus Guichenot probably synonymous with Labrichthys cyanotaenia Bleeker, Smith, J. L. B. (12).

Crenilabrus pavo skin carotenoids, Abolins & Abolins-Krogis.

Diproctacanthus xanthurus new Bornean record, Inger.

Duohemipteronotus subgen. nov. of Hemipteronotus q.v.

Duymaeria flagellifera new Bornean record, Inger.

Fowlerella gen. nov. type Labroides bicolor Fowler & Bean, Smith, J. L. B. (12) p. 104.

Halichoeres amboinensis, H. gymnocephalus, H. leparensis, H. nigrescens new Bornean records, Inger; H. maculipinna, H. bivittata from Puerto Rico, Erdman; H. marginatus recorded from Mozambique, Smith, J. L. B. (12); H. notophthalmus recorded from E. Africa, Smith, J. L. B. (12).

Hemipteronotus (Duohemipteronotus) subgen. nov. type Hemipteronotus evides Jordan & Richardson, Fowler (1) p. 281.

Labroides dimidiatus new Bornean record, Inger.

Lepidaplois albomaculatus sp. nov. Mauritius, Smith, J. L. B. (12) p. 101 figd.; L. luteopunctatus sp. nov. Delagoa Bay, Smith, J. L. B. (12) p. 102 figd.; L. (Aspiurochilus) subgen. nov. type Crenilabrus steinegeri Ishikawa, Fowler (1) p. 176.

Macropharyngodon bipartitus p. 104 figd., M. varialvus p. 105 figd., spp. nov. Mozambique, Smith, J. L. B. (12); M. meleagris recorded from Delagoa Bay, Smith, J. L. B. (12).

Pseudocheilinus evanidus recorded from Aldabra, Smith, J. L. B. (12).

Stethojulis albovittata, S. axillaris, S. renardi, S. strigiventer recorded from Natal, Smith, J. L. B. (12); S. kalosoma, S. phekadopleura, S. trilineata new Bornean records, Inger.

Suezia gen. nov. type Labrichthys caudovittatus Steindachner, Smith, J. L. B. (12) p. 106.

Thalassoma see Clepticus; T. hardwicki new Bornean record, Inger; T. quinquevittatus recorded from S. Africa, Smith, J. L. B. (12); T. schwanefeldi, T. albolineum recorded from Riu-Kiu Islands, Kamohara (3).

Wetmorella albofasciata recorded from Aldabra, Smith, J. L. B. (12); W. philippina recorded from Seychelles and Mozambique, Smith, J. L. B. (12); W. philippina bifasciata Schultz & Marshall not valid, Smith, J. L. B. (12).

Xiphocheilus quadrimaculatus recorded from Queensland, Marshall, T. C.

SCARIDAE

Keys to genera and species of China, Fowler (1).

Callyodon dubius, C. lazulinus, recorded from W. Indian Ocean, Smith, J. L. B. (2); C. formosus new Australian record, Marshall, T. C.; C. pectoralis recorded from Riu-Kiu Islands, Kamohara (3); C. scaber, C. madagascariensis, C. improvisus recorded from Aldabra, Smith, J. L. B. (2).

Chlorurus microcheilus recorded from W. Indian Ocean, Smith, J. L. B. (2); C. microrhinus recorded from Riu-Kiu Islands, Kamohara (3).

Pseudoscarus coelestinus from Puerto Rico, Erdman.

Scarus post larvae descr. and figd. from Luzon, Blanco & Villadolid; S. acutus from Puerto Rico, Erdman.

Sparisoma radians from Puerto Rico, Erdman; recorded from Corpus Christi Bay, Texas, Leary (1).

Xanothon bataviensis recorded from Riu-Kiu Islands, Kamohara (3); X. capistratoides, X. pentazona, X. oktodon recorded from W. Indian Ocean, Smith, J. L. B. (2); X. carifanus, X. parvidens recorded from Aldabra, Smith, J. L. B. (2).

AMMODYTIDAE

Anmodytes †pseudoacuminatus otolith figd. from London Clay of Bognor, Stinton (1).

Bleekeria renniei sp. nov. S. Africa, Smith, J. L. B. (14) p. 219 figd.

OPISTHOGNATHIDAE

Opisthognathus maxillosus from Puerto Rico, Erdman; O. rosenbergi new Bornean record, Inger; O. whitehursti, O. maxillosus oral incubation, Böhlke & Chaplin.

PINGUIPEDIDAE

Parapercis key to species of China, Fowler (1).

TRACHINIDAE

Trachinus draco from Co. Waterford, Ireland, Gibson (2).

LIMNICHTHYIDAE

Chalixodytes chameleontoculis sp. nov. Seychelles, Smith, J. L. B. (4) p. 890 figd.

URANOSCOPIDAE

Key to genera and species of China, Fowler (1).

Ichthyscopus fasciatus sp. nov. North Queensland, Haysom p. 139 figd.

CHIASMODONTIDAE

Chiasmodon niger niger first record from Greenland waters, Kotthaus & Krefft.

Pseudoscopelus scriptus from Puerto Rico, Erdman.

NOTOTHENHDAE

Notothenia †tenuis sp. nov. otolith Miocene New Zealand, Stinton (2) p. 516 pl. xxxii fig. 14.

ACANTHUROIDEA

ACANTHURIDAE

Natural hybrids, Slastenenko.

Acanthurus chirurgus, A. coeruleus extension of range in Gulf of Mexico, Caldwell & Briggs; A. chrysosoma new Bornean record, Inger; A. randalli sp. nov. Gulf of Mexico, Briggs & Caldwell p. 43 figd.

Rhinodactylus baixopindae gen. et sp. nov. Mozambique, Smith, J. L. B. (1) p. 686 figd.

SIGANOIDEA

SIGANIDAE

Siganus oramin, S. stellatus, S. corallinus recorded from Aldabra, Smith, J. L. B. (2).

TRICHIUROIDEA

GEMPYLIDAE

Notes on larval forms from Bermuda, **Voss**, N. A.

Gempylus serpens from Puerto Rico, Erdman.

TRICHIURIDAE

Evoxymetopon taeniatus notes on a specimen from Gulf of Mexico, Tucker.

SCOMBROIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

SCOMBRIDAE

Natural hybrids, Slastenenko.

Summary of research in progress, Voss, G. L. & Robins.

Acanthocybium solandri from Alabama coast, Boschung; recorded from Aldabra, Smith, J. L. B. (2).

Auxis thazard first record for Gulf of Maine, Mather & Gibbs; recorded from Aldabra, Smith, J. L. B. (2).

Euthynnus alleteratus, E. pelamis, recorded from Aldabra, Smith, J. L. B. (2).

Gasterochisma melampus recorded from Victoria, N.S.W., Whitley (2).

Germo albacora recorded from Aldabra, Smith, J. L. B. (2).

Gymnosarda unicolor from W. Indian Ocean, Smith, J. L. B. (2).

Neothunnus macropterus spawning areas and size at sexual maturity, Yuen & June; N. macropterus, Parathunnus sibi comparative food study, King & Ikehara.

Parathunnus mebachi first Australian record, Munro (2).

Rastrelliger kanagurta feeding habits' Kuthalingam; recorded from Aldabra, Smith, J. L. B. (2).

Sarda orientalis recorded from Aldabra, Smith, J. L. B. (2).

Scomber scombrus population study, Francesco.

Scomberomorus cavalla from Buzzards Bay, Mass., Mather & Gibbs; S. commerson, S. guttatus recorded from Aldabra, Smith, J. L. B. (2).

Thunnus alalunga taken off Delaware Bay, Mather & Gibbs; T. thynnus of E. and W. Atlantic alike in meristic characters, T. saliens distinct, Robins (2); biometric study, Genovese; recorded from Puerto Rico, Erdman; structure of pancreas, spleen, liver and intestine, Vallmitjana & Gadea; study on North Sea population, Tiews.

LUVARIDAE

Luvarus imperialis recorded from Chilean waters, Buen (3).

ISTIOPHORIDAE

Research in progress, Voss, G. L. & Robins.

Istiophorus americanus from Puerto Rico, Erdman; age and growth, de Sylva; study of a developmental series, spawning areas, food, Gehringer.

Makaira ampla, M. albida from Puerto Rico, Erdman.

XIPHIIDAE

Research in progress, Voss, G. L. & Robins.

†Palaeorhynchus glaronensis figd., Smigielska.

Xiphias gladius length/weight relationship; parasites, Postel (1).

GOBIOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

ELEOTRIDAE

Asterropteryx semipunctatus recorded from Aldabra, Smith, J. L. B. (3).

Electriodes longipinnis Japanese examples discussed, Tomiyama & Abe; E. strigatus, E. sexguttatus, recorded from W. Indian Ocean, Smith, J. L. B. (3).

Eleotris fusca recorded from Aldabra, Smith, J. L. B. (3).

Eviota post larvae descr. and figd. from Luzon, Blanco & Villadolid; E. distigma new Bornean record, Inger; E. gymnocephalus Weber a synonym of E. prasinus (Klunzinger), Smith, J. L. B. (3); E. zonura, E. prasinus, E. prasites recorded from Aldabra, Smith, J. L. B. (3).

Eviotops gen. nov. p. 825 type E. infulatus sp. nov. p. 826 figd., Mahé, Smith, J. L. B. (3).

Kribia revision of genus, Lambert.

Metagobius sclateri recorded from W. Indian Ocean, Smith, J. L. B. (3).

Platycephalops gen. nov. type Electris polyzonatus Klunzinger, Smith, J. L. B. (3) p. 822 figd.

Pogonoculius zebra recorded from Indian Ocean, Smith, J. L. B. (3).

Pterelectris microlepis recorded from Aldabra, Smith, J. L. B. (3); P. tricolor sp. nov. Mozambique, Smith, J. L. B. (3) p. 817 figd.

Trimma naudei sp. nov. Mahé, Smith, J. L. B. (3) p. 828 figd.

GOBIIDAE

Taxonomic notes, Ilyin (1).

Sensory papillae, Satô.

Acanthogobius flavimanus feeding habits, Imamura & Hashitani.

Acentrogobius ornatus, A. reichei new Bornean records, Inger.

Amblygobius albimaculatus recorded from Aldabra, Smith, J. L. B. (2); A. phalaena new Bornean record, Inger.

Barbulifer cauthoecus from Puerto Rico, Erdman.

Bathygobius fuscus new Bornean record, Inger; recorded from Aldabra, Smith, J. L. B. (2).

Boleophthalmus caeruleomaculatus new Queensland record, Marshall, T. C.

Callyogobius hasseltii from W. Indian Ocean, Smith, J. L. B. (2).

Chonophorus ocellaris post larvae described and figd. from Luzon, Blanco & Villadolid.

Cryptocentrus octofasciatus recorded from Aldabra, Smith, J. L. B. (2).

Ctenogobius nocturnus sp. nov. Aldabra, Smith, J. L. B. (2) p. 723 figd.

Elacatinus oceanops, association with Epinephelus, Eibl-Eibesfeldt.

Erotelis smaragdus from Puerto Rico, Erdman.

Favonigobius exquisitus new Queensland record, Marshall, T. C.

Fusigobius neophytus recorded from W. Indian Ocean, Smith, J. L. B. (2).

Gladiogobius ensifer recorded from W. Indian Ocean, Smith, J. L. B. (2).

Glossogobius key to Indo-Australian species, Inger; G. giuris recorded from Aldabra, Smith, J. L. B. (2); G. sandakanensis sp. nov. N. Borneo, Inger p. 393 figd.

Gnatholepis thompsoni from Puerto Rico, Erdman.

Gobiodon ceramensis new Queensland record, Fowler (2); G. citrinus recorded from Aldabra, Smith, J. L. B. (2); G. citrinus, G. erythrospilus new Bornean records, Inger.

Gobionellus claytoni from Puerto Rico, Erdman.

Gobiosoma novemlineatum, G. oceanops, G. horsti, from Puerto Rico, Erdman.

Gobius dorsorostralis sp. nov. p. 38 pl. vi figs. 49, 50 and G. d. sculpta subsp. nov. otoliths Pannonian Vienna Basin, Papp & Thenius; G. ebriosus from Puerto Rico, Erdman; G. lidwilli, early development, Dôtu (2); G. (Macrogobius) paganellus occurrence in Black Sea, Ilyin (2); G. syrman, G. fluviatilis biological rôle in Razelm waters, Roumania, Banarescu (2).

Lophogobius cyprinoides from Puerto Rico, Erdman,

Luciogobius elongatus figd., Dôtu (1); L. guttatus bionomics, Dôtu (3); L. pallidus Regan a synonym of L. albus Regan, Dôtu (1); L. saikaiensis sp. nov. Kyushu, Dôtu (1) p. 71 figd.

Neogobius melanostomus, N. syrman gas exchange, Shulman & others; cutaneous respiration, Shulman.

Oxyurichthys microlepis new Australian record, Marshall, T. C.; O. tentacularis new Japanese record, Kamohara (2).

Paragobiodon echinocephalus recorded from Aldabra, Smith, J. L. B. (2).

Periophthalmus cantonensis recorded from Aldabra, Smith, J. L. B. (2).

Proterorhinus marmoratus recorded from Austrian waters, Bauer & Schubert; occurrence in Neusiedler Lake, Austria, Sochurek.

Pseudogobius javanicus new Bornean record, Inger.

Rhinogobius mowbrayi from Puerto Rico, Erdman.

Seychellea Smith, J. L. B. a synonym of Koumansetta Whitley, Whitley (2); S. hectori gen. et sp. nov. Mahé, E. Africa, Smith, J. L. B. (2) p. 726 figd.

Triaenopogon barbatus bionomics, Dôtu (5).

Vaimosa piapensis post larvae descr. and figd. from Luzon, Blanco & Villadolid.

GOBIOIDIDAE

Gobioides brussonnettii from Alabama coast, Boschung.

Tyntlastes brevis taken in Gulf of Panama, Clemens.

TAENIOIDIDAE

Odontamblyopus rubicundus bionomics, Dôtu (4).

CALLIONYMOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

CALLIONYMIDAE

Callionymus festivus from Algerian coast, Dieuzeide & Roland (2); C. garthi Seale a synonym of C. atrilabiatus Garman; extension of range, Böhlke & Cohen.

Synchiropus altivelis taxonomy and bionomics, Azazaki; S. ocellatus new Bornean record, Inger.

BLENNIOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

Vol. 94

BLENNIIDAE

Key to Australian genera related to Tripterygion, Scott, T. D.

Acanthemblemaria key to Atlantic species, Böhlke (2); A. chaplini sp. nov. Bahamas, Böhlke (2) p. 37 figd.; A. spinosa, A. aspera, A. variegata new Bahaman records, Böhlke (2).

Alticops periophthalmus, A. edentulus recorded from Aldabra, Smith, J. L. B. (4).

Aspidontus taeniatus, A. filamentosus recorded from Aldabra, Smith, J. L. B. (4).

Atrosalarias fuscus new Bornean record, Inger.

Blennius adaptation to environment, Lecal; B. fluviatilis description of larvae, Wickler (2); B. pholis bionomics, Qasim.

Brachynectes gen. nov. p. 180 type B. fasciatus sp. nov. p. 181 figd., S. Australia, Scott, T. D.

Chaenopsis key to species, Böhlke (4); C. alepidota californiensis subsp. nov. California, Böhlke (4) p. 99 figd.; C. coheni Gulf of California p. 90 figd., C. deltarrhis Gorgona Is., Colombia p. 93, C. schmitti Galapagos Archipelago p. 94 spp. nov., Böhlke (4).

Cirripectus leopardus recorded from Aldabra, Smith, J. L. B. (4).

Coralliozetus Evermann and Marsh a synonym of Emblemaria Jordan and Gilbert, Böhlke (2).

Emblemaria, Hemiemblemaria key to Bahaman species, Böhlke (2); E. angelica sp. nov. Baja California, Böhlke & Mead p. 2 figd.; E. cardonae, E. diaphana new Bahaman records, Böhlke (2); E. marki Mowbray a synonym of E. atlantica Jordan and Evermann, Böhlke (2); E. pandionis new record from British Honduras, Böhlke (2).

Enchelyurus kraussi new Australian record, Haysom; recorded from W. Indian Ocean, Smith, J. L. B. (4).

Entomacrodus striatus, E. vermiculatus recorded from Aldabra, Smith, J. L. B. (4).

Gunellichthys pleurotaenia new Bornean record, Inger.

Halmablennius fasciatus recorded from Aldabra, Smith, J. L. B. (4); H. lineatus recorded from W. Indian Ocean, Smith, J. L. B. (4).

82

Hemiemblemaria simulus new Bahaman record, Böhlke (2).

Lophalticus gen. nov. type Salarias kirkii (Gunther), Smith, J. L. B. (4) p. 889.

Lucaya gen. nov. p. 83 type L. zingaro sp. nov. Green Cay, Bahamas, p. 84 figd., Böhlke (4).

Omobranchus kallosoma recorded from Aldabra, Smith, J. L. B. (4).

Parablennius Ribeiro, validity, Pinto (2).

Paremblemaria erythrops Fowler a synonym of Acanthemblemaria aspera (Longley), Böhlke (2).

Petroscirtes mitratus recorded from Aldabra, Smith, J. L. B. (4); P. rhynorhynchus new Bornean record, Inger.

Psednoblennius Jenkins and Evermann a synonym of Emblemaria Jordan and Gilbert, Böhlke (2).

Salarias chiliensis Clark a synonym of Homesthes biocellatus, Cohen; S. cyanostigma, S. fasciatus, S. guttatus new Bornean records, Inger; S. gibbifrons recorded from W. Indian Ocean, Smith, J. L. B. (4); S. petersoni Fowler a synonym of Scartichthys gigas, Cohen.

Scartichthys eques Steindachner a synonym of S. gigas, Cohen.

ANARRHICHADIDAE

Anarhichas lupus from Long Island Sound, Morrow (1); increase in numbers, Barsukov (1); A. lupus lupus, A. l. maris-albi compared, Barsukov (2); A. lupus, A. minor, A. latifrons distribution in Barents Sea, Barsukov (3).

CLINIDAE

Labrisomus trindadensis sp. nov. Trindade Isl., Brazil, Pinto (1) p. 1 figd.

Malacoctenus key to Puerto Rican species, Smith, C. L.; M. aurolineatus p. 2 figd., M. erdmani p. 8 figd. spp. nov. Puerto Rico, Smith, C. L.

Petraites roseus new Queensland record, Marshall, T. C.

STICHAEIDAE

Stichaeus grigorjewi structure of pituitary, Honma (1).

PHOLIDIDAE

Pholis gunnellus northern extension of range, Gordon & Backus.

LUMPENIDAE

Lumpenus fabricii first record from Labrador, Gordon & Backus; L. lumpretaeformis northern extension of range, Gordon & Backus.

ZOARCIDAE

Lycodonus dorsoscutatus sp. nov. Kamo, Japan, Oshima (1) p. 7 figs.

Lycogramma japonica sp. nov. Japan, Oshima (1) p. 7.

OPHIDIOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

BROTULIDAE

Barathrites key to three N. Atlantic species, Nybelin; B. parri sp. nov. N. Atlantic, Nybelin p. 283 figd.

Barathronus erikssoni sp. nov. North Atlantic, Nybelin p. 308 figd.

Bassogigas (Holcomycteronus) koefoedi Nybelin a synonym of Grimaldichthys squamosus, Nybelin.

Brosmophycis marginata Californian records, Best (1).

Brotula barbata from Puerto Rico, Erdman; B. multibarbata recorded from Aldabra, Smith, J. L. B. (4).

Dinematichthys cayorum from Puerto Rico, Erdmann; D. iluocoeteoides new Bornean record, Inger; recorded from W. Indian Ocean, Smith, J. L. B. (4).

Grimaldichthys a synonym of Bassogigas, Nybelin.

Holcomycteronus a synonym of Bassogigas, Nybelin.

Lamprogrammus key to species, Nybelin.

Leucochlamys jonassoni sp. nov. North Atlantic, Nybelin p. 311 figd.

Parabassogigas gen. nov. type Bassogigas grandis (Gunther), Nybelin p. 298.

Petrotyx sanguineus from Puerto Rico, Erdman.

Porogadus abyssalis sp. nov. N. Atlantic, Nybelin p. 288 figd.

Sciadonus kullenbergi sp. nov. North Atlantic, Nybelin p. 310 figd. Stygnobrotula gen. nov. p. 1 type S. latebricola sp. nov. Green Cay, Bahamas, p. 2 figd., Böhlke (1).

OPHIDIIDAE

Lepophidium profundarum from Puerto Rico, Erdman.

Otophidium grayi a valid species, Robins (1); O. welshi taken off Massachusetts, Mather & Gibbs; from Atlantic coast of Georgia, Caldwell & Briggs.

†Otolithus (Ophidiidarum) productus sp. nov. London Clay Bognor p. 27 pl. ii fig. 19; Ot. (O.) sheppeyensis, Ot. (O.) symmetricus figd., Stinton (1).

CARAPIDAE

Carapus acus, C. cuspis, Echiodon dentatus nomenclature, Arnold.

Jordanicus †exiguus sp. nov. otolith figd. Miocene New Zealand, Stinton (2) p. 516 pl. xxxii fig. 9; J. gracilis recorded from Aldabra, Smith, J. L. B. (4).

Onuxodon parvibrachium recorded from Aldabra, Smith, J. L. B. (4).

STROMATEOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

Nomeidae

Cubiceps gracilis juvenile recorded from Bonin Isl., Abe (2); C. ismaelensis from Algerian coast, Dieuzeide & Roland (2).

Palinurichthys bythites extension of depth range, Caldwell & Briggs.

Psenes cyanophrys bionomics, Legaspi; from Puerto Rico, Erdman; P. indicus recorded from Aldabra, Smith, J. L. B. (2).

TETRAGONURIDAE

Tetragonurus cuvieri juvenile taken off Bonin Isl., Abe (2); 146 specimens taken south of Aleutian Isl., Tanonaka.

ANABANTOIDEA

ANABANTIDAE

Histology of endocrine organs detailed study of reproductive and related behaviour, Forselius.

Colisa labiosa carcinoma of skin, Stolk (4).

Helostoma temmincki, main, accessory and emergency sources of food, Vaas.

Macropodus opercularis, colour-photo, breeding, Pinter.

Trichogaster trichopterus siamensis, Klausewitz (9); T. trichopterus, "T. t. sumatranus" not a subspecies but a colour-variety, Ladiges.

OPHICEPHALOIDEA

OPHICEPHALIDAE

Structure of scales, Bhatti & Aminul Huq.

Ophicephalus introduction in Japan, Okada; anatomy of swim bladder, George; O. punctatus afferent branchials, Bhuiya & Siddiqui.

PHALLOSTETHOIDEA

PHALLOSTETHIDAE

Neostethus post larvae descr. and figd. from Luzon, Blanco & Villadolid.

MUGILOIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

SPHYRAENIDAE

Sphyraena akerstromi recorded from Queensland, Marshall, T. C.; S. barracuda, S. jello, S. flavicauda, S. toxeuma recorded from Aldabra, Smith, J. L. B. (2); S. borealis adult taken off New England, Mather & Gibbs; S. aff. †malembeensis figd. from Miocene Îsle du Prince, Henriques da Silva.

MUGILIDAE

Key to Adriatic species; age and growth study, Morovic.

Feeding habits; dentition, with special reference to taxonomy, Ebeling.

Agonostomus monticola in Louisiana, Suttkus (2).

Crenimugil crenilabis recorded from Aldabra, Smith, J. L. B. (2).

Liza macrolepis recorded from Aldabra, Smith, J. L. B. (2),

Mugil auratus now introduced into Caspian Sea, Zenkevitch; M. cephalus from Oklahoma and northern Texas, Riggs; observations on schooling, Springer, S. (2); M. diadema new

Queensland record, Marshall, T. C.; M. †latus sp. nov. L. Miocene Caucasus, Svichenskaya p. 61; M. saliens, M. auratus, M. cephalus phylogenetic relationship, Burdak; M. †voesendorfensis sp. nov. otoliths Pannonian Vienna Basin, Papp & Thenius p. 34 pl. vi figs. 37, 38.

Plicomugil labiosus first W. Indian Ocean record, Smith, J. L. B. (2).

Valamugil buchanani, V. vaigiensis, recorded from Aldabra, Smith, J. L. B. (2).

ATHERINIDAE

Natural hybrids, Slastenenko.

Atherina presbyter Dutch coast, Kristensen.

Basilichthys bonariensis structure of alimentary canal, Carlo & Lopez.

Hepsetia pinguis recorded from Aldabra, Smith, J. L. B. (2).

POLYNEMOIDEA

POLYNEMIDAE

Eleutheronema tetradactylum chondrocranium, Marathe & Bal.

Polydactylus plebeius recorded from Aldabra, Smith, J. L. B. (2).

SCLEROPAREI

Larval and juvenile forms from the Mediterranean, Padoa & others.

SCORPAENOIDEA

SCORPAENIDAE

Key to genera and species of W. Indian Ocean, Smith, J. L. B. (9) & (10).

Amblyapistus taenianotus new Bornean record, Inger.

Dendrochirus, Pterois key to species, Klausewitz (6).

Dendroscorpaena gen. nov. type Scorpaena cirrhosa (Thunberg), Smith, J. L. B. (9) p. 60; D. gilchristi sp. nov. off the Tugela river, Smith, J. L. B. (9) p. 62 figd.

Helicolenus lahillei Norman a synonym of H. maculatus Cuvier, Smith, J. L. B. (10).

Parabrachirus Matsubara a synonym of Parapterois Bleeker, Smith, J. L. B. (10).

Parascorpaena maculipinnis sp. nov. Bazaruto, Indian Ocean, Smith, J. L. B. (9) p. 57 figd.

Parascorpaenodes hirsutus gen. et sp. nov. Pinda, Smith, J. L. B. (9) p. 63 figd.

Pterois lunulata Schlegel a synonym of P. russellii Bennett, Smith, J. L. B. (10); P. miles Bennett a synonym of P. volitans, Smith, J. L. B. (10).

Pteropterus mombasae sp. nov. Mombasa, Smith, J. L. B. (10) p. 80 figd.

Scorpaenodes caribbaeus new Florida record, Robins (1); S. corallinus Baixo Pinda, W. Indian Ocean p. 64 figd., S. varipinnis Zanzibar p. 65 figd., spp. nov., Smith, J. L. B. (9); S. guamensis new Bornean record, Inger.

Scorpaenopsis gibbosus new Bornean record, Inger.

Sebastes distribution, Baranenkova; food of in S.E. Greenland area, Magnússon (3); S. marinus description of larvae from Baie-des-Chaleurs, Jean; structure of urogenital system and reproductive cycle in ♂ and ♀; deposition of spermatozoa within ovary, Magnusson (1); structure of retina, Wunder; S. stoliczkae new Bornean record, Inger.

TRIGLIDAE

Paraheminodus gen. nov. type Satyrichthys laticephalus Kamohara, Kamohara (2) p. 4.

Peristethion cataphractum description of eggs, Thomopoulos (1).

Prionotus punctatus northern extension of range, Caldwell & Briggs.

SYNANCEJIDAE

Synanceja verrucosa account of two fatalities from stabs, Smith, J. L. B. (8); description of venom apparatus and its effects, Gail & Rageau.

PLATYCEPHALIDAE

Thysanophrys cirronasus new Queensland record, Marshall, T. C.

COTTIDAE

Cottopsis parvus Girard synonymous with Cottus asper, Robins & Miller.

Cottus key to Californian species, Robins & Miller; C. aleuticus aggregational behaviour, Greenbank; C. bubalis reproduction and embryonic development, Thomopoulos & Bauchot; C. gobio bionomies, Smyly; C. macrops Rutter a synonym of C. klamanthensis, Robins & Miller; C. perplexus Gilbert & Evermann a valid species, Robins & Miller; C. poecilopus in Finland, Koli; C. protrusus Schultz & Spoor a synonym of C. aleuticus, Robins & Miller; C. shasta synonymous with C. gulosus, Robins & Miller.

Enophrys diceraus recorded from arctic Alaskan waters, Mohr, Wilimovsky & Dawson.

Scorpaenichthys marmoratus avoidance of eggs by predators, Pillsbury.

Zesticelus japonicus sp. nov. Niigata, Japan, Oshima (1) p. 2 figd.

COMEPHORIDAE

Ecological adaptations in reproductive behaviour, Koryakov.

CYCLOPTERIDAE

Cyclopteropsis inarmatus sp. nov. Bering Sea, Mednikov & Prokhorov p. 717 figd.

Lethotremus fuscopunctatus sp. nov. Japan, Oshima (1) p. 4 figd.

LIPARIDAE

Crystallias kamoae sp. nov. Kamo, Japan, Oshima (1) p. 5 figd.

Liparis frenatus recorded from Nabuto, Japan, Kuroda (3).

DACTYLOPTEROIDEA

Larval and juvenile forms from the Mediterranean, Padoa & others.

THORACOSTEI

GASTEROSTEIDAE

Gasterosteus aculeatus breeding colouration of female, Drapkin; G. aculeatus × Pygosteus pungitius hybrids, Leiner.

HETEROSOMATA

Importance of lateral body orientation, Aleev (2).

Larval and juvenile forms from the Mediterranean, Padoa & others.

BOTHIDAE

Natural hybrids, Slastenenko.

Bothus pantherinus new Bornean record, Inger.

PLEURONECTIDAE

Natural hybrids, Slastenenko.

Clidoderma asperrimum new record from British Columbia, Welander, Alverson & Bergman.

Eopsetta jordani movements of spawning population, Alverson, D. L. & Chatwin.

Hippoglossoides platessoides breeding and fecundity, Bagenal (1).

Pleuronectes platessa structure of asymmetrical head, Blok; German Bight population study, Kotthaus; food of larvae, Shelbourne.

Reinhardtius hippoglossoides first record from Labrador, Gordon & Backus.

SOLEIDAE

Gymnachirus williamsoni recorded from Atlantic; compared with G. texae, Caldwell & Briggs.

Pardachirus pavoninus new Bornean record, Inger.

Solea humilis post-larvae descr. and figd. from Luzon, Blanco & Villadolid; S. lascaris description of eggs, Thomopoulos (1).

Synaptura selheimi from North Queensland, Marshall, T. C.

DISCOCEPHALI

Larval and juvenile forms from the Mediterranean, Padoa & others.

ECHENEIDAE

Echeneis naucrates recorded from Aldabra, Smith, J. L. B. (4).

Remora albescens recorded from Aldabra, Smith, J. L. B. (4); R. remora notes on respiration, Strasburg; R. remora, Rhombochirus osteochir from Puerto Rico, Erdman.

Remoropsis pallidus new Australian record, Haysom.

Rhombochirus osteochir new Australian record, Marshall, T. C.

PLECTOGNATHI

Larval and juvenile forms from Mediterranean, Tortonese in Padoa & others.

BALISTOIDEA

BALISTIDAE

Balistapus verrucosus new Bornean record, Inger.

Balistes capriscus from Puerto Rico, Erdman; skull, Garnaud (1).

Canthidermis sobaco from Puerto Rico, Erdman.

MONACANTHIDAE

Alutera schoepfi, A. punctata from Puerto Rico, Erdman.

Amanses pardalis recorded from Nabuto, Japan, Kuroda (3).

Cantherines pardalis new Bornean record, Inger.

Monacanthus cirrhifer effect of testosterone on dorsal fin, Ishii & Egami; M. tuckeri recorded from Florida, Robins (1).

Pervagor scanleni sp. nov. S. Africa, Smith, J. L. B. (14) p. 221 figd.

OSTRACIONTIDAE

Rhinesomus gibbosus new Australian record, Marshall, T. C.

TETRAODONTOIDEA

TETRAODONTIDAE

Arothron nigropunctatus new Bornean record, Inger.

Canthigaster rostratus recorded from Florida, Robins (1).

Carinotetraodon chlupatyi gen. et sp. nov. Thailand, Benl (2) p. 1 figd.

Sphaeroides halsteadi sp. nov. N.S.W., Whitley (2) p. 70 figd.

Tetraodon leiurus brevirostris subsp. nov. ? Siam, Benl (1) p. 63 figd.; T. ocellaris sp. nov. Thailand, Klausewitz (9) p. 201 figd.; T. somphongsi sp. nov. Thailand, Klausewitz (10) p. 205 figd., Klausewitz (3).

DIODONTIDAE

Chilomycterus antillarum recorded from Florida, Robins (1).

Diodon bleekeri post larvae descr. and figd. from Luzon, Blanco & Villadolid; D. holacanthus new Bornean record, Inger; D. hystrix spawning in captivity, Wolfsheimer.

Lophodiodon nigropunctatus sp. nov. S. Africa, Smith, J. L. B. (14) p. 222 figd.

MOLIDAE

Mola mola, Aramburu (2); occurrence in Texas waters, Kemp; M. ramsayi description of Chilean specimens, Buen (2).

XENOPTERYGII

Larval and juvenile forms from the Mediterranean, Padoa & others.

GOBIESOCIDAE

Key to genera of Lepadogastrinae, Briggs.

Acyrtops beryllinus from Puerto Rico, Erdman.

Arcos macrophthalmus from Puerto Rico, Erdman.

Gobiesox nigripinnis, G. strumosus from Puerto Rico, Erdman.

Lecanogaster gen. nov. p. 204 type L. chrysea sp. nov. Gold Coast, p. 205 figd., Briggs.

Lepadogaster l. lepadogaster, L. l. purpurea geographical range, Briggs.

Opeatogenys cadenati sp. nov. Senegal' French W. Africa, Briggs p. 206 figd.

Sicyases sanguineus results of electrocardiogram, Concha & Vargas; structure of adrenals, Vargas & Concha (1).

Tomicodon fasciatus from Puerto Rico, Erdman.

HAPLODOCI

BATRACHOIDIDAE

Porichthys myriaster luminescence and innervation of photophores, Nicol.

PEDICULATI

Larval and juvenile forms from the Mediterranean, Padoa & others.

ANTENNARIOIDEA

ANTENNARIIDAE

Key to genera and species, Schultz (2).

Abantennarius gen. nov. type Antennarius duescus Snyder, Schultz (2) p. 66 figd.; A. analis sp. nov. Hawaii, Gosline in Schultz (2) p. 67 figd.

Antennarius (Plumantennatus) subgen. nov. type Antennarius asper Macleay, Schultz (2) p. 89 figd.; A. bermudensis p. 98 figd., Bermuda, A. pauciradiatus p. 100 figd., Florida, spp. nov., Schultz (2); A. fuliginosus sp. nov. S. Africa, Smith, J. L. B. (14) p. 222 figd.

Antennatus gen. et subgen. nov. type A. strigatus Gill, Schultz (2) p. 80 figd.; A. (Xenophrynichthys) subgen. nov. type Antennarius cryptacanthus Weber, Schultz (2) p. 81 figd.

Golem gen. nov. type Antennarius cryptacanthus Weber, Whitley (2) p. 70.

Kanazawaichthys gen. nov. p. 62 type K. scutatus sp. nov. p. 63 figd., Gulf of Mexico, Schultz (2).

Lophiocharon (Uniantennatus) subgen. nov. type Antennarius horridus Bleeker, Schultz (2) p. 83 figd.; L. tenebrosus new Florida record, Robins (1).

Nudiantennarius gen. nov. type Antennarius subteres Smith & Radcliffe, Schultz (2) p. 66 figd.

Phrynelox (Triantennatus) subgen. nov. p. 74 type. Antennarius zebrinus sp. nov. p. 75 figd. N.S.W., Schultz (2); P. (Triantennatus) atra sp. nov. N.S.W., Schultz (2) p. 76 figd.

Plumantennatus subgen. nov. of Antennarius q.v.

Triantennatus subgen. nov. of Phrynelox q.v.

Uniantennatus subgen. nov. of Lophiocharon q.v.

Xenophrynichthys subgen. nov. of Antennatus q.v.

OGCOCEPHALIDAE

Ogcocephalus parvus from Puerto Rico, Erdman.

CERATIOIDEA

ONEIRODIDAE

Oneirodes bradburyae sp. nov. Gulf of Mexico, Grey p. 245 figd.

CERATIIDAE

Cryptopsaras couesii from Monterey Bay, first Californian record, described and figd., **Fast**.

OPISTHOMI

MASTACEMBELIDAE

Mastacembelus armatus secretory cells of gill filaments, Kapoor, B. G. (1).

CROSSOPTERYGII

Derivation of first tetrapods, Jarvik.

†RHIPIDISTIA

Laccognathus, Glyptolepis and Holoptychius remains descr. and figd. from Escuminae Bay, Ørvig (1).

Litoptychus and Onychodus remains figd. and referred to Holoptychiidae, Ørvig (1).

Onychodus and Hamodus spiral teeth figd., Gross.

HOLOPTYCHIIDAE

Hamodus lutkevitschi figd., Obruchev (1).

Holoptychius sp. figd., Obruchev (1).

RHIZODONTIDAE

Eusthenopteron wenjukowi figd., Obruchev (1).

Strepsodus siberiacus seale figd., Khabakov (1).

OSTEOLEPIDAE

Porolepis scales descr. and figd. from Spitsbergen, Ørvig (1); P. uralensis figd., Obruchev (1).

ACTINISTIA

COELACANTHIDAE

Latimeria chalumnae comparison of known specimens, Schindler (1); nucleic acids, Vialli.

†Rhabdoderma? aegyptiaca L. Carboniferous Egypt, Heide p. 74 pl. xvii fig. 28.

DIPNOI

Evolution, Lehman (1).
Palaeogeographic importance, Pidoplichko & Kurazhskovski.

†DIPTERIDAE

Dipterus serratus figd., Obruchev (1).

LEPIDOSIRENIDAE

Protopterus bionomics, Chopard; comparative morphology of mid-brain, Dorn; P. dolloi reproduction, Brien, Poll & Bouillon.

CERATODONTIDAE

†Ceratodus minor p. 310 pl. i fig. 1; C. youngi p. 310 pl. i fig. 2, spp. nov. Jurassic Szechuan, China, Liu & Yeh.

Epiceratodus Teller a synonym of Neoceratodus Castelnau, Miller, R. R. (1).

Neoceratodus forsteri (Krefft) correct name for Australian lungfish, Miller, R. R. (1).

ERRATA

PISCES

Vol. 92.

p. 89. Under Gobiidae: Croila should read Croilia.

Vol. 93.

p. 16. Under Geiger, W. insert "27:" after "and" in the fifth line.

p. 22. Under Illick, H. J., last line: "36" should read "56".

p. 23. Under Jordan & Evermann Bull, U.S. Fish, Comm. 22: in fourth line delete "1904" and substitute "1902 (1903)".

p. 66. Under †Osteostraci. The first two entries should read: "Nature of electric or sensory organs, Bohlin; Cory.

Kolymaspis gen. nov. type K. sibirica sp. nov.;"

p. 67. Under Selachii. Cooperodon should be preceded by "†".